# The Coues Check List and ORNITHOLOGICAL DICTIONARY

# Smithsonian Institution *Libraries*




Alexander Wetmore
1946 Sixth Secretary 1953



9, 1, Miller Jr. Peterboro. Dec,

a. Wetmore A=1



QL 677 C85 1882 BIRDS

# COUES CHECK LIST

OF

# NORTH AMERICAN BIRDS.

SECOND EDITION,

Rebised to Date, and entirely Rewritten, under Direction of the Author,

WITH A DICTIONARY OF THE

ETYMOLOGY, ORTHOGRAPHY, AND ORTHOEPY

OF THE

#### SCIENTIFIC NAMES,

THE CONCORDANCE OF PREVIOUS LISTS, AND A CATALOGUE OF HIS ORNITHOLOGICAL PUBLICATIONS.



BOSTON:

ESTES AND LAURIAT.

1882.

Copyright, 1882, By Estes and Lauriat.

University Press:

John Wilson and Son, Cambridge.

### INTRODUCTION.

In 1873, shortly after the publication of the author's "Key to North American Birds," appeared the original edition of this "Check List," which was almost immediately reissued in connection with the same writer's "Field Ornithology," in 1874. That list reflected the classification and nomenclature of the "Key" with much exactitude, although it included, in an Appendix, a few species additional to those described in the "Key," and made some slight changes in the names. Excepting some little comment in foot-notes and in the Appendix, the original "Check List" was a bare catalogue of scientific and vernacular names, printed in thick type on one side of the paper.

Meanwhile, the science of Ornithology has progressed, and our knowledge of North American birds has increased, both in extent and in precision, until the original list, faithful as it was at the time, fails now to answer the purpose of adequately reflecting the degree of perfection to which the subject has been brought. A new edition has therefore become necessary.

The list has been revised with the utmost care. The gratifying degree of accuracy with which it represented our knowledge of 1873 is exhibited in the fact, that it is found necessary to remove no more than ten names. On the other hand, the progress of investigation has resulted in adding one hundred and twenty names to the list, and in showing the necessity or expediency of making many changes in nomenclature. The exact analysis of the differences between the two lists is given beyond.

In revising the list for the main purpose of determining the ornithological status of every North American bird, the most scrupulous attention has been paid to the matter of nomenclature, — not only as a part of scientific classification, determining the technical relations of genera, species, and varieties to each other, but also as involved in writing and speaking the names of birds correctly. The more closely this matter was scrutinized, the more evidences of inconsistency, negligence,

or ignorance were discovered in our habitual use of names. It was therefore determined to submit the current catalogue of North American birds to a rigid examination, with reference to the spelling, pronunciation, and derivation of every name, — in short, to revise the list from a philological as well as an ornithological standpoint.

The present "Check List," therefore, differs from the original edition in so far as, instead of being a bare catalogue of names, it consists in a treatise on the etymology, orthography, and orthoepy of all the scientific, and many of the vernacular, words employed in the nomenclature of North American birds. Nothing of the sort has been done before, to the same extent at any rate; and it is confidently expected that the information given here will prove useful to many who, however familiar they may be with the appearance of these names on paper, have comparatively little notion of the derivation, signification, and application of the words; and who unwittingly speak them as they usually hear them pronounced, that is to say, with glaring impropriety. No one who adds a degree of classical proficiency to his scientific acquirements, be the latter never so extensive, can fail to handle the tools of thought with an ease and precision so greatly enhanced, that the merit of ornithological exactitude may be adorned with the charm of scholarly elegance.

The purpose of the present "Check List" is thus distinctly seen to be twofold: First, to present a complete list of the birds now known to inhabit North America, north of Mexico, and including Greenland, to classify them systematically, and to name them conformably with current rules of nomenclature; these being ornithological matters of science. Secondly, to take each word occurring in such technical usage, explain its derivation, significance, and application, spell it correctly, and indicate its pronunciation with the usual discritical marks; these being purely philological matters, affecting not the scientific status of any bird, but the classical questions involved in its name.

In the latter portion of his task, which, as is always the case when thorough work of any kind is undertaken, proved to be more difficult and more protracted than had been expected, and delayed the appearance of the list for nearly a year after the ornithological portion had been practically completed, the author of the original list has received invaluable assistance from Mrs. S. Olivia Weston-Aiken, who cordially shared with him the labor of the philological investigation, and to whose scholarly attainments he is so largely indebted, that it is no less a duty than a pleasure to recognize the co-operation of this accomplished lady.

#### ANALYSIS OF THE TWO EDITIONS.

The original edition of the "Check List" ostensibly enumerates only 635 species of North American Birds. This is owing to the fact that only full species are numbered, the many subspecies being given as a, b, &c., and some names being interpolated without corresponding numbers, both in the body of the list and in the Appendix. By actual count there are found to be, in the body of the list, 750; to which 28 are added in the Appendix: 750 + 28 = 778.

First, with regard to subtractions. It is in gratifying evidence of the general accuracy of the original list, that it is found necessary to remove only ten (10) names. Four of these are extra-limital; six are mere synonyms. The following is the—

#### LIST OF SUBTRAHEND NAMES.

- 1. Ægiothus fuscescens. Summer plumage of Æ. linaria.
- 2. Centronyx ochrocephalus. Fall plumage of Passerculus bairdi.
- 3. Sphyropicus williamsoni. Male of S. thyroïdes.
- 4. Lampornis mango. Extra-limital.
- 5. Agyrtria linnæi. Extra-limital.
- 6. Momotus cœruleiceps. Extra-limital.
- 7. Ibis thalassina. Young of Plegadis guarauna.
- 8. Ardea wuerdemanni. Dichromatism of A. occidentalis.
- 9. Sterna "longipennis." Meaning S. pikii Lawr. Young of S. macrura.
- 10. Podiceps cristatus. Extra-limital, as far as known.

On the other hand, the numerous accessions to the list are in no less gratifying evidence of the progress of our knowledge. There are no fewer than one hundred and twenty additions to be made. The large majority of these are bona fide species, and actual acquisitions to the North American list, being birds discovered since 1873 in Texas, Arizona, and Alaska, together with several long known to inhabit Greenland. It may be here remarked that although the Greenland Fauna has long been usually claimed and conceded to be North American, yet the full list of Greenland

birds has never before \* been formally incorporated with the North American, as is done in the present instance. Aside from such additions, the increment is represented by species or (chiefly) subspecies named as new to science since 1873; by a few restored to the list; and by two imported and now naturalized species. The following is the full—

#### LIST OF ADDEND NAMES. [Continued on p. 10.]

- 1. Turdus migratorius propinquus. Since described by Ridgway. Western U. S.
- 2. Turdus iliacus. Greenland.
- 3. Harporhynchus curvirostris (verus). Restored. Arizona.
- 4. Cyanecula suecica. Alaska.
- 5. Regulus satrapa olivaceus. Recognized as a subspecies.
- 6. Parus rufescens neglectus. Since described by Ridgway. California.
- 7. Parus cinctus. Alaska.
- 8. Psaltriparus melanotis. Restored. Nevada. Arizona.
- 9. Catherpes mexicanus (verus). Restored. Texas.
- 10. Thryothorus ludovicianus miamensis. Since described by Ridgway. Florida.
- 11. Anorthura troglodytes pacificus. Recognized as a subspecies.
- 12. Telmatodytes palustris paludicola. Recognized as a subspecies.
- 13. Alauda arvensis. Greenland; "Alaska;" Bermudas.
- 14. Motacilla alba. Greenland.
- 15. Mniotilta varia borealis. Recognized as a subspecies.
- 16. Parula nigrilora. Since described by Coues. Texas.
- 17. Helminthophaga lawrencii. Since described by Herrick. New Jersey.
- 18. Helminthophaga leucobronchialis. Since described by Brewster. Mass.
- 19. Helminthophaga cincinnationsis. Since described by Langdon. Ohio.
- 20. Peucedramus olivaceus. Arizona.
- 21. Dendræca palmarum hypochrysea. Since described by Ridgway.
- 22. Siurus nævius notabilis. Since described by Grinnell. Wyoming.
- 23. Cardellina rubrifrons. Arizona.
- 24. Vireo flavoviridis. Restored. Texas.
- 25. Vireo solitarius cassini. Recognized as a subspecies.
- 26. Passer montanus. Naturalized.
- 27. Leucosticte atrata. Since described by Ridgway. Colorado.
- 28. Leucosticte australis. Recognized as a species.
- 29. Leucosticte tephrocotis litoralis. Recognized as a subspecies.
- 30. Ægiothus linaria holboelli. Recognized as a subspecies.
- 31. Ægiothus hornemanni. Greenland.
- 32. Astragalinus notatus. Restored. Kentucky.
- 33. Passerculus sandvicensis alaudinus. Recognized as a subspecies.
- 34. Ammodramus caudacutus nelsoni. Since described by Allen. Illinois.
- 35. Peucæa æstivalis illinoensis. Since described by Ridgway. Illinois.
- 36. Peucæa ruficeps boucardi. Arizona.
- 37. Junco hiemalis annectens. Recognized as a subspecies.
- 38. Junco hiemalis dorsalis. Recognized as a subspecies.
- 39. Junco hiemalis cinereus. Arizona.
- 40. Passerella iliaca megarhyncha. Recognized as a subspecies.
- 41. Molothrus æneus. Texas.
- 42. Sturnella magna mexicana. Texas.

<sup>\* &</sup>quot;A Catalogue of the Birds of North America," by Robert Ridgway, in Pr. Nat. Mus., ii, pp. 163-246, published since the above was written, includes Greenland birds, together with various Mexican species not yet found within our limits.

- 43. Icterus vulgaris. Restored. South Carolina.
- 44. Quiscalus purpureus æneus. Recognized as a subspecies.
- 45. Cyanocitta stelleri annectens. Recognized as a subspecies.
- 46. Perisoreus canadensis fumifrons. Since described by Ridgway. Alaska.
- 47. Sturnus vulgaris. Greenland.
- 48. Pitangus derbianus. Texas.
- 49. Myiodynastes luteiventris. Arizona.
- 50. Myiarchus erythrocercus. Texas.
- 51. Empidonax flaviventris difficilis, Restored. Western U.S.
- 52. Ornithium imberbe. Texas.
- 53. Nyctidromus albicollis. Texas.
- 54. Selasphorus alleni. Since described by Henshaw. California.
- 55. Calothorax lucifer. Arizona.
- Amazilia fuscicaudata. Texas.
- 57. Amazilia yucatanensis. Texas.
- 58. Iache latirostris. Arizona.
- 59. Chordediles popetue minor. Florida.
- 60. Crotophaga sulcirostris. Texas.
- 61. Picus stricklandi. Arizona.
- 62. Scops asio maxwellæ. Since described by Ridgway. Colorado.
- 63. Scops trichopsis. Inserted on Ridgway's authority. Arizona.
- 64. Strix cinerea lapponica. Recognized by Ridgway. Alaska.
- 65. Strix nebulosa alleni. Since described by Ridgway. Florida.
- 66. Surnia funerea ulula. Recognized by Ridgway. Alaska.
- 67. Speotyto cunicularia floridana. Since described by Ridgway. Florida.
- 68. Astur atricapillus striatulus. Recognized as a subspecies. Western N. Am.
- 69. Falco sacer obsoletus. Recognized as a subspecies.
- 70. Falco islandicus. Restored. Greenland.
- 71. Falco sparverioïdes. Florida.
- 72. Buteo albocaudatus. Texas.
- 73. Urubitinga anthracina. Arizona.
- 74. Thrasyaëtus harpyia. Texas.
- 75. Haliaëtus albicilla. Greenland.
- 76. Engyptila albifrons. Texas.
- 77. Coturnix dactylisonans. Naturalized.
- 78. Charadrius fulvus (verus). Alaska.
- 79. Charadrius pluvialis. Greenland.
- 80. Ægialites hiaticula. Greenland.
- 81. Vanellus cristatus. Greenland.
- 82. Hæmatopus ostrilegus. Greenland.
- 83. Gallinago media. Greenland.
- 84. Arquatella couesi. Since described by Ridgway. Alaska.
- 85. Pelidna alpina (vera). Greenland.
- 86. Actodromas acuminata. Alaska.
- 87. Limosa ægocephala. Greenland.
- 88. Rhyacophilus ochropus. Nova Scotia.
- 89. Numenius phæopus. Greenland.
- 90. Ardea cinerea. Greenland.
- 91. Grus canadensis (vera = fraterculus). Recognized.
- 92. Parra gymnostoma. Texas.
- 93. Rallus longirostris saturatus. Since described by Henshaw. Louisiana.
- 94. Porzana maruetta. Greenland.
- 95. Cygnus ferus. Greenland.
- 96. Cynus bewicki. Restored. Arctic America.

- 97. Anser albifrons (verus). Greenland.
- 98. Bernicla brenta nigricans. Recognized as a subspecies.
- 99. Somateria mollissima dresseri. Recognized as a subspecies.
- 100. Phaëthon æthereus. Newfoundland.
- 101. Phalacrocorax violaceus resplendens. Recognized as a subspecies. California.
- 102. Larus cachinnans. Alaska.
- 103. Larus affinis. Greenland.
- 104. Larus canus. Labrador.
- 105. Œstrelata bulweri. Greenland.
- 106. Podicipes auritus (verus). Greenland.
- 107. Brachyrhamphus brachypterus. Restored. Pacific Coast.
- 108. Brachyrhamphus hypoleucus. California.
- 109. Brachyrhamphus craverii. California.
- 110. Lomvia troile californica. Recognized as a subspecies. California.

The original number of names, 778, minus 10, plus 120, gives the total of 888 of the present edition of the "Check List." The number seems large, in comparison, and I am free to confess that it includes some — some twenty or thirty, perhaps — which my conservatism would not have allowed me to describe as valid, and the validity of which I can scarcely endorse. I have nevertheless admitted them to a place, because I preferred, in preparing a "Check List" for general purposes, rather to present the full number of names in current usage, and let them stand for what they may be worth, than to exercise any right of private judgment, or make any critical investigation of the merits of disputed cases. Probably, however, there are not more than thirty cases of birds retained in this list whose claims to be recognized by subspecific names can be seriously questioned.

It should be observed, that the list is not yet to be regarded as finally filled. Our southern border has proved so fruitful of Mexican species, that various others doubtless remain to be there detected; and several species described as Texan by Giraud in 1841 remain to be confirmed. With the accessions that may reasonably be expected, and under current usage in the discrimination of subspecific forms, the list will probably in a few years contain about 900 names of birds occurring in North America north of Mexico and inclusive of Greenland.

It is to be added here, that the present southern boundary of "North America" is a political one, wholly arbitrary so far as natural Faunal areas are concerned. It would be far more satisfactory, from a scientific standpoint, to ignore the present political line, and construct the "North American" list upon consideration of the limits of the "Nearctic Region" of Sclater and Baird. This would be to extend our area along the table-lands and higher region of Mexico to about the Isthmus of Tehuantepec, but not so far in the tierras calientes of either coast of that country: on an average about to the Tropic of Cancer. Such course would give us the natural instead of the political Ornis of our country; and I have no doubt that it will some day be taken. A few Cape St. Lucas birds have been so long in the "North American" list, that it is not thought worth while to displace them; but with these exceptions, it is not intended to include any species not known to occur north of Mexico.

Aside from those modifications which affect the ornithological or scientific status of the "Check List," the changes in nomenclature are numerous and in many cases radical. Without counting merely literal changes in the spelling of words, nominal changes are made for one or another seeming good reason in upwards of 150 cases. In probably not more than 30 of these, however, is the ornithological status of any bird modified; the changes being simply nomenclatural.

This portion of the subject is concluded with the following table, showing the number of birds ascribed to North America by several authors who have published complete lists from 1814 to the present year.

#### SUMMARY COMPARISONS.

Total of	North Ame	rican Birds	g given by	WILSON	in	1814			022*
I Otal Ol	TTOTAL ZEING	nean Dira	given by						
66	"	66	66	BONAPARTE	66	1838			471 *
"	44	66	"	Brewer	"	1840			491
"	44	44	"	AUDUBON	"	1844			506*
"	"	44	66	Baird	"	1858	•		744 †
66	"	"	66	Coues	66	1874			778‡
46	"	44	66	RIDGWAY	66	1880			924§
"	66	66	"	Coues	"	1882			888 ¶

- \* Fide Baird: I have not made the count myself.
- † The number is ostensibly 738; but 5 numbers are duplicated in printing, and 1 species is not numbered, making 744; of which 22 are admitted to be extra-limital, but enumerated.
- ‡ Total of numbered species in the body of the Check List 635; actual number of species and subspecies 750; with 28 additional in the Appendix, making 778.
- § Total of numbered species in the Catalogue 764; actual number of species and subspecies 924; of which 37 are admitted to be extra-limital, for all that is known to the contrary; and several others do not appear to be fully established as North American.
  - ¶ Being the 778 of the orig. ed., minus 10 subtracted, plus 120 added, = 888.

Note. Mr. Ridgway's Catalogue contains the following 52 names of birds which I do not admit in the Check List, for reasons which may be inferred from the remarks set against each of them. But the Mexican (not insular) species may all be expected over our border; and the recognition of subspecies in some cases depends upon the perspective in which we may elect to view them.

- 1. Harporhynchus graysoni. Extra-limital. Socorro Is., NW. Mexico.
- 2. Regulus obscurus. Extra-limital. Guadalupe Is., Lower California.
- 3. Regulus cuvieri. "Pennsylvania" (Audubon). Not since identified.
- 4. Parus meridionalis. Extra-limital. Mexico. Since found in Arizona.
- 5. Certhia familiaris mexicana. Extra-limital. Mexico.
- 6. Salpinctes obsoletus guadalupensis. Extra-limital. Guadalupe Is.
- 7. Thryomanes brevicauda. Extra-limital. Guadalupe Is.
- 8. Troglodytes insularis. Extra-limital. Socorro Is.
- 9. Parula pitiayumi insularis. Extra-limital. Socorro Is.
- 10. Perissoglossa carbonata. "Kentucky" (Audubon). Not since identified.
- 11. Dendræca montana. "Pennsylvania" (Wilson). Not since identified.
- 12. Wilsonia minuta. "New Jersey" (Wilson). Not since identified.
- 13. Setophaga miniata. "Texas" (Giraud). Doubtless.
- 14. Ergaticus ruber. "Texas" (Giraud). Doubtless.
- 15. Basileuterus culicivorus. "Texas" (Giraud). Doubtless.

- 16. Basileuterus belli. "Texas" (Giraud). Doubtless.
- 17. Lanius ludovicianus robustus. "California" (Gambel). Doubtful.
- Progne subis cryptoleuca. Florida. If recognized as distinct.
   Euphonia elegantissima. "Texas" (Giraud). Doubtless.
- 20. Carpodacus purpureus californicus. California. If recognized as distinct.
- 21. Carpodacus amplus. Extra-limital. Guadalupe Is.
- 22. Chondestes grammicus strigatus = grammicus.
- 23. Junco insularis. Extra-limital. Guadalupe Is.
- 24. Pipilo maculatus consobrinus. Extra-limital. Guadalupe Is.
- 25. Pipilo maculatus carmani. Extra-limital. Socorro Is.
- 26. Passerina parellina. Extra-limital. Mexico. (Texas, doubtless.)
- 27. Icterus wagleri. Extra-limital. Mexico.
- 28. Quiscalus palustris. "California" (Gambel). "Louisiana?" (Ridgway). Dubious.
- 29. Aphelocoma ultramarina couchi. Extra-limital. Mexico.
- 30. Myiozetetes texensis. "Texas" (Giraud). Doubtless.
- 31. Empidonax fulvifrons (verus). "Texas" (Giraud). Doubtless.
- 32. Pachyrhamphus major. Extra-limital. Mexico.
- Hadrostomus aglaiæ. Extra-limital. Mexico.
- 34. Picus villosus leucomelas. NE. N. Amer. If recognized as distinct.
- 35. Colaptes auratus hybridus. Intermediate specimens of unstable character.
- 36. Colaptes rufipileus. Extra-limital. Guadalupe Is.
- 37. Momotus caruleiceps. Extra-limital. Mexico.
- 38. Rhynchopsitta pachyrhyncha. Extra-limital. Mexico.
- 39. Comurus holochlorus brevipes. Extra-limital. Socorro Is.
- 40. Bubo virginianus subarcticus. Wisconsin. If recognized as distinct.
- 41. Bubo virginianus saturatus. N. coast of N. A. If recognized as distinct.
- 42. Falco albigularis. Extra-limital. Mexico, and C. and S. Am.
- 43. Æsalon regulus. Extra-limital. "At sea, off Greenland, lat. 57° 41' N., long. 35° 23' W."
- 44. Tinnunculus alaudarius. Extra-limital. "At sea, off Cape Farewell, Greenland."
- 45. Polyborus lutosus. Extra-limital. Guadalupe Is.
- 46. Buteo vulgaris. Michigan (Maynard). Identification in question.
- 47. Buteo borealis socorroensis. Extra-limital. Socorro Is.
- 48. Oreortyx picta plumifera. S. and L. California. If recognized as distinct.
- 49. Sula cyanops. Extra-limital. Socorro Is.
- 50. Sula piscator. Extra-limital. Socorro Is.
- 51. Diomedea culminata. Extra-limital. "Off Columbia River" (Audubon).
- 52. Lomvia arra brunnichi. If recognized as distinct.

#### POSTSCRIPT.

During the printing of the List, and since the preceding pages were stereotyped, the following additions have been announced. They will be found at the end of the list, raising the addend names from 110 to 120, and the whole number from 878 to 888.

- 111. Parus meridionalis. Arizona.
- 112. Myiarchus crinitus cooperi. Arizona.
- 113. Antrostomus vociferus arizonæ. Since described by Brewster. Arizona.
- 114. Buteo brachyurus. Florida.
- 115. Buteo fuliginosus. Florida.
- 116. Eurinorhynchus pygmæus. Alaska.
- 117. Fulica atra. Greenland.
- 118. Fuligula rufina. New York.
- 119. Œstrelata gularis. New York.
- 120. Puffinus borealis. Since described by Cory. Massachusetts.

[December, 1881.

#### REMARKS ON THE USE OF NAMES.

#### § 1. ETYMOLOGY, OR DERIVATION.

Etymology, the ἐτυμολογία of the Greeks, consists in tracing the derivation of a word back to the root from which it springs, explaining its formation, inflection, and application, thereby more clearly illustrating its virtue or quality than can be done by merely considering any one of the various meanings it may in time acquire. For a good illustration of this definition, see the word Cardinalis.

The large majority of the scientific names of birds are Latin or Greek words, or modern compounds of such, derived conformably to the rules for the construction of classic terms. In general, therefore, it is easy to give the exact meaning of the names in their original acceptation, and to point out their applicability as terms descriptive of the objects designated. On the whole, it has not been our design to go beyond a good fair definition of these Greek and Latin words, considering that all practical purposes are thus subserved. Many of the classic words being themselves derivatives, and the field of philological inquiry being boundless, it was necessary to keep within certain limits; and we have therefore seldom found it advisable, even were it practicable, in a case like the present, to trace words back of their recognized stems. Yet there will be found in the present little treatise, it is believed, much philological information of interest and actual value to all who desire to be put at their ease in the use of the Greek and Latin names of birds.

Many pure Greek or Latin names of birds known in classic times have been transferred in ornithology, in a wholly arbitrary manner, to totally different species. Thus the *Trochilus* of the ancients was an Egyptian Plover; in ornithological nomenclature, it is a genus of American Humming-birds. So also, many proper names, and many of the epithets which classic writers were so fond of bestowing, have been adopted as generic or specific names of birds, with little reason or with none, except the will of the namer. The genus *Iache* has no more to do with the Greek battle-cry than the name of Smith or Brown has to do with trade or color.

The remaining names, not classic in origin, are a miscellaneous lot not easy to characterize tersely. Many are modern geographical or personal names in Latin form; as, wilsoni, genitive case of Alexander Wilson's name, Latinized Wilsonus; or wilsonianus, an adjectival form of the same; americana for American; hudsonicus, after the territory named for Henry Hudson; noveboracensis, which is literally, inhabiting New York. Some others are post-classic, or late Latin, though in perfectly good form; and there are more of these, we find, than is generally supposed. Not a few are wholly barbarous, as Pyranga, Guiraca; and some of these, as cheriway, wurmizusume, are barbarous in form as in fact. Some are monstrous combinations, like Embernagra from Emberiza and Tanagra, or Podilymbus from Podiceps and Colymbus. Some are simply Latin translations of vernacular names; as, Puffinus anglorum, the puffin of the English. Finally, some are anagrams, like Dacelo from Alcedo, or pure nonsense-words, as Dafila, Viralva, Xema.

The student who confidingly expects to discover erudition, propriety, and pertinence in every technical name of a bird, will have his patience sorely tried in discovering what lack of learning, point, and taste many words imply. Besides the barbarisms, anomalies, and absurdities already indicated, he must be prepared to find names used with as little regard for precision of meaning, almost, as those of Smith, Brown, and Jones. Nothing like the nice distinctions, for example, that the Romans made between ater and niger, both meaning "black," or between albus and candidus, "white," obtains in modern science, where names are too often mere sounds without sense, and where the inflexible rules of technical nomenclature compel us to recognize and use many terms of slight or obscure or entirely arbitrary applicability, if only they be not glaringly false or of express absurdity. Let him for example, compare the several birds whose specific name is fuscus, and see what color-blindness this word covers.

The large majority of the names being, as already said, of Greek or Latin derivation, we are enabled to give a reasonably full and fair account of their etymology, and to point out their significance and application. There are, perhaps, not two dozen words of the whole list which we are unable to explain and define.

#### § 2. ORTHOGRAPHY, OR SPELLING.

The literation of the scientific names is fixed and exact in nearly all cases. Their derivation being known, and their form having crystallized in a language "dead" for centuries, the proportion of cases in which the orthography is unsettled is comparatively small. In general, there is no alternative spelling of a Greek or Latin word, and the modern derivatives are or can be compounded according to rules so fixed as to leave little latitude. In some instances, of course, two or more admissible forms of the same word occur: as hyemalis or hiemalis, cæruleus or cæruleus, Haliæëtus or Haliaëtus. But, in general, there remains only one right way of spelling, and that way easily ascertained. We say, there remains; for of course

there were centuries when the classics were undergoing the incessant changes incident to all spoken or living languages, just as our tongue is now. But having, in the usual process of evolution, reached that point which we mean when we use the term "classic," the Greek and Latin have come down to us in a certain form, so measurably fixed as to permit no decided ulterior modification. Our orthography, as far as possible, should reflect the purity and lucidity of such crystallization; and a little care will enable us to make such reflection clear.

In the cases of actual Greek and Latin words employed as names of birds, there are probably not in the whole list a dozen instances of words which admit of defensible alternative spelling. In the modern compounds of Greek and Latin stems, there is necessarily some little margin for variability; but in all cases, perhaps, at least a defensible orthography may be attained, though some alternative may not be without its claims to consideration. We can only say, that in this matter we have endeavored to reach good results according to definite recognized rules.

In the much-vexed question of forming quasi-Latin genitives from the names of persons, we have adopted the following simple and uniform rule: If the word ends with a consonant add single i for a man's name,  $\alpha$  for a woman's name; if ending with a vowel, change that vowel to i; as bairdi, cassini, but lawrencii, bonapartii; blackburnæ, graciæ. There are but few exceptions to this, as annæ, costæ. letter y gives the most trouble: it is best generally to treat it as a consonant, and say suckleyi, ridgwayi; but it must sometimes be rendered by i, as luciæ for Lucy (Latin Lucia), derbianus from Derby. It is rarely that a case occurs that such practice cannot readily meet. Names of birds derived from those of persons may of course be from any language, and consequently offer combinations of letters unknown in Latin; but it is useless to attempt to Latinize them, further than by giving them a Latin genitive termination. We should be led into the pedantry of brunonis for browni, or even of nigri for blacki, if we attempt any systematic Latinization of "barbarous" proper names. It is best to apply the above rule even to names already Latin in form, and write, for instance, blasiusi, not blasii. desirability of such conventional proceeding may be illustrated in the case of a bird named after a Mr. Wilcox; better wilcoxi, and be done with it, than vilcocis.

Hitherto, we have spoken of Latin and Greek names of birds indiscriminately. It will be remembered, however, that we are supposed to write the names always in Latin, be they of that language or actually Greek. This brings up the subject of the transliteration of words from the latter into the former. Most of the letters of the Greek alphabet have their exact and simple equivalents in Latin; but some can only be represented by two Latin letters, and some combinations of Greek letters change in passing into Latin words.

The following are the simple equivalents: a=a;  $\beta=b$ ;  $\gamma=g$ ;  $\delta=d$ ;  $\epsilon=\check{\epsilon}$ ;  $\zeta=z$ ;  $\eta=\bar{\epsilon}$ ;  $\iota=i$ ;  $\lambda=l$ ;  $\mu=m$ ;  $\nu=n$ ;  $\dot{\xi}=x$ ;  $o=\check{o}$ ;  $\pi=p$ ;  $\rho=r$ ;  $\sigma$  or s=s;  $\tau=t$ ;  $\omega=\bar{o}$ .

The following are simple substitutions:  $\kappa = c$ ; v = y.

The following are expressed by two letters:  $\theta$  or  $\vartheta = th$ ;  $\phi = ph$ ;  $\chi = ch$ ;  $\psi = ps$ . The letter  $\zeta$ , though written single z, is double, and equals dz.

There being no letter h in Greek, the aspirate is expressed by the sign f, preceding a vowel or written over it; thus  $\hat{a}$ ,  $\hat{\epsilon}$ ,  $\hat{b}$ ,  $\hat{b}$  = ha, he, ho, hy. The letter  $\rho$  also takes the aspirate, in which case  $\hat{\rho} = rh$ ; and when  $\rho$  is doubled, the second is followed by h;  $\hat{\rho}\hat{\rho} = rrh$ .

It is needless to give formal examples of these rules here; for the reader will find one or more of them illustrated on any page following the introductory matter.

#### § 3. ORTHOEPY, OR PRONUNCIATION.

Correct pronunciation of Greek and Latin is a lost art. The best we can do now is to follow the usage of those scholars who conform most nearly with what they show reason for supposing to have been the powers of the letters as spoken by the Greeks and Romans. Unfortunately for the student, there are three reputable schools who pronounce certain letters, especially the vowels a, e, and i, so differently that their respective methods are irreconcilable.

- I. The English Method. In England, and generally in America, excepting in the Jesuit colleges, the letters have nearly or exactly their English powers. This school teaches us "how not to do it," that is, to pronounce as the Greeks and Romans never did. If we imagine a dialogue between an English Professor of Latin and the Manes of Cicero, we are bound to infer that they would not understand each other; in fact, that neither would know that the other was talking Latin; though they might write to each other in identical words. Obviously, therefore, the English method is to be shunned. If the student will pronounce any word in the following list as if it were English, he will give it a sound the furthest possible removed from the right sound. The only excuse for the English method we ever heard is, that, as we do not know the right pronunciation, a conventional and consistent substitute is better than any doubtful approximation; but such talk is a mere apology for the English pis aller, not a defence of that sorry makeshift.
- II. The Continental Method. This is universal in Europe, excepting in England, and has gained much ground in America through the teaching of the Jesuits and other learned scholars. It is also known as the Italian school. It may be defined, in brief, as a compromise between English Latin and Roman Latin; the vowels having nearly or quite what is believed to have been their sounds as spoken by the Romans, while the consonants are heard more nearly in their English powers. Leading features of the school are: long a as in father; long e as English a in fate; long i as in machine; long a as English a in moon; a, as a vowel, practically like a; a like

y; c and g hard or soft as they would be in English, and most other consonants as in English, nearly or exactly.

III. The Roman Method. This way of speaking Latin, if practicable, is obviously preferable; and it is believed that a close approximation to Latin orthoepy is feasible. "The world over, nearly all the Latin grammarians of the last quarter of a century have urged a return to first principles. The Latin has rights of its own, and a demonstrated pronunciation which should be respected." The credit of leading this reform in America has been ascribed to the late Professor S. S. Haldeman, of the University of Pennsylvania, whose "Elements of Latin Pronunciation" was published at Philadelphia in 1851.

Nevertheless, the practicability of introducing such radical reform among naturalists, to most of whom the writing and speaking of classical words is but an incident of their scientific studies, may be seriously doubted, however desirable it is to do so. We question whether ornithologists, of this generation at least, can be induced to say Kikeronia, Kirke, and Pikicorwus, or Chicheronia, Chirche, and Pichicorvus for Ciceronia, Circe, and Picicorvus, or wirraynce for virens. It may be most judicious at present, and best on the whole, to pave the way for the final consummation by carrying into practice the many points on which scholars agree, without insisting upon the extremes respecting which diversity of good authority is admitted.

Upon such understanding we offer, for pronouncing the Latin names of North American birds, a scheme which insists upon the Roman sounds of the vowels and diphthongs, but yields the point in the disputed cases of certain consonants; conceding, for example, that c may remain soft before e, i, and y, and that v need not be turned into w. We do not profess to go into the subtleties, or even all the niceties of Latin orthoepy. Much of the end we have in view will be attained, if we can succeed in preventing those barbarisms and vulgarisms which constantly come from the lips of some persons of great accomplishment in the science of ornithology. Having ourselves heard Oh-nanth and Fully-gewler for Enanthe and Fuligula, we need not affect to conceal our belief that some ornithologists may profitably look a little further into the matter than they appear to have hitherto done.

#### Vowels.

The difference between a "long" and a "short" vowel is essentially one of quantity only, not of quality: it is actually the prolongation of a sound, not necessarily involving a difference in sound. Thus, if we dwell never so long on the "short" a of fat, it does not convert the sound of that letter into that heard in the "long" a of fate. The phonetic quality of a vowel should therefore be distinguished from its prosodiac quantity. Practically, however, no such discrimination is to be made in the case of the Latin vowels. We only know them as "long" or "short;" we determine their quantity by prosodiac rules, and make their quality

<sup>\*</sup> W. G. Richardson on Latin Pronunciation: In Report of the Commissioner of Education for 1876. 8vo, Washington, 1878. p. 484.

correspondent. For all that is known to the contrary, the Romans may have had, for example, as many qualities of their a as we have in English; but as we know only their "long" and "short" a, it is simply a matter of more or less of the same sound of the letter, not a difference in sound. Our only resource, therefore, is to ascertain the natural or acquired quantity of the vowels according to the standard authorities, and pronounce them conformably therewith.

It is the rule, with few exceptions, that a vowel before two consonants, or before the double consonants x and z, is long. We are inclined to believe that in many cases the full length of the vowel itself is not implied, but rather the length of the whole syllable in which it occurs. For instance, in the word melanorhynchus, the vowel y is encased in five consonants; and the time required to speak the whole syllable -rhynch-, in metric composition, is what makes the y long. The Romans may have had the y as short in quality as the y's in our word pygmy. Nevertheless, we have no assurance of this, and can only mark the y long, which means that this syllable is to be pronounced -rheench-. Take the word fuscescens, again, where each vowel is followed by two consonants. In this country we seldom if ever hear any thing but sounds of all three of the vowels as short as if they were English. We must, however, mark them long, which is equivalent to directing the word to be called foosaysaynce. But it does not follow that a naturally short vowel lengthened only "by position" is to be sounded at full length. Thus, in ēffinis, īnsignis, ōbsoletus, from ăd-, ĭn-, ŏb-, the long mark indicates the quantity of the syllable rather than of the vowel. The chief exceptions above alluded to are furnished by the concurrence of a mute and a liquid, when the preceding vowel remains short, in prose, at least.

A vowel before a single consonant, or before another vowel, is short, as a rule; but there are so many exceptions to this, that each case of the kind requires to be considered on its own merits. An accented vowel is likely to be long from this cause alone. Diphthongs are long, except before another vowel.

In Latin words derived from the Greek, the vowels e and o are likely to be long or short, according to whether they stand for Greek eta or epsilon, omicron or omega. So, also, the Latin i is long when representing the Greek diphthong  $\epsilon_i$ , as it often does; and a vowel is likely to be long when in any case it comes by the contraction of two or more vowels into one. Thus, the frequent Latin termination -pus, from the Greek pous, is long, or should be, like the proper Latin pes (foot).

With these slight remarks, we take up the vowels, diphthongs, and consonants in alphabetical order.

A. Orthoepists reckon from four to seven sounds of this vowel in English, the four usually recognized being those heard in fate, fat, fat, fatl. The English sounds of a in fate, fat, and fatl are unknown in Latin. Long a in Latin is always sounded as a in psatm; it is almost exactly the English interjection ah!—the name of the letter r without any roll. Short Latin a is the same sound, but with less stress and less prolonged, like the a in diadem, or the final a in Maria, Amelia, Hannah. Thus

in the frequently recurring word americana, all three a's have the same quality, but differ in quantity; the first and the last a being short and the middle a long, simply because there is where the accent, or stress of voice, comes to prolong the sound. If the accent in this case were on the antepenult, all three a's would have exactly the same quantity and quality.

Long  $\bar{a}$  as in psalm.

Short ă as in diadem.

**E.** Long e has the sound of French  $\hat{e}$  in  $\hat{fete}$ , or English e in they, or English a in fate. Short e is like English e in them, not quite so short as in met; something between mate and met. Example of long e:  $\bar{e}xilip\bar{e}s$ , pronounced ache-seal-i-pace.

Long  $\bar{e}$  as in they.

Short ĕ as in them.

I. Long i is invariably like the English i in machine, police, oblique, pique; that is, the English ee in feet, ea in feat, &c.; but never the English i of fight, night. Short i is the same sound, but as brief and abrupt as possible, like English i in possible, ability, imitate. Short and long i are both heard in intrigue.

Long ī as in machine, pique.

Short i as in ability, imitate.

O. This letter, long or short, has always its pure English sound, there being no qualities of Latin o to correspond to such anomalies as the English o in move, more, come, &c.

Long  $\bar{o}$  as in old, no.

Short o as in odd, not.

U. It is not easy to correctly appreciate the powers of this vowel in Latin. Long u never has the sound of English u, eu, or ew, as in fury, feud, few; but is always broad as well as long, like o in move, oo in moon, fool. Short u is not the English u in tub or English o in love, but quite like the English u in bull, full. Take for example the common word  $r\bar{u}f\tilde{u}s$ , where the first u is long, the second short. This word is neither roof-uss, nor rewf-uss, nor rewf-uss, nor uss in the consonants permitted, it would rhyme exactly with uss-

Long  $\bar{u}$  as o in move, oo in moon, ue in rue.

Short  $\tilde{u}$  as in bull, full, pull.

Y. This letter, as a vowel, has practically the sound of i, long or short; more exactly, that of the German  $\ddot{u}$  (ue), as in  $M\ddot{u}ller$ , which is nearer  $M\ddot{i}ller$  than Muller. It is scarcely a Latin letter, and chiefly occurs in words from the Greek, corresponding to Greek upsilon; as hyperboreus, uropygialis.

It is to be remarked, that any vowel is or may be modified in quality as well as in quantity by its consonantal combination, this being especially the case when followed by the letter r. It is as if the r were rolling away, and dragging the vowel after. Compare fuscus with turtur; the first with the last syllable of turdus,

&c. We suspect that some of the less evident powers ascribed by orthoepists to various vowels, are not inherent in the vowels themselves, but due to consonantal modification of the sound.

Let us add that orthoepists commonly and with great propriety recognize what they call the "neutral" vowel-sound, a quality so slight and obscure, that any one of the vowels may express it indifferently. Thus, if we pronounce the word martyr as rapidly as possible, it makes scarcely any appreciable difference whether it be written martar, marter, martir, martor, martur, or martyr; as we say scarcely any thing more than martr, the six "neutral" vowels are phonetically interchangeable.

#### DIPHTHONGS.

In diphthongs, each vowel must be sounded, and the two sounds be smoothly combined. Two vowels coming together do not necessarily form a diphthong. For example,  $a\ddot{e}r$  is a word of two syllables, and  $a\ddot{e}don$  one of three; the vowels in these cases to be separately and distinctly uttered, as in English  $a\ddot{e}rial$ . Proper diphthongs, i.e., two vowel-sounds combined to make a third different from either, are comparatively rare; and all the following components of diphthongs also come together without combining.

Æ consists of ah-ay, which when rapidly spoken becomes so nearly like Latin long  $\bar{e}$  (see above) as to be practically the same. It was originally written ai, and is by some directed to be so sounded.

AI is a very composite sound. *i* itself is a compound, being *ah-ee*, the whole being therefore *ah-ah-ee*, which when run together becomes very nearly our English *eye* or the pronoun *I*. It seems quite like the French *naif*, *naive*, or English *knife*.

A and O do not combine, and seldom come together.

AU is oftenest heard, but wrongly, as in cause, or as aw in awl, law, awful. It is like the ow in how, now, owl. It is precisely the German au, as in auth.

E and A do not combine; they frequently come together, especially at the ends of words, but each is separately pronounced. E.g., Ene-as Bore-as, Arde-a.

EI is frequent. The analysis is ay-ah-ee, contracted to a drawling sound little different from long English a in mate; more exactly, English ei in vein, eight.

E and O do not combine. E-os, E-opsaltria, &c.

EU is equal to ay-oo. Strongly and rapidly uttered, it becomes the long English u in tube, ue in due, ew in few, eu in feud, ou in you; and especially when initial represents the whole word you. For example Eugenes = Yougenes = Ayoogenes. It seldom occurs, except in Greek words.

IA, IE, II, IO, IU do not combine. The very frequent ia, especially ending a word, and the ii, so frequent in the genitives of persons' names, are always two full syllables. The common iu, in the ending of words makes two syllables: e.g., spuri-us. So seri-es, rati-o have each three syllables. Some apparent diphthongs of vocal i with a following vowel, are really of consonantal i, which is j, pronounced g; as plebeius, plebejus, pronounced plebejus.

OA and OO and OU do not combine; bo-ops has two, arcto-us or arcto-a three, and o-ology four syllables. ou diphthong very early passed into long  $\bar{u}$ .

OE, when fully but rapidly said in combination, seems to yield the diphthong  $\alpha$  preceded by a slight w sound; the whole nearly as the English word way. If not this, it is indistinguishable from Latin  $\alpha$ . We are inclined to say way-nanthe for  $\alpha$ -nanthe; if not this, then ay-nanthe, not  $\alpha$ -nanthe nor  $\alpha$ -nanthe. The combination is sometimes interchangeable with  $\alpha$ , as  $\alpha$ -nanthe nor  $\alpha$ -nanthe. It is to be carefully distinguished from  $\alpha$  and  $\alpha$ -nanthe in  $\alpha$ -nanthe  $\alpha$ -nanthe

OI. These two letters may combine or not. Generally they do not, each being a distinct syllable. Thus, *Pic-o-i-des* is a word of four syllables, the second and third of which are *o-ee*. oi in combination is given by some as in English oil, but is perhaps more nearly the French oei in œil. As ai passed into æ, so oi early became æ, and some direct the letter to be sounded as oi.

UA and UE, in combination, yield sounds like English wah and way; as suavis, suecica.

UI, equivalent to oo-ah-ee, is like the French oui (yes), very nearly the English pronoun we. The rare UU seems to be simply  $\bar{u}$  at extreme length: equus.

Y making a diphthong with a following vowel gives the sound of such vowel preceded by w; as, Myiarchus = Mweearchus. It only occurs in Greek words, by transliteration for upsilon.

In some cases three or four vowels come together; but the pronunciation may usually be determined by the foregoing rules. Thus: Agelaus, Poacetes, Haliaetus. In these cases respectively ae and oe are combined, and pronounced as above said; the other vowels are distinct. Hal-i-a-e-tus is a word of five syllables. My-i-o-di-oc-tes is one of six syllables, though in practice reduced to five, by slurring the y and i together. In trudeaui, again, are four vowels together; but in this case eau combine into long o, and the word has but three syllables.

#### CONSONANTS.

Most of the consonants have their English powers, pure and simple. Some, however, call for remark, especially in certain of their combinations.

The letters c and g are now said to be "always hard," without qualification. It is a much vexed question. As it is not demonstrated that the Romans had no soft c and g, we do not see that we may not be permitted to retain these sounds.

C then is hard, like k, before a consonant or a, o, u, soft before e, i, y, and before the diphthongs a, ai, oe, oi. ch is always hard; there is no sound of ch as in church, still less as in chaise, in Latin.

G is hard or soft under the same circumstances as c, with the important exception, that it is hard before y in words derived from the Greek, when the y results from the Greek upsilon (v). Example: Gymnocitta, not Jymnocitta.

J is simply i, interchangeable with it, and always pronounced like the y in yes, or as in hallelujah.

N followed by c hard, k, g, or x has a nasal or twanging sound of ng; as in English ankle, anger, pronounced ang-kle, ang-ger. Preceded by m or g, it does not destroy these letters: as Mniotilta, Gnathodon.

P is not silent before s; thus in psaltria articulate both. So in the digraph ph, some direct to sound both, as in up-hill. It is difficult, if not impossible, to articulate both letters, especially when, as often happens, a th succeeds. For example, in erythrophthalmus we find that we cannot make four sounds for the phth as in up-hill and hot-house. Practically ph becomes something between f and v, just as in Stephen or Steven. So also the original Indo-European aspirates bh, dh, gh are not retained in any European language; there is nothing to correspond to log-house.

QU is sometimes followed by another u, as in altiloquius, propinquius. It would seem to be rendered by kwooce.

R is strongly pronounced with a trill. It is heard at the height of its power in the combination rrh; as in catarrhactes, pyrrhorrhoa.

S invariably retains its sharp hissing sound. Thus essence is a rhyme with fuscescens (as far as the s-sound is concerned); so also virens = virraynce, not vy-renz. Compare hiss or this with his. So particular were the Romans to avoid the z sound of s, that they even altered antecedent consonants; saying, for example, urps and pleps for urbs and plebs.

T always preserves its sound. There is nothing to correspond with the English -tion = shun, &c. E. g., gra-ti-a, rat-i-o, init-i-um.

V is directed by some to be sounded like English w in we. But this is rarely done.

X is always ks or cs, never gz or z, even when initial, as in Xema, Xanthocephalus.

Z, which only occurs in Latin words of Greek extraction, is a double letter equivalent to dz, and the best authorities recommend the d sound to be articulated. Thus Aphriza, Spiza, are pronounced Afreedza, Speedza.

A word in regard to the pronunciation of modern proper names, as of persons and places, so often recurring in ornithology. After mature deliberation, we have decided to mark them for their pronunciation in the language to which they belong. It seems finical and pedantic to attempt to Latinize them; for to carry out that plan to its logical result would be to give brunonis instead of browni; and even then some names would utterly defy us, unless changed beyond all recognition. So we have adopted the rule of preserving the orthography and orthoepy of all modern proper names, even though containing the letter w. Barbarous geographical words of unsettled or no known orthography may, however, be sometimes dressed in quasi-Latin; thus it is perfectly permissible to render aoonalaschkae by unalascæ. We make this remark to explain what must seem inconsistent in our use of diacritical marks in some places; for we mark the vowels long or short as the syllables are pronounced in the language to which the word belongs, not as they would be in Latin.

#### ACCENTUATION.

This is a matter of prime importance. For elegant, even for bearable, pronunciation, it is essential to place the accent or stress of voice on the right syllable. Fortunately the rules are simple, with comparatively few exceptions.

Accent the penult when it is long.

Accent the antepenult when the penult is short.

These two rules will carry us safely across the great majority of Latin words. In many cases lengthening the syllable, whether penult or antepenult, is actually equivalent to accenting it. We can scarcely recall a case of a short accented penult; but many short antepenults take the accent, which is simply because it cannot be thrown still further back. Modern proper names of three syllables with the accent on the first, keep it there after addition of the *i* of the genitive case; as, aud'uboni, rich'ardsoni.

So important is the matter of accent, that were all other diacritical marks dispensed with, we could still pronounce the words with measurable accuracy, knowing where to put the stress of voice.

The tendency in English is constantly to throw the accent back as far as possible; and there is much of this same practice in the usual pronunciation of Latin. For the latter language, and especially for words derived from the Greek, we consider it vicious and undesirable. It seems to us much more sensible and natural in the case of a word compounded of two Greek words, to keep the stress of the voice on the stem of each, than to throw it, for sake of glibness, on the most insignificant syllable, often the mere connective vowel, and a short one at that. for example Troglodytes, Lophophanes, Phylloscopus, or any similar words of four syllables, compounds of two words of two syllables each. It is glib to accent the antepenult, but it is done at the sacrifice of the strength and dignity of the stem which stands penult, and which we should prefer to accent, even if short. Where we have found it practicable on etymological grounds to lengthen and accent such penults, we have done so; in general, however, we have closely conformed to routine custom, especially as there is to be strongly set before the inexperienced student the necessity of avoiding the glaring impropriety of accenting the penult of erythrocephalus, for example. The tendency of all persons who find it difficult to handle a long new word, is to dissect it, with two or even three accents; and perhaps the inclination of the scholar to show his erudition has unconsciously led him to the opposite extreme. Any "rule" or custom aside, the natural accent of polysyllabic words is rhetorical - as if each syllable were a word. It may be seen in those words whose looseness of composition, so to speak, leaves them like sentences; as ne'vertheless", not'withstand"ing. The naturalness of a'naly"tic, ge'omet"ric contrasts favorably with the conventionality of analysis, geo'metry; and there is nothing in the quality of the final syllables to account for the differences in accent. But we are aware that our views of this matter will not pass current, even if they escape adverse criticism.

#### EXPLANATIONS.

- 1. The names in the Check List are consecutively numbered from first to last, whether they be of species or of subspecies. The latter are sufficiently distinguished by consisting of three terms instead of two.
- 2. The names in the Dictionary are numbered to correspond, each page containing the same numbers of the two series.
- 3. The person's name in parentheses immediately after each bird's name is that of the original describer of the species or subspecies. The unenclosed name succeeding is that of the authority for the particular combination of generic, specific, and subspecific terms adopted. When the original describer is also the authority for the combination, a single unenclosed name is given.—The following are the principal abbreviations:—

All., Allen.	Gamb., Gambel.	Lawr., Lawrence.	Sw., Swainson.
Aud., Audubon.	Gir., Giraud.	Licht., Lichtenstein.	Temm., Temminck.
Bd., Baird.	Gm., Gmelin.	Nutt., Nuttall.	Towns., Townsend.
Bodd., Boddaert.	Gr., Gray.	Reich., Reichenbach.	V., Vieillot.
Bp., Bonaparte.	L., Linnæus.	Ridg., Ridgway.	Vig., Vigors.
Cab., Cabanis.	Lafr., Lafresnaye.	Scl., Sclater.	Wagl., Wagler.
Cass., Cassin.	Lath., Latham.	Steph., Stephens.	Wils., Wilson.

- 4. After these terms come three letters, "B," "C," and "R," each followed by a number. These stand respectively for Baird's List, 1858, Coues's Check List, 1874, and Ridgway's Catalogue, 1880. The number following each of these letters is that which the bird bears in such lists. Thus, Turdus migratorius was named by Linnæus, who is also the authority for the combination, and is 155 of Baird's list, 1 of Coues's, and 7 of Ridgway's. The dash after any one of these letters shows that the species is not contained in B, C, or R, as the case may be.
- 5. The note of exclamation, in parentheses, indicates that the species is in North America only a straggler from the country that the following initial letter denotes: E., Europe, A., Asia, M., Mexico, W. I., West Indies. G. shows the bird to be only North American as occurring in Greenland.
- 6. The note of interrogation, similarly enclosed, means that the name is considered to be of slight or uncertain value, as of a subspecies scarcely distinguished from its stock, or of a species not well known.
- 7. The Index will be found to contain matter additional to, or corrective of, that in the body of the work. See p. 137.

## CHECK LIST

OF

# NORTH AMERICAN BIRDS.

- 1. Turdus migratorius L. B 155. C 1. R 7.
- 2. Turdus migratorius propinquus Ridg. B —. C —. R 7a. (?)
  Rocky Mountain Robin.
- 3. Turdus migratorius confinis (Bd.) Coues. B —. C 1a. R 8. St. Lucas Robin.
- 4. Turdus iliacus L. B —. C —. R 6. (G. !E.) Redwing.
- 5. Turdus nævius Gm. B 156. C 2. R 9. Varied Thrush.
- 6. Turdus mustelinus Gm. B 148. C 3. R 1. Wood Thrush.
- Tūr'-dŭs mī-gră-tō'-rī-ŭs. Lat. turdus, a thrush. Lat. migro, to move from one place to another; migrator, a wanderer, a migrant; migratorius, migratory.
- T. m. prŏ-pīn'-qŭŭs [propeenkwooce]. Lat. propinquus, near, neighboring; as related to T. migratorius.
- T. m. con-fi'-nis [confeenis]. Lat. confinis, subs. or adj., a neighbor, neighboring; here
  in sense of closely related to T. migratorius.
- 4. T. il-I'-ă-cŭs. Lat. iliacus, relating to the ilia, or haunches; also, Lat. Iliacus, Gr. Ἰλιακόs, relating to Troy, Trojan; application obvious in neither case. But Aristotle gives a kind of thrush, called lλιάs, supposed by old ornithologists, as Gesner and Belon, to be this species, said to be called by the modern Greeks lλλάs, τυλάs, κίχλη lλιάs, οτ κίχλα lλιάδα; and the actual form, Turdus iliacus, was an old name when Linnæus adopted it.
- T. naē'-vī-ŭs [nayveus]. Lat. nævius, spotted, from nævus, a mole (birth-mark). The sub-genus Hesperocichla is Gr. έσπερος, Lat. vesperus, evening, i.e., western, and κίχλα or κίχλη, a thrush.
- T. mūs-tē-lī'-nūs. Lat. mustelinus, weasel-like; i.e., in this case, tawny.—The sub-genus
  Hylocichla is Gr. ΰλη, a wood, and κίχλα.

- 7. Turdus fuscescens Steph. B 151. C 6. R 2. Wilson's Thrush.
- 8. Turdus unalascæ Gm. B 150. C 4b. R 5. Western Hermit Thrush.
- 9. Turdus unalascæ auduboni (Bd.) Coues. B 149 var. C 4a. R 5a. Audubon's Hermit Thrush.
- Turdus unalascæ nanus (Aud.) Coues. B 149. C 4. R 56.
   Eastern Hermit Thrush.
- Turdus ustulatus Nutt. B 152. C 5b. R 4.
   Oregon Olive-backed Thrush.
- 12. Turdus ustulatus aliciæ (Bd.) Coues. B 154. C 5a. R 3. Gray-cheeked Thrush.
- Turdus ustulatus swainsoni (Cab.) Coues. B 153. C 5. R 4α.
   Olive-backed Thrush.
  - 7. T. fūs-cēs'-cēns [foosaysaynce]. Present participle of a supposed Lat. inceptive verb fuscesco, I grow dark or swarthy; Lat. fusco, of same signification. It means, or should mean, less than fuscus; i.e., somewhat dark; is not otherwise applicable to the lightest-colored thrush of this group.
  - 8. T. ū-nă-lās'-caē. Of the Island of Unalaska. It is permissible, indeed desirable, to resolve Gmelin's barbarous word aonalaschkae into a purer form. With this orthography the word is of sufficiently classical aspect, and corresponds with alascensis. See Anorthura, No. 78, and Passerella, No. 283.

This is *T. pallasi* var. nanus of the orig. ed. of the Check List. For the change, see Pr. Nat. Mus., ii, 1880, p. 1.

9. T. u. aūd'-ŭ-bŏn-ī. To John James Audubon, the "American Backwoodsman," and famous author of the "Birds of America."

This is T. pallasi var. auduboni of the orig. ed. of the Check List.

10. T. u. nā/-nŭs. Lat. nanus, from the Gr. νάννος or νάνος, a dwarf.

This is *T. pallasi* of the orig. ed. It is true that *nanus* has of late been applied exclusively to the Western form, the true *unaluscæ* Gm. But the name *nanus* was originally based by Audubon on a bird from Pennsylvania, and only later amplified by him to include the Western form. The long survival of an error does not justify its continued perpetuation after detection.

 T. ūs-tŭ-lā/-tŭs. Lat. ustulatus, perfect participle of ustulo, I scorch, singe; with reference to the ashy coloration, as if the bird had been charred.

This stands as *T. swainsoni* var. ustulatus in the orig. ed. The case is precisely parallel with that of nanus vs. pallasi; for Nuttall named the Oregon bird ustulatus in 1840, and Cabanis did not apply the name swainsoni to the Eastern Olive-backed Thrush till several years afterward.

 T. u. ă-lī'-cĭ-aē. To Miss Alice Kennicott, sister of Robert Kennicott, of Illinois. See Scops, No. 466.

This is T. swainsoni var. aliciæ of the orig. ed. See No. 11.

13. T. u. swaīn'-sŏn-ī. To William Swainson, the zealous and accomplished English naturalist.

This is T. swainsoni of the orig. ed. See No. 11.

- 14. Oroscoptes montanus (Towns.) Bd. B 255. C 7. R 10. Mountain Mocking-bird.
  - 15. Mimus polyglottus (L.) Boie. B 253. C 8. R 11. Mocking-bird.
  - 16. Mimus carolinensis (L.) Gr. B 254. C 9. R 12. Cat-bird.
  - 217. Harporhynchus rufus (L.) Cab. B 261, C 10, R 13, Brown Thrush; Thrasher.
  - 18. Harporhynchus rufus longirostris (Lafr.) Coues. B 260, C 10a, R 13a, Texas Thrasher.
  - 19. Harporhynchus curvirostris (Sw.) Cab. B 259. C —. R. 15. Curve-billed Thrasher.
  - 20. Harporhynchus curvirostris palmeri Ridg. B —. C 11. R 15a. Bow-billed Thrasher.
  - 21. Harporhynchus bendirii Coues. B —. C 11bis. R 14a. Arizona Thrasher.
  - 22. Harporhynchus cinereus Bd. B —. C 12. R 14. St. Lucas Thrasher.
  - 14. O-rō-scōp'-tēs [-tace] mōn-tā'-nŭs. Gr. ὅρος, a mountain, σκώπτης, a mimic; σκώπτω, I mock, deride, jeer at. The orthography differs; authority may be found for either Oroscoptes or Oreoscoptes; the former was originally written by Baird; it is shortest: and we usually say orology, orography, &c. Lat. montanus, relating to mons, a mountain.
  - 15. Mi'-mŭs [meemus] pŏl-ȳ-glōt'-tŭs. Lat. mimus, Gr. μᾶμος, a mimic.—Gr. πολύγλωττος, polyglot, from πολύς, many, γλῶττα, tongue.
  - 16. M. ca-ro-lin-ēn'-sis. Lat. for Carolinian, of Carolina; Carolus, Charles, is the modern Lat. form of Germ. Rari, or Reri, a peasant; A. S. ceorl, Scot. carle, Eng. churl. Carolina is by some derived from Charles II. of England; but Ribault, in 1562, built in Port Royal a fort he called Charlesfort, and Laudonnière, who came to relieve Ribault's colonists in 1564, one which he says, "je nommay la Caroline, en honneur de nostre prince le roy Charles [IX., of France]."
  - 17. Hār-pŏ-rhȳn'-chus [rh very strong; ch as k] rū'-fus. Gr. ἄρπη, a sickle; ῥύγχοs, a beak; i. e., bow-billed. The former word is seen in harpy, so called from its hooked beak. Some purists will have the r doubled in this and all such cases, making Harporrhynchus; but the current of modern usage has set too strongly against it to be stemmed without liability of seeming pedantic. Lat. rufus, rufous, reddish.
  - H. r. lon-gi-ros/-tris [loang-gi-roas-tris]. Lat. longus, long, rostris, beaked, from rostrum, beak.
  - 19. H. cūr-vĭ-rōs'-trĭs. Lat. curvus, curved; and rostris. Not in the orig. ed.
  - 20. H. c. pāl'-měr-ī [sound the l]. Dedicated to one Edward Palmer.
  - 21. H. běn-dī'-rī-ī. To Capt. Charles Bendire, U. S. Army.
  - 22. H. cĭn-ĕr'-ĕ-ŭs. Lat. cinereus, ashy, or ash-colored; from cinis, genitive cineris, ashes. Gr. κόνιs, of same meaning, apparently from καίω, κάω, I burn. Related English words are incinerate, cinder, &c.

- 23. Harporhynchus redivivus (Gamb.) Cab. B 256. C 13. R 16. Californian Thrasher.
- 24. Harporhynchus redivivus lecontii (Lawr.) Coues. B 257. C 13a. R 16a. Yuma Thrasher.
- 25. Harporhynchus crissalis Henry. B 258. C 14. R 17. Crissal Thrasher.
- 26. Saxicola cenanthe (L.) Bechst. B 157. C 15. R 21. (!E.) Stone Chat; Wheat-ear.
- 27. Sialia sialis (L.) Hald. B 158. C 16. R 22. Eastern Blue-bird.
- 28. Sialia mexicana Sw. B 159. C 17. R 23. Western Blue-bird.
- 29. Sialia arctica Sw. B 160. C 18. R 24.

  Arctic Blue-bird.
- 30. Cinclus mexicanus Sw. B 164. C 19. R 19. Water Ouzel; Dipper.
- 31. Cyanecula suecica (L.) Brm. B —. C —. R 20. (!A.) Blue-throated Redstart.
- 23. H. rĕ-dĭ-vī'-vŭs. Lat. redivivus, revived, from re-, red-, redi-, in sense of back again, and vicus, living. Gambel discovered in this bird a long-lost species of an older author.
- 24. H. r. lĕ-cŏn'-tĭ-ī. To Dr. John L. Le Conte, of Philadelphia, the famous entomologist.
- 25. H. crīs-sā/-līs. No such Latin word; there is a verb criso or crisso, used of a certain motion of the haunches; crissum is a technical word lately derived therefrom, signifying in ornithology the under tail-coverts, which in this bird are red. Cf. Gr. κρισσός, κιρσός.
- 26. Sāx-ĭ'-cŏ-lā oē-nān'-thē [oo-ay-nanthe, as if way-nanthe]. Lat. saxicola, a rock-inhabitant; saxum, a rock, and incola (in and colo), an inhabitant.—Lat. vitiflora, and Gr. olvάνθη, signify precisely the same thing: the bird is prettily named "flower of the vine:" Lat. vitis, the vine, flora, a flower. The Gr. olvάνθη, whence Lat. αnanthe, is an uncertain bird mentioned by Aristotle and Pliny; the name was definitely applied to this species in 1555. The word primarily relates to the grape, οἴνη, as if the bird were one which frequented vineyards, or appeared with the flowering (ἄνθος) of the vine.
- 27. Sĩ-ăl'-ĭ-ă sĩ'-ăl-ĭs. Gr. σιαλίς, a bird, in "Ath. 392 F;" from σίαλον, saliva; verb σιαλίζω, I slaver, or make some sibilant noise. To call this Anacreon a slobberer!
- 28. S. mēx-ĭ-cā'-nă. Latinized from Mexican. The country is called Mexico, Mejico, or Mehico, from Mexilli, the Aztec god of war.
- 29. S. ārc'-tĭ-că. Lat. arctica, northern, arctic; i. e., Gr. ἄρκτος, a bear, ἀρκτικός, near the bear.
- 30. Cīn'-clŭs mēx-ĭ-cā'-nŭs. Gr. κίγκλος, Lat. Cinclus, the name of a bird, by some supposed to be the European Cinclus aquaticus, by others a kind of Sandpiper; κιγκλίζω is to wag the tail.— Lat. mexicanus, see No. 28.
- 31. Cỹ-ăn-ĕ'-cŭ-lă suē'-cĭ-că. Cyanecula is a diminutive substantive lately (perhaps not before Brisson, 1760) formed from the Lat. adjective cyaneus, Gr. κυάνεος οr κυανός, blue; meaning, as we might say, "bluet." Rubecula is a word similarly coined. —Lat. suecica or svecica, Swedish; Sweden having been called Suecia or Svecia. In that country the bird is said to be called "Charles's-bird," Carls-fogel, whence Avis Carolina of some of the treatises written in Latin. "Redstart" is a corruption of Rothster, meaning "redtail," and Ruticilla and Phanicurus are among the translated book-names of the species.

Not in the orig. ed. of the Check List. See Ibis, 1878, p. 422. Alaska.

- 32. Phylloscopus borealis (Blas.) Dress. B —, C 20. R 34. (!A.) Kennicott's Warbler.
- Regulus calendula (L.) Licht. B 161, C 21, R 30.
   Ruby-crowned Kinglet.
- 34. Regulus satrapa Licht. B 162. C 22. R 33. Golden-crested Kinglet.
- 35. Regulus satrapa olivaceus Bd. B C R 33a. (?)
  Western Golden-crested Kinglet.
- 36. Polioptila cœrulea (L.) Scl. B 282. C 23. R 27. Blue-gray Gnat-catcher.
- 37. Polioptila melanura Lawr. B 284. C 24. R 29. Black-capped Gnat-catcher.
- 38. Polioptila plumbea Bd. B 283. C 25. R 28. Plumbeous Gnat-catcher.
- 39. Chamæa fasciata Gamb. B 274. C 26. R 35. Wren-tit.
- 40. Lophophanes bicolor (L.) Bp. B 285. C 27. R 36. Tufted Titmouse.
  - 32. Phỹl-lō'-scŏ-pus bŏr-ĕ-ā'-līs. Gr. φύλλον, a leaf; σκοπός, a watchman; σκοπέω, I look out, survey, examine; as these birds peer about in the foliage. Lat. boreas, the northwind, h. e., the north; borealis, northern.
  - 33. Rĕg'-ŭ-lŭs căl-ēn'-dŭ-lă. Lat. regulus, diminutive of rex, a king; exactly equivalent to "kinglet."— Calendula is a substantive which may be formed from the gerund of the verb caleo, I am warm; figuratively, glowing; in allusion to the fiery color on the head. It was apparently coined by Brisson, 1760, for the European Regulus cristatus, but was in 1766 appropriated by Linnæus to the present species. The early ornithologists had a great variety of names for these diminutive birds, mostly indicating royalty or other high station, in obvious reference to the "crown;" as Rex, Regulus, Regillus, Tyrannus or Τύραννος, Basiliscus or Βασιλίσκος, Presbys or Πρέσβυς, Βασιλεύς; to say nothing of Orchilus or 'Ορχίλος, Trochilus or Τροχίλος, Parus, Sylvia, Motacilla, Passerculus, Troglodytes, &c. The French Roitelet or Roytelet, and the German βιϋπίξειπ, correspond to "kinglet."
  - 34. R. sāt-rā'-pă. Lat. satraps, satrapes, or satrapa, Gr. σατράπης, from the Persian kbsbatram, meaning a crown or a kingdom: English satrap. Alluding to the bird's golden crown.
  - 35. R. s. ŏl-ī-vā-cĕ-ŭs. Late Lat. olivaceus, olivaceous, olive-colored. See Vireo, No. 170.
  - 36. Pŏ-lǐ-ōp'-tǐ-lă coē-rŭl'-ĕ-ă [sayrulea]. Gr. πολιόs, hoary, gray; πτίλον, feather; in allusion to the whitish edgings of the primaries.—Lat. cærulea or cærulea or cærulea, blue, azure. Any of these forms of the word is admissible. We prefer cærulea.
  - 37. P. měl-ăn-ū'-ră. Gr. μέλαs, fem. μέλαινα, black; οδρα, tail. See Index, p. 137.
  - 38. P. plūm/-bĕ-ă. Lat. plumbeus, plumbeous, lead-colored; from plumbum, lead.
  - 39. Chăm-aē'-ă [kam-ay-ah] fās-cĭ-ā'-tă. Gr. χαμαί, adverb, on the ground. Lat. fuscis, a bundle of faggots; hence, fusciatus, striped. The allusion is to the indistinct bands across the tail-feathers of the bird that lives in bushes close to the ground.
  - 40. Löph-ŏ'-phă-nēs [-nace] bī'-cŏl-ŏr. Gr. λόφος, a crest; and φαίνω, I appear; in allusion to the conspicuous crest.—Lat. bicolor, two-colored.
    - N.B.—The accentuation of this and many similar words is questionable, and perhaps arbitrary. We give the above in deference to technical rule, conformably with Aristo'phanes, &c. The actual usage, in this country at least, is Lŏph-ŏ-phā'-nēs; and

- 41. Lophophanes inornatus (Gamb.) Cass. B 287. C 28. R 38. Plain Titmouse.
- 42. Lophophanes atrocristatus Cass. B 286. C 29. R 37. Black-crested Titmouse.
- 43. Lophophanes wollweberi Bp. B 288, C 30, R 39. Bridled Titmouse.
- 44. Parus atricapillus L. B 290. C 31, R 41.

  Black-capped Chickadee
- 45. Parus atricapillus septentrionalis (Harr.) All. B 289. C 31a. R 41a. Long-tailed Chickadee.
- 46. Parus atricapillus occidentalis (Bd.) Coues. B 291. C 31c. R 41b. Western Chickadee.
- 47. Parus carolinensis Aud. B 293. C 31b. R 42. Carolina Chickadee.

[See Addenda, No. 879.

- 48. Parus montanus Gamb. B 294. C 32. R 40. Mountain Chickadee.
- 49. Parus hudsonicus Forst. B 296. C 33. R 45. Hudsonian Chickadee.

we instinctively incline to the latter, both as throwing the stress of voice on the radical syllable, instead of on the connecting vowel, and as the a in -phanes represents two vowels, ai or a as in phanomenon, phanogamous.

- 41. L. ĭn-ōr-nā'-tŭs. Lat. in, negative, and ornatus, ornate, adorned; orno, I ornament.
- 42. L. ā-trō-crīs-tā'-tŭs. Lat. ater, atra, atrum, black; and cristatus, crested; crista, a crest. Commonly written atricristatus; see Parus, No. 44.

N. B. — The tenability of the position taken by Dr. Coues (B. C. V., i, p. 117; 1878) respecting atro-cristatus has been queried by several correspondents; among them Mr. W. C. Avery, of Greensboro', Ala., who some time since furnished an extensive commentary on the names of the old Check List, and whose suggestions have often proved valuable. Mr. Avery maintains atricristatus, adducing albicerata (sc. ficus) from Pliny, 15, 18; and atri-, albi, magni-, &c., is undoubtedly a correct form of such compounds. But we take cristatus to be a perfect participle, and put ater in the ablative of instrument; there being no such word as atricristatus, unless we coin it. We consider the word equal to cristatus atro, conformably with usage in Picus albo-larvatus, Tyrannus aurantio-atrocristatus, &c. Compare also the actual Latin auro-clavatus, striped with golden.

- 43. L. wöll-web'-er-i. To --- Wollweber.
- 44. Pā'-rŭs ā-trǐ-căp-īl'-lŭs. Lat. parus, a titmouse; etymology in question, but apparently parus for parvus, small, petty, like the actual adverb parum, little; Gr. παῦρος, of same signification, th. παὑω; cf. pau-cus, pau-lus, pau-per, &c. Lat. atricapillus, black-hair(ed); capillus, hair of the head; a diminutive, allied to caput, and Gr. κεφαλή, the head. Compare English capillary, thready, hair-like, i. e., as fine as a hair. Notice atri-, not atro-; cf. Lophophanes, No. 42. If the compound were with capillatus, it would be atrocapillatus.
- 45. P. a. sep-ten-tri-o-na'-lis. Lat. septentrionalis, northern; septemtriones (septem and trio) being the constellation of seven stars near the north pole.
- 46. P. a. oc-cid-en-ta'-lis. Lat. occidentalis, western; occide, I fall; i. e., where the sun sets.
- 47. P. că-rŏl-ĭn-ēn'-sĭs. See Mimus, No. 16.
- 48. P. mon-ta/-nus. Lat. montanus, relating to a mountain; mons, genitive montis, a mountain.
- 49. P. hud-son/-i-cus. Latinized from the name of Henry Hudson, discoverer of the region.

- 50. Parus rufescens Towns. B 295. C 34. R 46. Chestnut-backed Chickadee.
- 51. Parus rufescens neglectus Ridg. B —. C —. R 46a. (?)
  Californian Chickadee,
- 52. Parus cinctus Bodd. B —. C —. R 44. (!A.) Siberian Chickadee.
- .53. Psaltriparus minimus (Towns.) Bp. B 298. C 35. R 47.
- 54. Psaltriparus plumbeus Bd. B 299. C 36. R 48. Plumbeus Bush-tit.
- 55. Psaltriparus melanotis (Hartl.) Bp. B 297. C —. R 49. (!M.)
  Black-eared Bush-tit.
- 56. Auriparus flaviceps (Sund.) Bd. B 300. C 37. R 50. Yellow-headed Verdin.
- 57. Sitta carolinensis Gm. B 277. C 38. R 51. White-bellied Nut-hatch.
- 58. Sitta carolinensis aculeata (Cass.) All. B 278. C 38a. R 51a. Slender-billed Nut-hatch.
- 50. P. rū-fēs'-cēns. Lat. rufescens, present participle of the inceptive verb rufesco, to grow red; be rufous.—"Chickadee" is an obvious onomatopœia, from the bird's note.
- 51. P. r. nēg-lēc'-tŭs. Lat. neglectus, neglected; verb negligo; equal to nec (non), not, and lectus, chosen, picked, taken; lego, I gather in, select, &c. Neglect is a nearly exact opposite of collect.
- 52. P. cinc'-tŭs. Lat. cinctus, girdled; perfect participle of cingo, I surround, encompass, encircle. A cingulum is a little something that goes around as a girdle does, whence surcingle, cinche.
- 53. P-sāl-trī-pā'-rūs mīn'-ĭ-mūs [sound the initial p; the a in parus is properly long; sometimes shortened in composition]. Lat. psaltria, Gr. ψάλτρια, one who plays on the lute; from the verb psallo, ψάλλω, to strike such an instrument; English psaltery, &c.; and parus, a titmouse. See No. 44. Lat. minimus, least, superlative of parvus, small.
- 54. P. plūm'-bĕ-ŭs. Lat. plumbeus, plumbeous, lead-colored.
- 55. P. měl-ăn-ō'-tīs. Gr. μέλαs, genitive μέλανος, black; οἶs, genitive ἀτός, ear. Not in the orig. ed. of the Check List, and scarcely established as North American, though given by Baird in 1858. Supposed to have been seen by Ridgway in Nevada, August, 1868. See Rep. Surv. 40th Par., iv, 1877, p. 415. See Index, p. 137.
- 56. Aūr-ī-pā'-rŭs [owriparus] flā'-vǐ-cēps. Lat. aureus, golden, from aurum, gold; and parus, a titmouse. Lat. flavus, yellow, for flagvus, from flagro, to glow; whence English deflagrate, flagrant, &c. Ceps is a Lat. termination, from Gr. κεφαλή, the head; compare caput, cephalic, occiput, &c. A more strict method of compounding aure-us with parus would give aureiparus; but it may be taken direct from aurum, making auriparus admissible; as we should say "gold-tit," like "bush-tit," "coal-tit."
- 57. Sit'-tă că-rŏ-lin-ēn'-sis. Gr. σίττα, σίττη; Lat. sitta, a nut-hatch; the word occurs in Aristotle. It is related to σιττάκη, ψίττακος, Lat. sittace, psittacus, a parrot; the implication being some sharp sound made by the bird, as English psit! There is a Greek verb ψιττάζω, to make such a noise. Lat. carolinensis, see Mimus, No. 16.
- 58. S. c. ă-cū-lĕ-ā'-tă. Lat. aculeatus, sharpened, dim. aculeus, sharp, acus, a needle; from acer, sharp. Gr. ἀκίs, a point; compare ἀκών, ἀκμή, ἄκροs, &c., English acme, acropolis, acerbity, acrimony, and numberless words in many languages, from √ak.

- 59. Sitta canadensis L. B 279. C. 39. R 52. Red-bellied Nut-hatch.
- 60. Sitta pusilla Lath. B 280. C. 40. R 53. Brown-headed Nut-hatch.
- 61. Sitta pygmæa Vig. B 281. C 41. R 54. Pygmy Nut-hatch.
- 62. Certhia familiaris L. B 275. C 42. R 55. Brown Creeper.
- 63. Campylorhynchus brunneicapillus (Lafr.) Gr. B 262. C 43. R 56. Brown-headed Cactus Wren.
- 64. Campylorhynchus affinis Bd. B —. C 44. R 57. St. Lucas Cactus Wren.
- 65. Salpinctes obsoletus (Say) Cab. B 264. C 45. R 58. Rock Wren.
- 66. Catherpes mexicanus (Sw.) Bd. B 263. C —. R 59. (!M) Cañon Wren.
- 59. S. că-năd-ēn'-sĭs. Latinized from Canadian. Nut-hatch is nut-hatcher or nut-hacker (Fr. hacher, Swed. hacka), the bird that hacks, pecks, nuts; also called nut-jobber, to job being to peck, or thrust at.
- 60. S. pŭ-sīl'-lă [puceellah, not pewziller]. Lat. pusillus, petty, puerile; directly formed from puer, pusus, or pusio (Gr. παῖs), a boy; here and commonly used simply as signifying small. The Sanskrit root reappears in endless forms of kindred meaning.
- 61. S. pyg-maē'-ā. Gr. πυγμή, the fist; hence πυγμαῖοs, Lat. pygmæus, a pygmy, fistling, or tom-thumb. As a measure of length, from elbow to clenched fist, a πυγμή was about 13½ inches; the original Pygmies were a race of African dwarfs at war with the Cranes; pygmæus came afterward to mean any thing pygmy, dwarfed, and is here applied to a very small nut-hatch. Compare Machetes pugnax, No. 639.
- 62. Cēr'-thǐ-ă făm-ĭl-i-ā'-rĭs. Gr. κέρθιος, Lat. certhius, become later certhia. The name occurs in Aristotle, who apparently uses it for this very species, which he also calls κυιπολόγος, cnipologus; that is to say, a gatherer of insects; κυίψ, a bug, and λέγω, I collect.—Lat. familiaris, familiar, domestic, hence common; familia, or older familias, the family, the household.
- 63. Cām-pỹ-lŏ-rhỹn/-chus brūn-neī-căp-īl/-lus [broonaycapeellus]. Gr. καμπύλοs, bent, from κάμπτω, I bend; and ῥύγχοs (rhynchus), beak. Lat. brunneus, brown; capillus, hair. The adjective brunneus is post-classic, Latinized from It. bruno, Fr. brun, Germ. braun; A. S. byrnan, to burn; related are brand, brunt, and many similar words, among them brant; see Bernicla, No. 700.
- 64. C. āf-fīn'-ĭs [affeen'is]. Lat. affinis, i. e., ad and finis, at the end of, hence bordering on, neighboring; here in the sense of related to, resembling, having affinity with, No. 63.
- 65. Sāl-pinc/-tēs ōb-sŏ-lē/-tŭs. Gr. σαλπιγκτήs, a trumpeter, becoming in Latin salpinctes, from σάλπιγξ (salpigx = salpinx), a trumpet; in allusion to the bird's loud, ringing song.— Lat. obsoletus, unaccustomed, from ob, against, and soleo, I am wont; hence obsolete, in sense of effaced, all the colors of the bird being dull. Wren is A. S. wrenna.
- 66. Căth-ēr'-pēs mēx-ĭ-cā'-nŭs. Gr. καθερπήs, a creeper; καθέρπω, I creep down, from κατά, down, and ἕρπω, I creep, crawl. The stem of the word is seen in herpes, the disease which creeps over the skin; herpetology, the science of creeping things, reptiles; repto or repo, I creep, in Latin, simply altered from ἕρπω. Lat. mexicanus, see No. 28.

- 67. Catherpes mexicanus conspersus Ridg. B —. C 46. R 59a. Speckled Canon Wren.
- 68. Thryothorus ludovicianus (Gm.) Bp. B 265. C 47. R 60. Carolina Wren.
- 69. Thryothorus ludovicianus miamiensis Ridg. B —. C —. R 60b. Floridan Wren.
- 70. Thryothorus ludovicianus berlandieri (Couch) Coues. B 266. C 47a, R 60a. Texan Wren.
- 71. Thryothorus bewicki (Aud.) Bp. B 267. C 48. R 61. Bewick's Wren.
- 72. Thryothorus bewicki leucogaster Bd. B —. C 48a. R 61b. White-bellied Wren.
- 73. Thryothorus bewicki spilurus (Vig.) Bd. B —. C 48b. R 61a. (?) Speckled-tailed Wren.
- 74. Troglodytes domesticus (Bartr.) Coues. B 270, 272. C 49. R 63. House Wren.
- 75. Troglodytes domesticus parkmani (Aud.) Coues. B 271. C 49a. R 63a. Western House Wren.
- 67. C. m. con-spēr'-sus. Lat. conspersus, speckled; perfect participle of conspergo, from con and spargo (Gr. σπείρω), I strew, scatter, sprinkle; whence English sparse, scattered, and many other words, as disperse, aspersion. The Span. cañon, brutalized as Eng. canyon, is constantly used in the West for a rocky gorge or mountain-pass.
- 68. Thrỹ-ŏ-thō'-rŭs lū-dō-vǐ-cǐ-ā'-nŭs. Gr. θρύον, a reed, rush, and θοῦρος, a leaping, springing, from (θόρω), θρώσκω, I run or rush through. The penult is marked long, as equivalent to Gr. οῦ. Lat. Ludoviciana, Louisiana, of or relating to Ludovicus, Louis (XIV., of France). The old Territory was vastly more extensive than the present State is.
- 69. T. 1. mi-ă-mi-ēn'-sis. Latinized from the name of the Miami river in Florida.
- 70. T. l. ber-lan'-dī-er-ī. To Dr. Louis Berlandier, a naturalist, sometime resident in Mexico.
- 71. T. bē'-wick-i. To Thomas Bewick, "the father of wood-engraving."
- 72. T. b. leū-cŏ-gās'-tēr [lewco-]. Gr. λευκόs, white, and γαστήρ, stomach, belly; whence English gastric, gastronomy.
- 73. T. b. spǐl-ū'-rŭs. Gr.  $\sigma\pi i\lambda \sigma s$ , spotted;  $\sigma \partial \rho \alpha$ , tail.
- 74. Trög-lð'-dỹ-tēs [-tace] dŏm-ēs'-tǐ-cŭs. Gr. τρωγλοδύτης, a cave-dweller, from τρώγλη, a cave (literally, a hole made by gnawing τρώγω, I gnaw), and δύτης, an inhabitant, from δύνω or δύω, I go in or under. The Τρωγλοδύται or Troglodytæ were a cave-dwelling people of Æthiopia. The name was later applied to a kind of wren. Lat. domesticus, domestic, from domus, a house. The specific name aëdon, applied by Vicillot to this bird, is the Gr. ἀηδών, a songster, par excellence the nightingale; from ἀείδω, I sing. The pronunciation of Troglodytes wavers; we mark it as commonly heard, and also as seems to be defensible, in Latin, the penult being indubitably short; though to do so violates one of the leading principles of Greek accentuation, that no word with the ultimate long is a proparoxytone. Many persons say Trog'lodỹ'tes, conformably with English Trog'lodye''. The case is precisely parallel with that of Lopho'phānēs, q. v., No. 40; and the analogy of Aristo'phānes is not decisive, the Greek being 'Αριστοφάνης or 'Αριστοφανής, not 'Αριστοφανής.
- T. d. pārk'-măn-ī. To Dr. George Parkman, of Boston, murdered by Professor John W. Webster, in 1849.

- 76. Anorthura troglodytes hiemalis (Wils.) Coues. B 273. C 50. R 65. Winter Wren.
- 77. Anorthura troglodytes pacificus (Bd.) Ridg. B —, C —, R 65a. (?) Western Winter Wren.
- 78. Anorthura troglodytes alascensis (Bd.) Coues. B —. C 50a. R 66.
  Alaskan Winter Wren.
- -79. Telmatodytes palustris (Bartr.) Cab. B 268. C 51. R 67. Long-billed Marsh Wren.
- 80. Telmatodytes palustris paludicola Bd. B —. C —. R 67a. (?)
  Tule Marsh Wren.
- 81. Cistothorus stellaris (Licht.) Cab. B 269. C 52. R 68. Short-billed Marsh Wren.
- 82. Eremophila alpestris (L.) Boie. B 302. C 53. R 300. Horned Lark; Shore Lark.
- 83. Eremophila alpestris leucolæma Coues. B C 53b. R 300a. (?)
  Western Shore Lark.
  - 84. Eremophila alpestris chrysolæma (Wagl.) Bd. B —. C 53a. R 300c.
  - 76. Ăn-ōr-thū'-ră trōg-lŏ'-dỹ-tēs hĭ-ĕm-ā'-līs. Gr. ἀ or ἀν, privative, ὀρθόs, straight, οὖρα, tail. The name was invented by Rennie, because he considered Troglodytes etymologically inapplicable to a wren. Lat. hiemalis or hyemalis, of or pertaining to winter; from (hiemps) hiems or hyems, winter, a weakened form of the Gr. χεῦμα, a gushing, a torrent, or χεῦμάν, the rainy, tempestuous, or winter season; Skr. hima, snow. We oftener use the y than the i, but the latter is correct.
  - 77. A. t. pā-ci'-fi-cus. Lat. pacificus, pacific, peaceful, literally peace-making, from pax, genitive pacis, peace, and facio, I make, do. The application is to the occurrence of the bird on the west coast of the United States.
    Not in the orig. ed. of the Check List. (Baird, Rev. Am. B., i, 1864, p. 145.)
  - 78. A. t. ă-lās-cēn'-sis. Alascensis, relating to Alaska.
  - 79. Tēl-mă-tŏ'-dỹ-tēs păl-ūs'-tris. Gr. τέλμα, genitive τέλματοs, a marsh or swamp; δύτηs, an inhabitant, from δύω, I go in or under. Lat. palustris, adjective from palus, a marsh, whence palustrine, like lacustrine from lacus, marine from mare.
  - 80. T. p. păl-ū-dĭ'-cŏ-lă. Lat. palus, genitive paludis, a marsh; and (in)cola, an inhabitant. See No. 79.
    - Not in the orig. ed. of the Check List. (Baird, Rev. Am. B., i, 1864, p. 148.)
  - 81. Cis-tŏ-thō'-rŭs stēl-lā'-rĭs. Gr. κίστος, a shrub, and θοῦρος, from (θόρω) θρώσκω, I run or rush through; compare Thryothorus, No. 68. Cabanis, who coined the word in 1850, gives Ciflĕdiüpfer as the German translation. Lat. stellaris, stellar, starry, adjective from stella, a star, like aster, Gr. ἀστήρ; here in the sense of speckled.
  - 82. Ĕr-ē-mŏ'-phǐ-lă āl-pēs'-trǐs. Gr. ἐρῆμος, a desert; φιλέω, I love.—Lat. Alpestris (not classic), from Alpes, Alps; perhaps from ἀλφός, albus, white; that is, snowy.
  - 83. E. a. leū-cŏ-laē'-mā. Gr. λευκόs, white; λαιμόs, the throat. This is a slight variety, lately described by Coues from the high central plains; it is the bleached form of that region. (B. N. W., 1874, p. 38.)
  - 84. E. a. chrỹ-sŏ-laē'-mă. Gr. χρύσεος, golden; that is, of a golden color, from χρυσός, gold; and λαιμός, the throat. Λ. S. laferc, Scot. laverock, Germ. lerde, Eng. lark.

- 85. Alauda arvensis L. B —. C —. R 299. (!E.) European Skylark.
- 86. Motacilla alba L. B —. C —. R 69. (G.) White Wagtail.
- 87. Budytes flavus (L.) Gr. B —. C 54. R 70. Yellow Wagtail.
- 88. Anthus pratensis (L.) Bechst. B —. C 55bis. R 72. (!E.) Meadow Pipit.
- 89. Anthus ludovicianus (Gm.) Licht. B 165. C 55. R 71. Louisiana Pipit; American Titlark.
- 90. Neocorys spraguii (Aud.) Scl. B 166. C 56. R 73. Missouri Skylark; Sprague's Pipit.
- 85. A-laū'-dă ār-vēn'-sīs. Lat. alauda, a lark, said to be literally "a great songstress," or one who sings on high; from the Celtic al, great, high, and aud, song. Some say from Gr. ἄλη, roaming, and ἀδή, song; i. e. the bird that sings as she soars. The form of the word might suggest ala, wing, and laus, genitive laudis, praise; as if the bird sang praises on wing. But the Celtic is the only tenable etymon.—Lat. arvensis, relating to a ploughed field; arvum, arable land; arvus for aruus, ploughed; aro, I plough.

  Not in the orig. ed. Said to occur in Greenland, Alaska, and the Bermudas.

86. Mō-tă-cīl'-lă āl'-bă. Lat. motacilla = wagtail, "quod semper caudam movet," carly applied

to some small bird; Lat. moveo, motus, I move, motion, and Gr. κίλλω of similar signification. There is a Greek word κίλλουρος, for the wagtail; on the contrary there are the Lat. albicilla, atricilla, meaning white-tail, black-tail, &c. The implication in either case seems to be tail, considered as a movable part. Compare Fr. hochequeue.

Not in the orig. ed. The species is North American only as occurring in Greenland.

87. 'Bū'-dy-tēs flā'-vŭs. Budytes is an unknown word to us, unless conjectured to be δύτης, with the augmentative particle βου. See Troglodytes, No. 74. The particle βου, however, is from βοῦς, a bull, ox, cow, and becomes "augmentative," just as we say "horse-laugh," "bull-finch," "elephant-folio," &c., being therefore of obvious inapplicability to this delicate little bird.

Since the above was written, Mr. Henry T. Wharton, of London, has kindly replied to queries respecting various words of which we were in doubt. In this case, his MS. confirms the above etymology, but in a different application; the actual form, βουδύτης, being found in "Opp., Ix. 3. 2," for some small bird; qu., one that goes among cattle?

There is some question whether the yellow wagtail of Alaska be the true B. flavus.

88. An'-thus prā-tēn'-sis. Lat. anthus, Gr. ἄνθος, a kind of bird. — Lat. pratensis, adjective from pratum, a meadow. For anthus, compare αnanthe = vitiflora, under Saxicola, No. 26.

This is North American as found in Greenland, and said to also occur in Alaska.

- 89. A. lu-dō-vī-cī-a'-nūs. Lat. Ludovicus, nom. prop. See Thryothorus, No. 68. Pipit, little used in this country, though always said for these birds in England, is an onomatopæia (δνοματοποιία, word-making to express the sense by the sound), like the Lat. pipio, I pip, peep, chirp; see Pipilo, No. 301. Titlark is good English for a small kind of lark, like tit-mouse, tom-tit; tit in all its forms, and with numerous related words, conveying the sense of something little or otherwise insignificant.
- 90. Ně-ŏ'-cŏ-rÿs sprā'-guĭ-ī [three syllables]. Gr. νέος, new; κόρυς, primarily a helmet; hence applied to the crested lark.—To Isaac Sprague, companion of Audubon on the Missouri.

- '91. Mniotilta varia (L.) V. B 167. C 57. R 74. Black-and-white Creeper.
- 92. Mniotilta varia borealis (Nutt.) Ridg. B —. C —. R 74a. (?)
  Small-billed Greeper.
- 93. Parula americana (L.) Bp. B 168. C 58. R 88. Blue Yellow-backed Warbler.
- 94. Parula nigrilora Coues. B.—. C.—. R 89a. Sennett's Warbler.
- 95. Protonotaria citrea (Gm.) Bd. B 169. C 59. R 75. Prothonotary Warbler.
- 96. Helmintherus vermivorus (Gm.) Bp. B 178. C 60. R 77. Worm-eating Warbler.
- 91. Mnĭ-ŏ-tīl'-tă văr'-ĭ-ă. Gr. μνίον, moss, and τίλλω, I pluck, or τιλτόs, plucked. Neither the orthography nor the applicability of the word is obvious. Vieillot wrote sometimes mniotilta, sometimes mniotilta. The conjectured application is to the weaving of moss into a nest. Lat. varia, variegated, as this bird is with black and white.
- 92. M. v. bŏr-ĕ-ā'-līs. Lat. borealis, northern. See Phylloscopus, No. 32.

  Not in the orig. ed. of the Check List.
- 93. Pā'-rǔ-lǎ ǎm-ĕr-ĭ-cā'-nǎ. Lat. parula, diminutive from parus, a titmouse, q. v., No. 44.

   Lat. americana, American. America is generally supposed to derive its name from Amerigo Vespucci, Latinized Americus Vespucius; and is said to have first appeared in the form of America Provincia, on a map published at Bâsle in 1522. The counter-argument is: (1) The name if from the Italian navigator's would have been from his surname. (2) His name was Alberico Vespuzio. (3) Americ, or Amerique, is the native name of a range of mountains in Nicaragua. "It is most plausible that the State of Central America, where we find the name Americ signifying great mountain, gave the continent its name." (Blavatsky, Isis Unveiled, i, p. 592.) The author cited seeks to establish a connection with the Hindu Meru, or Meruah, of similar signification.
- 94. "P. nīg-rī-lō'-rā. Lat. niger, black; and lorum, a thong, strap, a bridle-rein; hence the cheeks, along which the bridle passes. The "lore" has become in ornithology a technical name for a small space on the side of a bird's head between the eye and the bill.

  Not in the first ed. of the Check List. Lately discovered in Texas by Mr. George B. Sennett. See Coues, Bull. U. S. Geol. Surv. Terr., iv, 1878, p. 11.
- 95. Prō-tŏ-nŏ-tā'-rĭ-ă cīt'-rĕ-ă. Low Latin for prothonotary; from Gr. πρώτοs, first, and Lat. notarius, a scribe, a notary-public. The bird is le Protonotaire of Buffon, Latinized by Gmelin as protonotarius in 1788; but for the name, as Pennant observed in 1785, "the reason has not reached us."—Lat. citrea, of or pertaining to the citron, in allusion to the yellow color.
- 96. Hēl-mīn-thē'-rŭs vēr-mī'-vŏr-ŭs. Gr. ἔλμινs, genitive ἕλμινθοs, and θήριον, from θήρ, an animal. The word is very incorrectly compounded. Its full form is helminthotherium; we may perhaps reduce it by elision to helmintherus, but helmitherus, as originally written by Rafinesque, is inadmissible. This is the accepted derivation; but we may suggest a short cut to the same etymon, θήρ, an animal; ἐλμινθοθήραs, a worm-hunter, like the actual ὀρνιθοθήραs, a fowler, in Aristoph., Av. 62; being ἕλμινs and θήρα, the chase, from θήρ; though we hesitate to act upon this by writing Helmintheras.—Lat. vermivorus, worm-eating, from vermis, a worm (verto, I turn, in the sense of squirming or wriggling) and voro, I eat.

- .97. Helmintherus swainsoni (Aud.) Bp. B 179. C 61. R 76. Swainson's Warbler.
- 98. Helminthophaga pinus (L.) Bd. B 180. C 62. R 79. Blue-winged Yellow Warbler.
- 99. Helminthophaga lawrencii Herrick. B—, C—, R 80. (?) Lawrence's Warbler.
- 100. Helminthophaga leucobronchialis Brewster. B —. C —. R 82. (?)
  White-throated Warbler.
- 101. Helminthophaga cincinnatiensis Langdon. B —. C —. R —. (?)
  Cincinnati Warbler.
- 102. Helminthophaga chrysoptera (L.) Bd. B 181, C 63, R 81. Blue Golden-winged Warbler.
- 103. Helminthophaga bachmani (Aud.) Cab. B 182. C 64. R 78. Bachman's Warbler.
- 104. Helminthophaga luciæ Coop. B —. C 65. R 83. Lucy's Warbler.
- 105. Helminthophaga virginiæ Bd. B —. C 66. R 84. Virginia's Warbler.

owam 1 1110= Tode ....

- 106. Helminthophaga ruficapilla (Wils.) Bd. B 183. C 67. R 85. Nashville Warbler.
  - 97. H. swāin'-sŏn-ī. To Wm. Swainson, Esq., the celebrated English naturalist. Notice that this word, like others containing the letter w, cannot be Latinized without change; the nearest Latin would be suā-īn'-sŏn-ī, in four syllables. See also lawrencii, next but one below; this should be laū-rēn'-cī-ī or lāv-rēn'-cī-ī. But it is futile, finical, and pedantic to undertake such transliterations in the cases of modern proper names.
  - 98. Hēl-mīn-thờ'-pha gà pĩ'-nus. Gr. ελμις, a worm, and φαγείν, to eat. Lat. pinus, Gr. πίτυς, a pine-tree. Notice that pinus is a substantive, not an adjective; it may be put in the genitive, pinûs, of a pine, but is just as well left nominative.
  - 99. H. lăw-rĕn'-cĭ-ī. To George N. Lawrence, Esq., of New York, long time one of the leading ornithologists of America. Not in orig. ed. Since described, Pr. Phila. Acad., 1874, p. 220, pl xv.
  - 100. H. leū-cŏ-brōn-chī-ā/-līs. Gr. λευκόs, white, and βρόγχοs, the throat; this becomes in Latin bronchus, whence the adjective bronchialis, English bronchial, bronchitis, &c. Not in orig. ed. Since described, Bull. Nutt. Club, i, 1876, p. 1, pl.
  - 101. H. cīn-cīn-nă-tǐ-ēn'-sīs. Of Cincinnati, Ohio, where discovered.
    Not in the orig. ed. Lately described by F. W. Langdon, in Journ. Cinc. Soc. Nat.
    Hist., ii, July, 1880, p. 119, and Bull. Nuttall Club, v, October, 1880, p. 208, pl. iv.
- 102, H. chry-sop'-tě-ră. Gr. χρυσόπτερος, golden-winged, from χρυσός, gold, and πτερόν, wing.
- 103. H. băch'-măn-ī. To Rev. John Bachman, D.D., of Charleston, S. C., collaborator with Audubon in the "Quadrupeds of North America."
- 104. H. lū'-cĭ-aē. To Miss Lucy Baird, daughter of Professor S. F. Baird.
- 105. H. vīr-gĭn'-ĭ-aē. To Mrs. Virginia Anderson, wife of Dr. W. W. Anderson, who discovered the bird.
- 106. H. rū-fī-căp-īl'-lă. Lat. rufus, reddish, and capillus, hair of the head. See Parus, No. 44.

- 107. Helminthophaga celata (Say) Bd. B 184. C 68. R 86. Orange-crowned Warbler.
- 108. Helminthophaga celata lutescens Ridg. B —. C 68a. R 86a. Pacific Orange-crowned Warbler.
- 109. Helminthophaga peregrina (Wils.) Cab. B 185. C 69. R 87. Tennessee Warbler.
- 110. Peucedramus olivaceus (Gir.) Coues. B —. C —. R 92. Olive Warbler.
- 111. Dendræca æstiva (Gm.) Bd. B 203. C 70. R 93. Summer Warbler.
- 112. Dendræca virens (Gm.) Bd. B 189. C 71. R 107.
  Black-throated Green Warbler.
- 113. Dendræca occidentalis (Towns.) Bd. B 190. C 72. R 109. Western Warbler.
- 114. Dendræca townsendi (Nutt.) Bd. B 191. C 73. R 108. Townsend's Warbler.
- 107. H. ce-la'-ta. Lat. celatus, concealed, from celo; the orange color of the crown being hidden.
- 108. H. c. lū-tēs'-cēns. Lat inceptive verb lūtesco, present participle lūtescens, from lūteus, yellow; from lūteum, an herb used in dyeing yellow. There is actually no such verb as lūtesco, the describer of the species having apparently mistaken lūtesco, I grow muddy, become miry, for a supposed lūtesco, I grow yellow, by some confounding of lūteus, muddy, loamy (hence possibly clay-colored or yellowish) with lūteus, golden-yellow. The bright yellowness of the bird in comparison with H. celatu being its prime characteristic, the propriety of assuming the derivation to be from lūtum, and hence writing lūtescens, from a supposed lūtesco, is obvious.

A form lately distinguished by Ridgway, Am. Journ. Sci., 3d ser., iv, 1872, p. 457.

- 109. H. pěr-ě-grī'-nă. Lat. peregrinus, wandering, alien, exotic, that comes from foreign parts; from per, through, and ager, a field or land; literally, "across country."
- 110. Peū-cē'-dră-mŭs ŏl-ī-vā'-cĕ-ŭs. Gr. πεὐκη, a pine-tree, and δραμεῖν, 2d aorist infinitive, from τρέχω, I run. The allusion is to the pine-creeping habits of the bird. N. B. Many genera are compounded from the same root, and spelled either-dramus or dromus. Either is correct. Lat. olivaceus, pertaining to the olive; in this case, in color, olivaceous.
  Not in the orig. ed. Since discovered in Arizona by H. W. Henshaw.
- 111. Dēn-droē'-că aēs-tī'-vă [dayndrwaykah aysteevah]. Gr. δένδρον, a tree, and οἰκέω, I inhabit; οἶκος, a habitation. The word was originally compounded Dendroica by G. R. Gray: later emended as above. The full form would be Dendrocetes, like Poacetes, Nephacetes (οἰκητής, an inhabitant). Lat. æstira, adjective from æstas, the summer season; æstus, heat, ardor (Gr. αἴθω, I burn). Notice the long accented penult.
- 112. D. vĭr'-ēns [pronounced virraynce]. Lat. virens, participle present of vireo, I grow green.
- 113. D. ōc-cĭ-dēn-tā'-līs. Lat. occidentalis, occidental, western; that is, in the place where the sun sets; from occido, I fall down.
- 114. D. town'-sĕnd-ī. To J. K. Townsend, Esq., companion of Nuttall during his travels. The first syllable of this word represents the exact pronunciation of Latin au diphthong—like English ow; as if we made it tă-ūn-.

- 115. Dendræca chrysoparia Scl. & Salv. B —, C 74. R 106. Golden-cheeked Warbler.
- 116. Dendræca nigrescens (Towns.) Bd. B 192. C 75. R 105.
  Black-throated Gray Warbler.
- 117. Dendræca cærulescens (L.) Bd. B 193. C 76. R 94.
  Black-throated Blue Warbler.
- 118. Dendræca cærulea (Wils.) Bd. B 201. C 77 R 98 Coerulean Warbler.
- / 119. Dendrœca coronata (L.) Gr. B 194. C 78. R 95. Yellow-rumped Warbler.
  - ,120. Dendræca auduboni (Towns.) Bd. B 195. C 79. R 96. Audubon's Warbler.
- /121. Dendræca blackburnæ (Gm.) Bd. B 196. C 80. R 102. Blackburn's Warbler.
  - 122. Dendræca striata (Forst.) Bd. B 202. C 81. R 101. Black-poll Warbler.
  - 123. Dendræca castanea (Wils.) Bd. B 197. C 82. R 100. Bay-breasted Warbler.
    - 115. D. chry-sŏ-păr-ī'-ă. Gr. χρυσόs, gold, and παρειά, cheek. Greek diphthong ει becomes long i in Latin: hence, -parīa, not -pareia; see also beyond, among the names of pigeons ending in -pelīa.
    - 116. D. nĭg-rēs'-cēns. Lat. nigresco, I grow black; an inceptive verb, present participle nigrescens, equivalent to being blackish, or partly black. See No. 126.
    - 117. D. coē-rŭl-ēs'-cēns [pronounced sayrullaysaynce]. Lat. cœrulesco, I grow blue; a coined inceptive verb from cœruleus, blue; this from cœlum, the (blue) sky; compare Gr. κοίλος, hollow, i. e., the vault of heaven, and cœlare or celare, to conceal, as if in a hollow place, &c. N. B. There is constant difference of orthography: either cœ- or cæ- is defensible; the former seems preferable. In English we may write indifferently cœrulean, cœrulean, or cerulean.
    - 118. D. coē-rŭl'-ĕ-ă. See last word.
    - 119. D. cŏr-ō-nā'-tă. Lat. coronatus, crowned, from corona, a crown, garland, or wreath. Gr. κορώνη.
    - 120. D. aūd'-ŭ-bŏn-ī. To John James Audubon, "the American backwoodsman," as he liked to be called.
    - 121. D. black'-burn-aë. To Mrs. Blackburn, an English lady. Commonly written blackburniæ, in four syllables, with accent on the antepenult; more correctly as above. Diacritical marks are futile in such a case as this; the English name is never pronounced blarkboorn, as it would be according to rule for the quantity of the vowels in Latin.
    - 122. D. strī-ā'-tă. Lat. participial adjective from strio, I furrow, channel, flute, groove, striate, stripe; stria, substantive, a furrow, stripe, &c.
    - 123. D. cas-tăn'-ĕ-ă. Lat. castanea, a chestnut; in allusion to the bay or chestnut color. The word is a noun, but is constantly used adjectivally. Gr. κάστανον, the nut of Castana, a city of Thessaly.

- 124. Dendrœca pennsylvanica (L.) Bd. B 200. C 83. R 99. Chestnut-sided Warbler.
- 125. Dendræca maculosa (Gm.) Bd. B 204. C 84. R 97. Black-and-Yellow Warbler.
- 126. Dendræca tigrina (Gm.) Bd. B 206. C 85. R 90. Cape May Warbler.
- 127. Dendræca discolor (V.) Bd. B 210. C 86. R 114.
  Prairie Warbler.
- 128. Dendræca graciæ Coues. B —, C 87. R 104. Grace's Warbler.
- 129. Dendrœca dominica (L.) Bd. B 209. C 88. R 103. Yellow-throated Warbler.
- 130. Dendræca dominica albilora Bd. B —. C 88a. R 103a. White-cheeked Warbler.
  - 131. Dendræca kirtlandi Bd. B 205. C 89. R 110. Kirtland's Warbler.
  - 132. Dendræca palmarum (Gm.) Bd. B 208, C 90. R 113. Yellow Red-poll Warbler.
- 124. D. pěnn-sỹl-vā'-nĭ-că. An adjective coined from sylvanus, sylvan, this from sylva, which is sibilated and digammated from Gr. ὕλη = (σ)υλ Fη, a wood; preceded by the name of William Penn; "Penn's woods." The modern use of the y is less correct than i would be. The whole word would preferably be written pensilvanica, as it is in some ornithological works of the last century.
  - 125. D. mă-cŭl-ō'-să. Lat. maculosus, spotted or full of spots; macula, a spot.
  - 126. D. tig-ri'-nă. Lat. tigrinus, striped (like a tiger, tigris, Gr. τίγριs). The quantity of the antepenult is doubtful, perhaps common. By ordinary rule, it is long, and Tigris makes the final spondee of some hexameter lines. On the other hand, the combination of a mute or f and a liquid does not necessarily lengthen a preceding vowel in prose; and some other combinations of consonants also permit the vowel to remain short, in cases of Greek words, as Cýcnus or Cýgnus. We leave it short, as usually heard. Perissoglossa, a generic name now often used for this species, is the Gr. περισσόs and γλῶσσα, in allusion to the peculiarity of the laciniate tongue.
  - 127. D. dis'-cŏl-ŏr. Lat. discolor (post-classic), party-colored; opposed to concolor, whole-colored.
  - 128. D. gră'-cĭ-aē. To Mrs. Charles A. Page, née Grace Darling Coues, the author's sister. Would more strictly be written gratiæ (Lat. gratia, grace, favor, thanks).
  - 129. D. dom-in'-i-ca. Lat. dominicus, relating to the lord or master of the household, dominus; domus, a house. So, to dominate, to have dominion. The application is here to the West Indian island named originally Hayti, then San Domingo. The bird was early described from that locality.
  - 130. D. d. āl-bǐ-lō'-ră. Lat. albus, white, and lorum, the lore or cheek. See Parula, No. 94.
  - 131. D. kirt'-land-ī. To Dr. Jared P. Kirtland, of Ohio. See remarks under D. blackburnæ.
  - 132. D. pāl-mā'-rum. Lat. palmarum, of the palms, genitive plural of palma, a palm.

- 133. Dendræca palmarum hypochrysea Ridg. B —. C —. R 113a. (?)
  Yellow-bellied Red-poll Warbler.
- 2134. Dendrœca pinus (Bartr.) Bd. B 198. C 91. R 111.
  Pine-creeping Warbler.
- > 135. Siurus auricapillus (L.) Sw. B 186. C 92. R 115. Golden-crowned Thrush.
- · 136. Siurus nævius (Bodd.) Coues. B 187. C 93. R 116. Water Thrush.
- 137. Siurus nævius notabilis Grinnell. B —. C —. R 116a. (?)
  Wyoming Water Thrush.
- Large-billed Water Thrush.
- ,139. Oporornis agilis (Wils.) Bd. B 174. C 95. R 118. Connecticut Warbler.
- 140. Oporornis formosa (Wils.) Bd. B 175. C 96. R 119. Kentucky Warbler.
- 133. D. p. hỹ-pō-chrỹ/-sĕ-ă. Gr. ὁπό, becoming Lat. hypo-, under, below, beneath, and χρόσεος, golden; referring to the under parts of this variety, which are yellower than those of palmarum. Properly, hypo- in such connection simply diminishes the force of the adjective; hypoleucus, hypochryseus, meaning whitish, yellowish; but the present is an established usage in ornithology.

Not in the orig. ed. - Since described by Ridgway, Bull. Nutt. Club, i, 1876, p. 84.

- 134. D. pī'-nŭs. See Helminthophaga pinus, No. 98.
- 135. Sī-ū'-rūs aūr-ĭ-căp-īl'-lūs. Gr. σείω, I wave or brandish, and οἶρα, tail. The word is precisely equivalent to Lat. motacilla, French hochequeue, English wagtail. It was originally and has since commonly been written Seiurus. (See Coues, Bull. Nuttall Club, ii, no. 2, 1877, p. 29.) We keep the i long as representing Gr. ει. Lat. aurum, gold, and capillus, hair; golden-haired. (See Coues, ibid., p. 30.) See also Lophophanes, No. 42, and Parus, No. 44.
- 136. S. naē'-vĭ-ŭs. Lat. nævus, a birth-mark, nevus, or spot; whence nævius, so marked, or, in general, spotted in any way.
- 137. S. n. nŏ-tā/-bĭ-lĭs. Lat. notabilis, notable, from nota, a note, and the termination -bilis. Not in the orig. ed. of the Check List. Later described by R. Ridgway, from Grinnell's MS., in Pr. Nat. Mus., ii, 1880, p. 12. Very doubtful.
- 138. S. mō-tă-cīl'-lă. See Motacilla, No. 86.
- 139. Ŏp-ōr-ōr'-nĭs ă'-gĭ-lĭs. Gr. ὀπώρα, the autumn, and ὅρνις, a bird; in allusion to the abundance of the species in the fall, in comparison with its scarcity in the spring. Lat. agilis, agile, from ago, I act; literally, do-able, that is, act-ive; the adjectival termination being simply applied to the root of the verb, both in Latin and English.
- 140. O. fōr-mō'-să. Lat. formosa, beautiful; primitively, in the sense of shapely, well-formed, in good or full proportion; forma, form. So said of Juno, in whose "lofty mind" remained judicium Paridis, spretæque injuria formæ, h. e., of her slighted beauty. Verg., Æn., i, 27.

- 141. Geothlypis trichas (L.) Cab. B 170. C 97. R 122.

  Maryland Yellow-throat.
- 142. Geothlypis philadelphia (Wils.) Bd. B 172. C 98. R 120. Mourning Warbler.
- 143. Geothlypis maegillivrayi (Aud.) Bd. B 173. C 99. R 121.

  Macgillivray's Warbler.
- 144. Icteria virens (L.) Bd. B 176, C 100. R 123. Yellow-breasted Chat.
- 145. Icteria virens longicauda (Lawr.) Coues. B 177. C 100a. R 123a. Long-tailed Chat.
  - 146. Myiodioctes mitratus (Gm.) Aud. B 211. C 101. R 124. Hooded Flycatching Warbler.
  - 141. Gĕ-ōth'-ly-pīs trīch'-ās. Gr.  $\gamma \hat{\eta}$  or  $\gamma \acute{\epsilon} \alpha$ , the earth, and  $\Theta \lambda \nu \pi \ell s$ , "a proper name." Gr.  $\theta \rho \ell \xi$ , genitive τριχός, hair; there is also the actual word τριχάς, for some kind of a thrush, occurring in Aristotle. Some take the τριχάς of Aristotle to be the bird named by Linnæus Turdus pilaris, i.e., the hairy thrush; but Sundevall reasonably identifies it with T. musicus. Of course it had originally nothing to do with the present species, to which Linnæus applied the term trichas in 1766. - Cabanis coined Geothlypis in 1847, simply explaining  $\Theta \lambda \nu \pi i s$  as a "proper name." The meaning of the term is obscure, but we think it may be explained, considering that  $\theta \lambda \nu \pi is$  is the same as  $\theta \rho \alpha \nu \pi is$ , which latter occurs in Aristotle as the name of some conirostral granivorous bird never satisfactorily identified. Sundevall says θραυπίς ,, würde in einigen codices Θλυπίς (Thlypis) acfdrieben"; and the identity of the two words appears to be established, seeing that  $\theta \lambda \dot{\alpha} \omega$ , I break, bruise, crush, whence  $\theta \lambda \alpha \upsilon \pi i s$ , has the same meaning as  $\theta \rho \alpha \dot{\omega} \omega$ , whence θραυπίς. (See Aristoph., Av. 466.) In each case the name is that of a bird considered as granivorous - as a seed-eater, i. e., seed-breaker, samenfressender, coccothraustes, σποροθλάστης, κ.τ.λ. But the name, though thus perfectly explicable, is very badly chosen to designate a strictly insectivorous species, its only pertinence being in geo-, signifying the humility of this bird of brake and briar.
    - 142. G. phīl-ā-dēl'-phǐ-ā. Named for the "city of brotherly love." Gr. φιλέω, I love, ἀδελφός, brother; the latter from ὰ connective (for ἄμα) and δελφύς, the womb, that is, having one mother. But the compound itself, Philadēl'phāa, is classic, as the name of a city, and there are the actual words φιλαδέλφεια, φιλαδελφία, amor fraternus, charitus fraterna. The Lat. is marked for quantity as above in the authority consulted; but some contend for the Greek accent, philadelphū'-a.
    - 143. G. măc-gil-līv-rāy'-ī. To William Macgillivray, Esq., of Edinburgh, author of much of Audubon's scientific work, besides several other important treatises.
    - 144. Ic-těr'-ĭ-ă vĭr'-ēns. A dialectic form, invented by Vieillot, of Gr. Ἰκτεροs or Lat. icterus; primarily, the disease jaundice; also a certain yellow bird, probably the golden oriole of Europe, by the sight of which jaundiced patients were fancied to be cured. The name was in 1760 by Brisson applied to the American orioles as a generic term, Icterus; and by Vieillot later, in the form Icteria, to the present genus.—Lat. virens, present participle of vireo, I grow green.
    - 145. I. v. lon-gi-caud'-ă [-cowda]. Lat. longus, long, and cauda, tail.
    - 146. Mỹĩ-ŏ-dĩ-ōc'-tēs mĩ-trā'-tǔs. Gr. μυῖα, a fly, and διώκτης, a pursuer. Lat. mitratus, wearing a turban; Gr. μίτρα, a turban or other head-dress. cf. μιτόω, I weave. The word is sometimes six-syllabled, but properly reducible to five, the ỹĩ, from Gr. vî, being slurred; the sound is that of mwee-, not mī- or mē-.

- 147. Myiodioctes pusillus (Wils.) Bp. B 213. C 102. R 125.

  Green Black-capped Flycatching Warbler.
  - 148. Myiodioctes pusillus pileolatus (Pall.) Ridg. B —, C 102a. R 125a.

    Pacific Black-capped Flycatching Warbler.
- 149. Myiodioctes canadensis (L.) Aud. B 214, 215. C 103. R 127. Canadian Flycatching Warbler.
- 150. Cardellina rubrifrons (Gir.) Scl. B —, C —, R 131.

  Red-fronted Flycatching Warbler.
- 151. Setophaga picta Sw. B 218. C 105. R 129.
  Painted Flycatching Warbler.
- 152. Setophaga ruticilla (L.) Sw. B 217. C 104. R 128.

  American Redstart.
- 153. Certhiola bahamensis Reich. B 301. C 106. R 159. (!W. I.)

  Bahaman Honey Creeper.
  - 147. M. pŭ-sīl'-lŭs. See Sitta, No. 60.
  - 148. M: p. pī-lĕ-ŏ-lā'-tŭs. Lat. pileum or pileolum, Gr. πίλος, a kind of cap, a skull-cap; pileolatus, capped. In late days, pileum has become a technical word in ornithology, meaning the top of the head.
- 149. M. căn-ă-dēn'-sīs. Latinized from Canada, with the termination -ensis. Canada is said to be the Iroquois word Kanata, a village or collection of huts.
- 150. Cār-dēl-lī'-nă rŭb-rī'-frōns. Apparently an arbitrary variation from Lat. carduelis, a kind of finch, from carduus, a thistle.—Lat. ruber, red, and frons, the forehead. The pronunciation of rubrifrons is in question; everybody says roo'bryfrŏnz; as it is not a classic word, we can only mark it by analogy with such words as rūbrico, &c. But see above, Dendræca, No. 126, in favor of rūb'rffrōns, as the i here comes before f and a liquid.

Not in the orig. ed. of the List; since discovered by H. W. Henshaw in New Mexico.

- 151. Sē-tŏ'-phă-gă pīc'-tă. Gr. σήs, genitive σητόs, an insect; and φαγεῖν, to eat. The connecting vowel o need not lengthen before ph, as this is only equivalent in force to f.—Lat. pictus, painted, pictured, here in the sense of brightly or highly colored; pingo, I paint, depict.
- 152. S. rtt-Y-cil/-la. Lat. rutilus, reddish; for the rest see Motacilla, No.86. The word is exactly equal to redstart, which is Anglicized from the Germ. Rothster or Rothster, all three words meaning simply redtail.
- 153. Cēr-thī'-ŏ-lă bă-hă-mēn'-sīs. Certhiola is a coined diminutive of Certhia, which see, No. 62; we usually hear it accented on a long penult, which is certainly vicious. — Bahamensis is Latinized from Bahama.

In the first ed. of the Check List, this species stands as *C. flaveola*, corrected in the Appendix. If we were to use the latter, it would be *flāvūlā*, not *flāveola*. Certhiola is correctly formed as a diminutive from Certhia, like lineola from linea; for the general rule, however, in cases when the stem ends in a consonant, we may recall the exquisite lines attributed to the death-bed of Hadrian:—

Animula vagula blandula, Hospes comesque corporis, Quæ nunc abibis in loca, Pallidula rigida nudula, Nec, ut soles, dabis jocos?

- 154. Pyranga rubra (L.) V. B 220. C 107. R 161.
  Scarlet Tanager.
- 155. Pyranga æstiva (L.) V. B 221. C 108. R 164. Summer Tanager.
- 156. Pyranga æstiva cooperi (Ridg.) Coues. B —. C 108a. R 164a. Cooper's Tanager.
- 157. Pyranga hepatica Sw. B 222. C 109. R 163. Hepatic Tanager.
- 158. Pyranga ludoviciana (Wils.) Bp. B 223. C 110. R 162. Louisiana Tanager.
  - 159. Hirundo erythrogastra horreorum (Bartr.) Coues. B 225. C 111. R 154. Barn Swallow.
  - 160. Iridoprocne bicolor (V.) Coues. B 227. C 112. R 155. White-bellied Swallow.
  - 161. Tachycineta thalassina (Sw.) Cab. B 228. C 113. R 156. Violet-green Swallow.
  - 154. Py-rān'-gă rŭb'-ră. The word Pyranga has a classic twang, as if formed in part from the Gr. πῦρ, fire; but it is a barbarous word, taken from some South American dialect. Several similar combinations of letters occur in Marcgrave. Vieillot wrote it Piranga in 1807, and Pyranga in 1816. The latter has come into general use.—The English tanager is simply altered from the South American tanagra or tangara, both of which words occur in the older authors, the latter being in general use until Linnæus, perhaps by a misprint, gave the former currency.
  - 155. P. aēs-tī'-vă. See Dendræca, No. 111.
  - 156. P. a. coop'-ĕr-ī. To Dr. J. G. Cooper, of California.
  - 157. P. hē-păt'-ĭ-că. Gr. ηπαρ, genitive ηπατος, the liver, or Lat. hepar, hepatis, the same; whence ηπατικός or hepaticus, the direct adjective. The allusion is to the liver-colored plumage.
  - 158. P. lū-dō-vĭ-cĭ-ā'-nă. See Thryothorus, No. 68.
  - 159. Hĭr-ūn'-dō ĕr-y-thrŏ-gās'-tră hōr-rĕ-ō'-rŭm. Lat. hirundo, a swallow, from the Gr. χελιδών, of same meaning.—Gr. ἐρυθρός, red or ruddy, and γαστήρ, the belly.—Lat. horreum, a barn, in the genitive plural. (On the etymology of hirundo, and various other, including the English, names of swallows, see Birds Col. Vall., i, 1878, p. 369.)
  - 160. Ir-ĭ-dō-prōc'-nē bī'-cŏl-ŏr. Gr. ¸¹Iριs, genitive ¸'Iριδos, Lat. Iris, Iridis, Iris, the messenger of the gods; also the rainbow; from εἴρω or ἐρω, to announce. The allusion is to the sheen of the plumage. Gr. Πρόκνη, or Lat. Procne or Progne, a proper name, the daughter of Pandion, fabled to have been transformed into a swallow.—Lat. bicolor, two-colored.
  - 161. Tāch-y-cīn-ē'-tā thāl-ās'-sī-nā. Gr. ταχυκινητοs, moving rapidly, i.e., a swift runner; ταχύs, swift (θέω, to run); κινητήρ, from κινέω, to move.—Gr. θαλάσσινοs, sea-green, θάλασσα, the sea, from äλs, the sea, or salt. Observe accentuation of thalassina. We keep the penult of Tachycinē'ta long as being Gr. η, but are not sure that it should not be transliterated Tachycin'ēta.

- 162. Petrochelidon lunifrons (Say) Cab. B 226. C 114. R 153. Cliff or Eave Swallow.
- 163. Cotile riparia (L.) Boie. B 229. C 115. R 157.
  Bank Swallow.
  - 164. Stelgidopteryx serripennis (Aud.) Bd. B 230. C 116. R 158. Rough-winged Swallow.
- 165. Progne subis (L.) Bd. B 231. C 117. R 152. Purple Martin.
- 166. Ampelis garrulus L. B 232. C 118. R 150. Bohemian Waxwing.
- 167. Ampelis cedrorum (V.) Bd. B 233. C 119. R 151. Cedar Waxwing.
- 168. Phaïnopepla nitens (Sw.) Scl. B 234. C 120. R 26. Black Ptilogonys.
- 162. Pēt-rō-chěl-ī'-dōn lū'-nĭ-frōns. Gr. πέτρα, a rock, and χελιδών, a swallow; alluding to the places where the nests are often built.—Lat. luna, the moon, that is, a crescent, and frons, the forehead or front; referring to the white frontal crescent. Luna is contracted from Lucina, a proper name, epithet of Juno, from luceo, I shine; lux, light.
- 163. Go'ti-le rī-păr'-ĭ-ă. The generic name was originally written Cotile by Boie, afterward by him Cotyle. The latter orthography came into general use, the alleged etymology being κοτύλη, a cup, in supposed allusion to the excavations in which the bird nests. The proper orthography is Cotile, from κωτιλάs, the swallow; literally, the twitterer, babbler, prattler, from κωτίλλω, I prate. (See Wharton, Ibis, October, 1879, p. 451; and Coues, Bull. Nuttall Club, April, 1880, p. 96.) Lat. riparia, riparian: ripa, the bank of a stream.
  - 164. Stēl-gǐ-dōp'-tě-ryx sēr-rǐ-pēn'-nĭs. Gr. στελγίς or στλεγγίς, a scraper; and πτέρυξ, wing.

     Lat. serripennis, saw-feathered; serra, a saw, penna, a feather. Both words mean substantially the same thing, having reference to the peculiar structure of the outer web of the first primary.
  - 165. Prōg'-nē sŭb'-ĭs. Lat. Proque; see Iridoprocne, No. 160. Lat. subis, a word not known except as applied by Pliny to a bird said to break eagles' eggs; application in this case unknown.
  - 166. Ām'-pĕ-līs gār'-rǔ-lūs. Gr. ἀμπελίς, or ἄμπελος, the grapevine; also, a small bird which frequented vineyards, by some conjectured to be the present species; ἀμπελίων also occurs as the name of a bird.—Lat. garrulus, garrulous, loquacious, from garrio, I chatter (Gr. γηρύω or γαρύω, I speak, γῆρυς or γάρυς, voice); also, as substantive, a jay-bird, which is the implication in this case.
  - 167. A. cēd-rō'-rŭm. Lat. cedrus, genitive plural cedrorum, the cedar; Gr. κέδρος.
  - 168. Phă-ī-nŏ-pēp'-lă nīt'-ēns. Dr. Sclater says (Ibis, 1879, p. 223) that he formed the word from φαεινός, shining, and that it should be written as above, as he originally did. This, however, is merely a poetic form, from φαείνω, itself poetic for φαίνω. It would appear to be most naturally written Phænopepla, like phænomenon, phænogamous, &c., from the same source; but if the orthography Phaïnopepla, in five syllables, be preserved, it can be easily defended. Gr. πέπλα, poetic plural of πέπλος, a robe. Lat. nitens, present participle from niteo, I shine.

- 169. Myiadestes townsendi (Aud.) Cab. B 235. C 121. R 25. Townsend's Flycatching Thrush.
- 170. Vireo olivaceus (L.) V. B 240. C 122. R 135. Red-eyed Greenlet.
- 171. Vireo flaviviridis Cass. B 241. C —. R 136. Yellow-green Greenlet.
- 172. Vireo altiloquus barbatulus (Cab.) Coues. B 243. C 123. R 137. Black-whiskered Greenlet.
- 173. Vireo philadelphicus Cass. B 244. C 124. R 138. Brotherly-love Greenlet.
- 174. Vireo gilvus (V.) Bp. B 245. C 125. R 139. Warbling Greenlet.
- 175. Vireo gilvus swainsoni Bd. B —. C 125a. R 139a. Western Warbling Greenlet.
- 176. Vireo flavifrons V. B 252. C 126. R 140. Yellow-throated Greenlet.
- 177. Vireo solitarius V. B 250. C 127. R 141.
  Blue-headed Greenlet.
- 178. Vireo solitarius cassini (Xantus) Ridg. B 251. C —. R 141a. (?) Cassin's Greenlet.
- 169. MỹI-ă-dēs'-tēs [mweeadaystace] town'-send-ī. Gr. μυῖα, a fly, and ἐδεστήs, an eater; ἔδω, or ἔδομαι, I eat; see Myiodioctes, No. 146. (Not to be written Myiadectes, as if fly-"taker," Muscicapa, from μυῖα and δεκτήs, from δέχομαι). — To J. K. Townsend, from whom Audubon received many new birds, and to whom he dedicated several.
- 170. Vīr'-ĕ-ō [vĭr'ryoh, not vī'reo] ŏl-ī-vā'-cĕ-ŭs. Lat. vireo, a kind of bird, from vireo, I am green or flourishing. Late Lat. olivaceus, olive-like, olive-colored; green obscured with neutral tint; oliva, the olive, from olea, Gr. ἐλαία, the olive-tree; whence oleum, Gr. ἔλαιον, Eng. oil, oleaginous, &c.
- 171. V. flā-vī-vī-vī-t-dīs. Lat. flavus, yellow, and viridis, green, from vireo. See Auriparus, No. 56. Commonly but wrongly written flavoviridis.

  This species is not in the first ed. of the Check List; it has only recently been discovered in the United States, in Texas, by J. C. Merrill.
- 172. V. āl-tǐ-lò-quus bār-bā/-tu-lus. Lat. altus, high, from alo, I bear up, sustain, and loquus, an adjective from loquor, I speak; pronounced ahlty/lockwooce, like ventri/loquist, grandi/loquent, &c.—Lat. barbatulus, having a small beard; barbatus, bearded; barba, a beard. The allusion is to the dusky maxillary streaks.
- 173. V. phil-ă-dēl'-phi-cus. See Geothlypis philadelphia, No. 142.
- 174. V. gīl'-vŭs [g hard]. Lat. gilvus, gilbus, galbus, helvus, yellowish, greenish-yellow; German gtlb, Ital. giallo, A. S. gelew, geoluwe; related to fulvus, flavus, &c.
- 175. V. g. swain/-sŏn-i. To William Swainson.
- 176. V. flā'-vǐ-frōns. Lat. flavus, yellow; frons, forehead. See Auriparus, No. 56.

  Obs.—It would appear from B. C.V., i, 1878, p. 494, that the proper name of this species is V. ōch-rŏ-leū'-cŭs (Gm.) Coues. Gr. ἀχρός, ochraceous, yellowish, and λευκός, white.
- 177. V. sō-lǐ-tā/-rǐ-ŭs. Lat. solitarius, solitary; solus, alone.
- 178. V. căs'-sin-i. To John Cassin, of Philadelphia, sometime the "Nestor of American ornithology"; the only ornithologist America ever produced who knew any considerable number of Old World birds. Not in the orig. ed.; since recognized.

- 179. Vireo solitarius plumbeus (Coues) Allen. B —. C 127a. R 141b. Plumbeous Greenlet.
- 180. Vireo vicinior Coues. B —. C 128. R 147.
  Gray Greenlet.
- .181. Vireo noveboracensis (Gm.) Bp. B 248. C 129. R 143. White-eyed Greenlet.
- 182. Vireo huttoni Cass. B 249. C 130. R 144.

  Hutton's Greenlet.
- 183. Vireo belli Aud. B 246. C 131. R 145.
  Bell's Greenlet.
- 184. Vireo pusillus Coues. B —. C 132. R 146. Least Greenlet.
- 185. Vireo atricapillus Woodh. B 247. C 133. R 142.
  Black-capped Greenlet.
- 186. Lanius borealis V. B 236. C 134. R 148. Great Northern Shrike; Butcherbird.
- 187. Lanius ludovicianus L. B 237. C 135. R 149. Loggerhead Shrike.
- 188. Lanius ludovicianus excubitorides (Sw.) Coues. B 238. C 135a. R 149a. White-rumped Shrike.
- 179. V. s. plūm'-bĕ-ŭs. Lat. plumbeus, plumbeous, lead-colored; plumbum, lead.
- 180. V. vī-cī/-nǐ-ōr. Lat. comparative degree of vicinus, neighboring; vicinia, a neighborhood or vicinity; this from vicus, digammated from Gr. οἶκοs, a house. The allusion is to the close resemblance of the species to others.
- 181. V. nov-e-bor-a-cen'-sis. Very late Latin for of, or pertaining to, New York; novus, new, and eboracensis, pertaining to Eboracum, the old name of York, England; Noveboracum is literally New York.
- 182. V. hŭt'-tŏn-ī. To William Hutton, of Monterey, California.
- 183. V. běl'-lī. To J. G. Bell, of New York, for many years the most skilful and most distinguished taxidermist of America.
- 184. V. pŭ-sīl'-lŭs. See Sitta pusilla, No. 60.
- 185. V. ā-trī-căp-īl'-lŭs. Lat. ater, atra, black; the opposite of albus. It properly means dead black, as niger does glossy black, which latter would have been better in this case. Capillus, hair of the head, from caput, head; whence English capillary, thready.
- 186. Lăn'-ĭ-ŭs bŏr-ĕ-ā'-lĭs. Lat. lanius, a butcher; from lanio, I rend, lacerate. See Falco, No.502.—Lat. boreas, the north wind, h. e., the north; whence borealis, northern. For reason of the generic change from Collurio of the orig. ed. of the Check List, and for Shrikes' names in general, see Birds Colorado Valley, i, 1878, p. 537 et seq.
- 187. L. lū-dō-vĭ-cĭ-ā'-nŭs. Lat. Ludovicus, Louis, a proper name. The application here is to the Territory of Louisiana, formerly of great extent. See Thryothorus, No. 68.
- 188. L. ēx-cŭb-ĭ-tō-rī/-dēs. Lat. excubitor, a watchman, sentinel, from ex, out of, and cubitor, one who lies down, from cubo; i. e., an out-lier. The termination of the word is the Gr. elδos, appearance or resemblance (εἴδω, I see). There is a difference in the orthography of the word: it has oftenest been written excubitoroides, and pronounced in six syllables, with the accent on the penult. But if this spelling is used, it should be excubitoroides,

- 189. Hesperophona vespertina (Coop.) Bp. B 303. C 136. R 165. Evening Grosbeak.
- 190. Pinicola enucleator (L.) V. B 304. C 137. R 166. Pine Grosbeak.
- 191. Pyrrhula cassini (Bd.) Tristr. B—. C 138. R 167. (!A.)
  Cassin's Bullfinch.
- 192. Passer domesticus (L.) Koch. B —. C 187. R —. [Imp. and Nat.]
  Philip Sparrow.

with the diæresis over the i, and consequently making seven syllables. So long a word is therefore preferably shortened by omitting the connecting vowel o; which, with the usual change of Gr.  $\epsilon \hat{t}$  to long i in Latin, gives the above spelling and pronunciation. The full number of letters in the compound is *excubitorieides*.

- 189. Hēs-pěr-ŏ-phō'-nă vēs-pēr-tī'-nă. Gr. ἐσπέρα, Hesperus, the west, the place of sunset (χώρα, region, being understood); hence, the evening; and φωνή, the voice; φωνέω, I speak; φάω, φημί, related to φαίνω, &c.—Lat. Vespertinus, pertaining to the evening, Vesperus being the same as Hesperus.—The genus-name is universally written Hesperiphona, as Bonaparte originally spelled it, but the above is certainly correct, as it is pure Greek for what Vesperisona would be the Latin of. The pleonastic name signalizes a belief, formerly entertained, that the bird sings chiefly at evening.— Grosbeak or grossbeak is corrupted from the Fr. grosbec, thick-bill.
- 190. Pī-nǐ'-cŏ-lǎ ē-nū-clē-ā'-tōr. Lat. pinus, a pine, and incola, an inhabitant, from colo, I cultivate; formed like many other words in -cola, as saxicola, agricola, &c. Lat. enucleator, one who "shells out," or enucleates; from enucleo, I take out the kernel; nucleus, the nucleus or kernel, this from nux, a nut. The two words indicate the characteristic habitat and habit of the bird.
- 191. Pỹr'-rhǔ-lã cǎs'-sǐn-ī. Lat. pyrrhula, a bullfinch; a diminutive of Pyrrhus, a proper name; Gr.  $\pi v \hat{\rho} \hat{\rho} \delta s$ , fiery-red, from  $\pi \hat{v} \rho$ , fire; in allusion to the bright color of the bird. To John Cassin.

It is still uncertain what relation this bird may best be considered to bear to the Old World form  $P.\ coccinea$ , as no Alaskan specimens, since the type, have been forthcoming. We give it as it stands in the body of the orig. ed. of the Check List.

Note. — Another species of this genus has lately been reported from Greenland by Kumlein (Bull. U. S. Nat. Mus., No. 15, p. 74; 1879); but the case remains very dubious.

192. Pās'-ser dom-ēs'-ti-cus. Many interesting words are grouped about this ubiquitous bird, which has been named in nearly or quite all civilized languages; some of them may be here noticed. — 1. Passer domesticus, literally "house sparrow," is itself a very old Latin biblionym, though used for less than a century as a technical term. Passer is good Latin for sparrow, and particularly for this very species, which is said to have been noted, if not named, for its salacity; but the etymology of this word is unknown to us, as it also appears to be to the authors of several lexicons; one says passer for padser, from pando, I spread. Passer seems to have become of general signification, almost as broad as English "bird" or "fowl." The Ital. is passera, passere, passara, and this language also had passer domesticus in passara cazarenga. The word passes directly into the Fr. passerat, passereau, and to the Eng. technical adjective passerine, sparrow-like; while the Span. paxaro (as if pacsaro) or pajaro is apparently the same. - 2. The Gr. name for this species was  $\sigma \tau \rho o \nu \theta \delta s$ , in Aristotle; which in modern technic has become, in the form struthio, the name of the ostrich, Struthio camelus L., and has given our Eng. adjective struthious, ostrich-like. The actual application to the ostrich, however, dates back to Aristotle, whose στρουθός δ εν Λιβύη, or Libyan fowl, was the ostrich — like the Lat. passer marinus, i. q., the bird brought from over the sea. -3. The Gr. word  $\pi \nu \rho \gamma i \tau \eta s$ , from  $\pi \dot{\nu} \rho \gamma \sigma s$ , a tower, and meaning a dweller in the tower, has been of late years used to some

- 193. Passer montanus (L.). B —. C —. R —. [Imp. and Nat.] European Tree Sparrow.
- -194. Carpodacus purpureus (Gm.) Gr. B 305. C 139. R 168. Purple Finch.
- 195. Carpodacus cassini Bd. B 307. C 140. R 169. Cassin's Purple Finch.
- 196. Carpodacus frontalis (Say) Gr. B 308. C 141. R 170. Crimson-fronted Finch; House Finch; Burion.

extent as the generic name, under the form Pyrgita: though having originally no reference to the species whatever, it is a very apt designation of a bird which nests so habitually about buildings. — 4. The word Fringilla, one of a large group, giving name to the Finch family, Fringillidæ, and to the English adjective fringilline, is the origin of the word finch itself; though it is only for about a century that it has had any thing to do with the present species. Fringilla is the Latin name of the same bird that the Greeks called  $\sigma\pi'(\zeta\alpha)$  or  $\sigma\pi'(\zeta\eta)$ , spiza, the F. calebs L., English Chaffinch. Fringilla or fringuilla has been derived by some from frango, I break, as the bird does seeds (just as we have in Gr.  $\theta \lambda \nu \pi is$  or  $\theta \rho \alpha \nu \pi is$ ). But its etymology appears when we regard the non-nasalized form frigilla, from frigutio or frigultio (= fringulio or fringultio, formed like singultio, I hiccup), I twitter, chirp, stammer; these words being themselves lengthened from friquio, I croak, as a crow, and this from frigo, I squeak, squeal. (Cf. Gr. φρύγω, and the actual φρυγίλοs, the name of a bird in Aristophanes, and source of the modern genus Fregilus, a jackdaw. The idea seems to be some short sharp sound, as the hissing, sizzling of something cooking, — frigo or φρύγω, I cook.) Fringilla reappears in several Italian forms, from two of which two series of words branch off; from such as fringuello, frinco, are derived, with loss of the r, Germ. find, fint, and Eng. finch; while from such as frinsone we pass through grinson, quinson, pinson, or later Fr. pincon to Eng. spink, a name of F. calebs. — 5. An entirely different set of words gives the pedigree of modern Eng. sparrow, back from which we pass to sparrowe, or sparowe, or sparwe, Gothic sparwa or sparva, A. S. spearwa; related forms being spörr, spar, sper, spurr, spurr, sparf, spatz, sperg, sperk, sperlingk, round again to the present Germ. sperling or haussperling, housesparrow, passer domesticus. - 6. Eng. sparrow also curiously leads us back again to Latin, through such a form as sparva, Latinized as sparrius; so, also, Falco sparverius, i. q. fringillarius, σπιζίας, Fr. espervier or épervier, anglicè sparrow-hawk. — 7. There is said to be an old Flemish name mousche for this bird, which may not improbably connect with O. Fr. moucet, moisson. — 8. The present Fr. is moineau, or moineau franc, or moineau de ville. — 9. Several languages have applied cant names to this sturdy vulgarian; Span. gorrion, thief, rogue, scamp; Fr. gamin; American tramp, hoodlum. - 10. An onomatopæia as interesting as Fringilla itself has arisen from the sharp, abrupt, dissyllabic note. This is represented by the syllables yellop (cf. Gr. ἐλλόπ-os), yellup, or phyllup, easily becoming Philip. Early in the sixteenth century appear the "Boke of Phyllup Sparrowe" and the "Praise of Philip Sparrow"; and this name is Shakspearian.

Introduced, but now thoroughly naturalized everywhere.

- 193. P. mon-ta'-nus. Lat. montanus, of mountains.
  - Not in the orig. ed.; since introduced from Europe, and naturalized in some places.
- 194. Cār-pŏ'-dă-cŭs pūr-pŭr'-ĕ-ŭs. Gr. καρπός, a fruit, and δάκος, from δάκνω, I bite; 2d aorist ἔδακον, or δάκον. — Lat. purpureus, purple; Gr. πορφύρεος, English porphyry, &c.; cf. πυρφόρος (πῦρ, φέρω) the fire-bearer, an epithet of Prometheus. — The quantity of the penult is in question; we usually hear carpodā'-cus in this country; but carpŏ'-ducus is preferable.
- 195. C. căś'-sĭn-ī. To John Cassin.
- 196. C. fron-ta/-lis. Lat. frontalis, relating to the forehead; frons, forehead, front.

- 197. Carpodacus frontalis rhodocolpus (Cab.) Ridg. B C 141a. R 170a. Rose-breasted Finch.
- 198. Loxia leucoptera Gm. B 319. C 142. R 173. White-winged Crossbill.
- 199. Loxia curvirostra americana (Wils.) Coues. B 318. C 143. R 172. Common American Crossbill.
- 200. Loxia curvirostra mexicana (Strickl.) Coues. B —. C 143a. R 172a. Mexican Crossbill.
- 201. Leucosticte atrata Ridg. B —. C —. R 176. Ridgway's Rosy Finch.
- 202. Leucosticte australis Allen. B —. C —. R 177. Allen's Rosy Finch.
- 203. Leucosticte tephrocotis Sw. B 322. C 144. R 175. Swainson's Rosy Finch.
- 204. Leucosticte tephrocotis litoralis (Bd.) Coues. B —. C R 175a. Baird's Rosy Finch.
  - 197. C. f. rhŏ-dŏ-cōl'-pŭs. Gr.  $\dot{\rho}\dot{\phi}\delta\sigma\nu$ , the rose, and  $\kappa\dot{\phi}\lambda\pi\sigma s$ , the breast; in allusion to the rose-red color of that part.

The form C. f. hamorrhous, given in the orig. ed. of the Check List, is the Mexican race; the above should replace No. 141a.

- 198. Lōx'-ĭ-ă leū-cōp'-tě-ră. Gr. λοξίαs, an epithet of Apollo, whose oracles were sometimes obscure or equivocal; from λοξόs, oblique, devious, deviating from a straight line; very pertinent to the Crossbill. Gr. λευκόs, white, and πτερόν, wing.
- 199. L. cūr-vī-rōs/-tră. Lat. curvus, curved; and rostrum, bill. In this and numberless similar cases of a noun compounded with an antecedent adjective, the whole word is treated as an adjective, capable of inflection according to gender. Thus curvirostra is as if curviroster or curvirostr-us, -a, -um. So we even find longicaud-us, -a, -um, like auricom-us, -a, -um, and the Vergilian centiman-us, -a, -um. In such a case as the present, the adjectival form curvirostris (like -ventris) might be more elegant. But curvirostra has the sanction of several centuries' use as a noun, having apparently been invented as a Latin synonym of Loxia; it is not, however, classic. Other synonyms are crucirostra, crucifera, cruciata; Fr. Bec-croisé, Germ. Arcusfanabel, &c.
- 200. L. c. mēx-ĭ-cā'-nă. Lat. mexicana, of Mexico. See Sialia, No. 28.
  - 201. Leū-cō-stīc'-tē ā-trā'-tă. Gr. λευκός, white, and στικτή, variegated; from στίζω, I puncture, brand, or mark. Lat. atrata, blackened; a participial adjective, from an obsolete or rather hypothetical verb atro.

Not in the orig. ed. of the Check List; described from Colorado by Ridgway, Amer. Sportsm., iv, No. 16, p. 241, July 18, 1874.

- 202. L. aūs-trā'-līs. Lat. australis, southern; from auster, the south wind, hot and dry; this from Gr. αἴω, I dry up or parch.
  Not in the orig. ed. of the Check List, as then not supposed valid.
- 203. L.  $t\bar{e}ph-r\check{o}-c\bar{o}'-t\check{s}$ . Gr.  $\tau\epsilon\phi\rho\delta s$ , gray, ashy, from  $\tau\epsilon\phi\rho\alpha$ , ashes; and  $-\delta\delta s$ , genitive  $-\delta\epsilon\tau\delta s$ , the ear; the connective consonant c being introduced for euphony.
- 204. L. t. lī-tŏr-ā'-līs. Lat. litoralis, littoral; from litus, the shore, of sea, lake, or river. The word is commonly written littoralis, but preferably as above.

  Not in the first ed. of the Check List, as not then supposed to be valid.

- 205. Leucosticte griseinucha (Brandt) Bd. B 323. C 144a. R 174. Brandt's Rosy Finch.
- 206. Leucosticte arctoa (Pall.) Bp. B 324. C 145. R —.
  Pallas's, Rosy Finch.
- 207. Ægiothus linaria (L.) Cab. B 320. C 146, 146a. R 179. Common Red-poll.
- 208. Ægiothus linaria holboelli (Brehm) Coues. B —, C —, R 179a. (?) Holböll's Red-poll.
- 209. Ægiothus hornemanni (Holb.) Coues. B 321. C —. R 178. (G.) Greenland Mealy Red-poll.
- 210. Ægiothus exilipes Coues. B —. C 146b. R 178a.

  American Mealy Red-poll.
- 211. Linota flavirostris brewsteri (Ridg.) Coues. B —. C 147. R 180. (?)
  Brewster's Linnet.
- 205. L. grīs-ĕī-nū'-chă. Lat. griseus, gray, and nucha, the nape or scruff of the neck. Neither part of the word is classic; griseus is Latinized from such a word as seen in Fr. gris, Ital. griso, English grisly; and nucha, a technical word in ornithology, is Latinized from Fr. nuque, the nape (A. S. cnæp, a knob, knoll), which is the same as Gaelic cnoc, Welsh cnwc. Nape is thus closely related to neck itself; A. S. hnecca, Dan. nakke, Dutch nak or nek, Germ. naten, &c.
- 206. L. ārc-tō'-ă. Gr. ἄρκτος, a bear; also, the constellation; hence, the north; adjective ἀρκτφος, same as ἀρκτικός, northern, whence Lat. arctous and arcticus, of same signification.
- 207. Aeg-1'-5-thŭs lī-nā'-rĭ-ă. Gr. Aiγίοθος, given by Cabanis as a proper name: supposably derived from aiγίς, a goat-skin, or ægis, and τίθημι, to put or place, as if the shield-bearer, like Ægisthus. The application is far from being evident. The word is probably only another form of αἴγιθος, the name of an unknown bird, occurring in Aristotle, Hist. ix. 1, conjectured by some to be this very species.—Lat. linaria; from linum (Gr. λίνον), flax; the root is seen in many words, as line, linear, linen, lint, linnet, &c.
- 208. A. 1. hŏl'-boĕl-lī. To Carl v. Holböll, a Danish naturalist, chiefly known in ornithology of for his researches in Greenland.

  Not recognized in the first ed. of the Check List.
- 209. A. hōrn'-ĕ-măn-nī. To —— Hornemann, who had to do with Greenland birds.

  This species is not in the orig. ed. of the Check List. It is only American inasmuch as it is found in Greenland. It is absolutely confined to that country, and is the bird usually quoted as Greenlandic "canescens."
- 210. A. ex-il'-I-pes. Lat. exilis (for exigilis, from exigo), small, slender, &c., and pes, foot. See Ardetta, No. 667.
- 211. Lī-nō'-tă flā-vĭ-rōs'-trĭs brews'-tĕr-ī. See Linaria, above: the word is not classic, being directly Latinized from the Fr. linotte, one of the numberless words from linum, linea, &c. Lat. flavirostris, yellow-billed. To William Brewster, of Cambridge, Mass., an excellent ornithologist.

This is questionably North American, and questionably a good species.

- 212. Chrysomitris pinus (Bartr.) Bp. B 317. C 148. R 185.
  Pine Linnet; American Siskin.
- 213. Astragalinus tristis (L.) Cab. B 313. C 149. R 181.

  American Goldfinch.
- 214. Astragalinus lawrencii (Cass.) Coues. B 316. C 150. R 183. Lawrence's Goldfinch.
- 215. Astragalinus psaltria (Say) Coues. B 314. C 151. R 182.

  Arkansaw Goldfinch.
- 216. Astragalinus psaltria arizonæ Coues. B —. C 151a. R 182a. Arizona Goldfinch.
- 212. Chrỹ-sŏ-mī'-trīs pī'-nŭs. Gr. χρυσομίτρις, having a golden head-dress or girdle; χρυσέος, golden, and  $\mu l \tau \rho \alpha$ , a mitre. There are other forms of the word, varying in the vowels, as χρυσομίτρης and χρυσομητρις. The latter, which occurs in Aristotle, is translated aurivittis by Gaza; as Sundwall remarks, heightening the probability that it is the same word as χρυσομίτρηs, and is based upon the bright appearance of the European Goldfineh, F. carduelis L. — Some other names of classic origin for the Goldfineh and its relatives may be here conveniently noted. Aristotle had three species of "Acanthophaga" as he called them; i. e., birds living upon prickly plants; as we should say, "thistle-birds." 1. One of these was the θραυπίς or θλυπίς, concerning which see Geothlypis, No. 141. 2. The χρυσομήτριs, as just said. 3. His ἀκανθίς, which was undoubtedly the Fringilla cannabina L. This in Latin becomes spinus, of late years taken as the specific name of F. spinus L. — The exact Latin of "thistle-bird" is carduelis, occurring in Pliny; it is from carduus, a thistle, and reappears in numerous shapes; as Ital. carduello, cardello; carduelino, cardellino (compare Cardellina, No. 150), and also gardello and gardellino; Fr. chardonneret, &c. Aristotle speaks of the sharp voice of his ἀκανθίς — λιγυρά; whence ligarinus, another of the many names for birds of this kind. So have we later derived siskin from the sharp note; Swedish siska, Dutch sijsken, Germ. şiefig, Polish czyz, &c. — Another Greek name for some kind of thistle-bird, perhaps the European Goldfinch, is ἀστραγαλίνος, in 1850 applied by Cabanis to the American Goldfinch, as a generic term: see next word. - Lat. pinus, a pine-tree.
- 213. As-tră-gă-li'-nŭs trīs'-tĭs. Gr. ἀστραγαλῶνοs is given by Cabanis as the word, and as a name of a thistle-bird; it is evidently an adjectival form from ἀστράγαλοs, a die, one of the ankle-bones, and also, in Dioscorides, the name of some kind of plant; whence the modern botanical genus Astragalus. The original application of ἀστραγαλῶνοs is undoubtedly to some bird that lived upon, or frequented, the plant in mention, its recent transference to an American Goldfinch being of course arbitrary. When the present species was first described it was called chardonneret de l'Amérique, i. e., carduelis americana: see No. 212. Lat. tristis, sad, in allusion to the plaintive cry of the bird.
- 214. A. law-ren'-ci-i. To George N. Lawrence, of New York, the eminent ornithologist.
- 215. A. psāl'-trǐ-ă. See explanation of Psaltriparus, No. 53. Psaltria is not a Lat. adj. to be made agreeable in gender with Astragalinus, but a Greek noun, ψάλτρια, signifying a female lutist. "Arkansaw" is not, as it would seem to be, "Kansas" with a prefix, nor is it the name by which the aborigines of that country knew themselves; nor is "Kansas" the right name of any tribe of Indians. The meaning of neither of these words is known. "Arkansaw" is preferable to Arkansas, as nearer the original "Arkanso."
- 216. A. p. ă-rī-zō'-naē. Named after the Territory of Arizona, where discovered in 1864. See Peucæa, No. 253.

- 217. Astragalinus psaltria mexicanus (Sw.) Coues. B 315. C 151b, R 182b. Mexican Goldfinch.
- 218. Astragalinus notatus (Du Bus) Coues. B 310. C —. R 184. (!M.) Black-headed Goldfinch.
- 219. Plectrophanes nivalis (L.) Meyer. B 325. C 152. R 186. Snow Bunting; Snowflake.
- 220. Centrophanes lapponicus (L.) Kaup. B 326. C 153. R 187. Lapland Longspur.
- 221. Centrophanes pictus (Sw.) Cab. B 327. C 154. R 188. Painted Longspur.
- 222. Centrophanes ornatus (Towns.) Cab. B 328, 329. C 155. R 189. Chestnut-collared Longspur.
- 223. Rhynchophanes maccowni (Lawr.) Bd. B 330. C 156. R 190. Maccown's Longspur.
- 224. Passerculus bairdi (Aud.) Coues. B 331. C 157, 157bis. R 191. Baird's Savanna Sparrow.
- 217. A. p. mēx-ĭ-cā'-nă. Lat. mexicanus, of Mexico. See Sialia, No. 28.
- 218. A. not-ā'-tus. Lat. notatus, noted, marked; noto, I make note of. In allusion to the distinction between this species and C. magellanicus.

  Not in the orig. ed. of the Check List. Said by Audubon to have been actually taken in Kentucky. (?)
- 219. Plēc-tro'-phā-nēs nīv-ā'-līs. The Gr. πλήκτρον, or Lat. plectrum, was an instrument for striking the lyre, from πλήσσω, I strike; also used for a quill, a spur, &c.; the meaning in this case is the hind claw of the bird, which is remarkably long and straight. The rest of the word is from value, to appear, to seem, &c., the claw in mention being likened to the instrument spoken of. Obs. There is continual difference in opinion respecting the pronunciation of this and similar words, according to whether we consider them as Greek or as Latin. The rule in Greek would retain the accent upon the root of each word entering into the composition, giving Plec'-tro-phal'-nes. But in Latinizing it is allowable, and indeed preferable, to accent as above; as we have also done in the cases of Helmintho'phaga, Lopho'phanes, &c. The gender of the many coined words ending in -phanes is practically in question among ornithologists; we make them masculine.
- 220. Gēn-tro'-phā-nēs lāp-pon'-ĭ-cus. Gr. κέντρον, a prick, nail, claw, &c., from κεντέω, I. καλλας prick or goad. The reference, as in the case of Plectrophanes, is to the long hind claw. See Plectrophanes.—Lat. lapponicus, pertaining to Lapland, formerly Lapponia.
  - 221. C. pic'-tŭs. Lat. pictus, painted, from pingo, I paint or ornament; in allusion to the variegated colors.
  - 222. C. or-na/-tus. Lat. ornatus, adorned, decorated, from orno, I ornament.
  - 223. Rhyn-cho'-pha-nes mac-cown'-i. Gr. ρύγχος, snout, muzzle, beak, and φαίνω; in allusion to the large bill. See *Plectrophanes*.—To Capt. J. P. McCown, then of the U. S. Army.
  - 224. Pās-sēr'-cŭ-lŭs baird'-ī. Lat. passerculus, a little sparrow; diminutive of passer. To Spencer F. Baird, long time the leader in North American ornithology.

    Centronyx ochrocephalus, No. 157 bis of the first ed., is this species in fall plumage.

- /225. Passerculus princeps Mayn. B —. C 158. R 192. Ipswich Savanna Sparrow.
  - 226. Passerculus sandvicensis (Gm.) Bd. B 333. C 159b. R 193. Sandwich Savanna Sparrow.
- 227. Passerculus sandvicensis savana (Wils.) Ridg. B 332. C 159. R 193α. Common Savanna Sparrow.
- 228. Passerculus sandvicensis anthinus (Bp.) Coues. B 334. C 159a. R 194. Pipit Savanna Sparrow.
- 229. Passerculus sandvicensis alaudinus (Bp.) Ridg. B 335. C —. R 193b. Lark Savanna Sparrow.
- 230. Passerculus rostratus (Cass.) Bd. B 336. C 160. R 196.
  Beaked Savanna Sparrow.
- 231. Passerculus guttatus Lawr. B —, C 160a. R 195. St. Lucas Savanna Sparrow.
- 232. Poœcetes gramineus (Gm.) Bd. B 337. C 161. R 197. Bay-winged Bunting; Grass Finch.
- 233. Poœcetes gramineus confinis Bd. B —. C 161a. R 197a. Western Grass Finch.
- 234. Coturniculus passerinus (Wils.) Bp. B 338. C 162. R 198. Yellow-winged Sparrow.
  - 225. P. prīn'-cēps. Lat. princeps, first, principal; from primus, first, and -ceps.
- 226. P. sānd-vǐ-cēn'-sĭs. Named after Sandwich Island, one of the Kurile or Aleutian Archipelago.
- 227. P. s. să-vā'-nă. Properly Span. sabana or savana, anglicized savanna or savannah, a meadow. As a quasi-Latin word, it should have but one n, as in the Spanish. The quantity of the penult is marked by the general rule for accentuation in Spanish, that words ending in a vowel have the accent on the penult.
- 228. P. s. ān-thī/-nŭs. Arbitrarily formed from anthus, a pipit, which see, No. 89.
- 229. P. s. ăl-aūd-ī'-nŭs. Arbitrarily formed from Lat. alauda, a lark; this from the Celtic al, high, and aud, song.

  Not in the orig. ed., as then not recognized as valid.
- 230. P. rōs-trā/-tŭs. Lat. rostratus, beaked, i. e., having a large beak; rostrum, a beak; this from rodo, to gnaw, corrode, &c.
- 231. P. gūt-tā'-tŭs. Lat. guttatus, spotted, speckled; from gutta, a drop; as if marked with droppings.
- 232. Pŏ-oē'-cĕ-tēs grā-min'-ĕ-ŭs. Gr. πόα, ποία, πόη, ποίη, grass, herbage; and οἰκότης, an inhabitant; from οἴκος, a dwelling. The orthography of this word has been unsettled: it was first written Poocætes by Baird in 1858, and has since been variously spelled. The stem of the first word is πο, giving po-; and οἰκότης becomes in Latin æcetes; the above form seems eligible, as first emended by Sclater in 1859. It may be susceptible, but not preferably, of further contraction into Pæcetes. Lat. gramineus, grassy, figuratively applied to a bird that lives much in the grass; gramen, grass.
- 233. P. g. con-fi'-nis. Lat. confinis, like affinis, allied to, &c.; con, with, and finis, the boundary, limit, edge, or end of a thing.
- 234. Cō-tūr-nĭ'-cŭ-lŭs pās-sĕr-ī'-nŭs. Arbitrary diminutive of coturnix, a quail; said to be so called from the resemblance of the sound of its voice to the sound of the word.—

  Passerinus, an arbitrary adjective from passer, a sparrow; sparrowlike.

- 235. Coturniculus passerinus perpallidus Ridg. B —. C 162a, R 198a.
  Bleached Yellow-winged Sparrow.
  - 236. Coturniculus henslowi (Aud.) Bp. B 339. C 163. R 199. Henslow's Sparrow.
  - 237. Coturniculus lecontii (Aud.) Bp. B 340. C 164. R 200. Le Conte's Sparrow.
- -238. Ammodramus maritimus (Wils.) Sw. B 342. C 165. R 202. Seaside Finch.
- 239. Ammodramus maritimus nigrescens Ridg. B —. C 165a. R 203. Floridan Seaside Finch.
- 240. Ammodramus caudacutus (Wils.) Sw. B 341. C 166. R 201. Sharp-tailed Finch.
- 241. Ammodramus caudacutus nelsoni Allen. B —, C —, R 201a.

  Nelson's Sharp-tailed Finch.
- 242. Melospiza lincolni (Aud.) Bd. B 368. C 167. R 234. Lincoln's Song Sparrow.
- 243. Melospiza palustris (Bartr.) Bd. B 369. C 168. R 233. Swamp Song Sparrow.
- 244. Melospiza fasciata (Gm.) Scott. B 363. C 169. R 231. Song Sparrow.
- 235. C. p. pēr-pāl'-lī-dŭs. Lat. pallidus, pallid, pale, and the intensive particle per.
- 236. C. hěn/-slow-i. To Prof. J. S. Henslow, of Cambridge, Eng.
- 237. C. lĕ-cŏn'-tĭ-ī. To Dr. John L. Le Conte, of Philadelphia.
- 238. Ām-mŏ'-dră-mŭs măr-ĭt'-ĭ-mŭs. Gr. ἄμμος, sand, sea-sand; for the rest of the word, see under Peucedramus, No. 110. The name was originally written as above by Swainson, and we see no necessity of changing it to Ammodromus. It is commonly accented on the penult. Lat. maritimus, maritime; mare, the sea.
- 239. A. m. nig-rēs'-cēns. Lat. nigrescens, present participle of nigresco, I grow black; niger, black.
- 240. A. caūd-ă-cū'-tǔs [kowdakootus not cordakewtus]. Lat. cauda, tail, and acutus, acute, sharp; acus, a pin or point, Gr. ἀκή or ἀκίς, whence the Lat. verb acuo, of which acutus is the perfect participle.
- 241. A. c. něl'-sŏn-ī. To E. W. Nelson, of Illinois, who discovered it near Chicago.

  Not in the orig. ed. Since described by Allen, Pr. Bost. Soc. Nat. Hist., xvii, 1875, p. 93.
- 242. Měl-ŏ-spī/-zã līn/-cŏl-nī. Gr. μέλος, a song, melody, and σπίζα or σπίζη, some small bird; from σπίζω, I chirp. Not to be confounded, as some writers have done, with σπιζίας, a kind of hawk. The σπίζα of Aristotle is supposed to be Fringilla calebs. To Robert Lincoln, sometime a companion of Audubon. In strictness, the above generic name should be pronounced melospeedzah; and the l in lincolni be heard.
- 243. M. păl-ūs'-tris. Lat. palustris, pertaining to a swamp; from palus, a swamp.
- 244. M. fās-cǐ-ā'-tă. Lat. fasciatus, striped; fascis, a bundle of fagots. The allusion is to the indistinct bands upon the tail-feathers; so obsolete are they, in most cases, that it is only recently that it has been admitted that this is the species described by Gmelin. But the markings are as obvious, in some cases, as those on the tail of Chamwa fasciata. The species is given as M. melodia in the orig. ed. of the Check List.

- 245. Melospiza fasciata fallax (Bd.) Ridg. B 367. C 169a. R 231a. Gray Song Sparrow.
- 246. Melospiza fasciata guttata (Nutt.) Ridg. B —. C 169b. R 231d. Oregon Song Sparrow.
- 247. Melospiza fasciata rufina (Brandt) Ridg. B 366. C 169c. R 231c. Rufous Song Sparrow.
- 248. Melospiza fasciata heermanni (Bd.) Ridg. B 364. C 169d. R 231b. Heermann's Song Sparrow.
- 249. Melospiza fasciata samuelis (Bd.) Ridg. B 343, 365. C 169e. R 231c. Samuels' Song Sparrow.
- 250. Melospiza cinerea (Gm.) Ridg. B —. C 169/. R 232. Bischoff's Song Sparrow.
- 251. Peucæa æstivalis (Licht.) Cab. B 370. C 170. R 226.
  Bachman's Summer Finch.
- 252. Peucæa æstivalis illinoensis Ridg. B —, C —, R 226a. Illinois Summer Finch.
- 253. Peucæa æstivalis arizonæ Ridg. B —. C 170a. R 227.

  Arizona Summer Finch.
- 245. M. f. fāl'-lāx. Lat. fallax, false, fallacious, deceitful; in allusion to the perplexity attending the attempt to distinguish it specifically from M. fasciata.
- 246. M. f. gūt-tā'-tă. Lat. guttatus, spotted; gutta, a drop.
- 247. M. f. rū-fī'-nă. Lat. rufus, reddish, of which rufinus is an arbitrary form.
- 248. M. f. heēr'-măn-nī. To Dr. A. L. Heermann, of Philadelphia, sometime naturalist of the Pacific R. R. Survey.
- 249. M. f. săm- $\bar{u}$ - $\bar{e}'$ -lĭs. To E. Samuels. Samuelis is more euphonic than the usual form samuelsi would be.
  - This is M. gouldii of the first ed. of the Check List, the name now adopted having priority.
- 250. M. cĭn-ĕr'-ĕ-ŭs. Lat. cinereus, ashy(-colored): from cinis, genitive cineris, ash. So cinder, in-cin-erate, &c.
  - This is *M. insignis* Bd. of the first ed. of the Check List. As Ridgway has shown (Pr. Nat. Mus., ii, 1880, p. 3) the "Cinereus Finch" of Pennant, on which Gmelin named a *Fringilla cinerea*, from Unalashka, is this bird.
- 251. Peū-caē'-ă aēs-tǐ-vā'-lĭs. Gr. πενκή, a pine; supposed to be from πύκω, to prick, in allusion to the "needles" of this tree. Lat. æstivalis = æstivus, pertaining to summer; æstas, summer.
- 252. P. a. ĭl-lĭ-nŏ-ēn'-sĭs. To the State of Illinois, with the termination -ensis, indicating locality. Illinois is the French corruption of the name by which the aborigines called themselves Illini, "the men."
  - Not in the orig. ed. of the Check List; since described by Ridgway, Bull. Nutt. Club, iv, 1879, p. 219.
- 253. P. a. ă-rī-zō/-naē. To the Territory of Arizona. Arizona is probably a corruption of Orazona, the significance of which is unknown; but it may be observed that zona is the word in the Opata language for the fruit of the mezcal, a characteristic product of the region.

- 254. Peucæa cassini (Woodh.) Bd. B 371. C 170bis. R 228. Cassin's Summer Finch.
- 255. Peucæa ruficeps (Cass.) Bd. B 372. C 171. R 230. Rufous-crowned Summer Finch.
- 256. Peucæa ruficeps boucardi (Scl.) Ridg. B —, C —, R 230a.

  Boucard's Summer Finch.
- 257. Peucæa carpalis Coues. B—. C 171bis. R 229. Bendire's Summer Finch.
- 258. Amphispiza bilineata (Cass.) Coues. B 355. C 172. R 224. Black-throated Finch.
  - 259. Amphispiza belli (Cass.) Coues. B 356. C 173. R 225. Bell's Finch.
  - 260. Amphispiza belli nevadensis Ridg. B —, C 173a. R 225a.

    Nevada Finch.
  - 261. Junco hiemalis (L.) Scl. B 354. C 174. R 217.
    Common Snowbird.
  - 262. Junco hiemalis aikeni Ridg. B —. C 174α. R 216. White-winged Snowbird.
  - 254. P. căs'-sĭn-ī. To John Cassin, of Philadelphia.
  - 255. P. rū'-fī-cēps. Lat. rufus, reddish, and -ceps, a termination denoting the head; from κεφαλή.
  - 256. P. r. boū-cār'-dī. To Adolphe Boucard, a French naturalist, who collected in Mexico and Central America.
  - 257. P. cār-pā'-līs. Gr. καρπόs, fruit, berry, grain; also, the wrist; Latinized as carpus. The derivation supposed to be κάρφω, I gather, as fruit; Lat. carpo, I take, seize. The quasi-Latin carpus is only used as signifying the wrist; the adjective carpulis is an arbitrary form, denoting of or pertaining to the wrist; carpus and carpal are common terms in anatomy. The allusion is to the bright color on the carpal-joint of the bird's wing.
  - 258. Am-phĭ-spī'-ză bĭ-lĭn-ĕ-ā'-tă. Gr. àµφί, on both sides, and σπίζα, a finch; in allusion to the close relation of the genus to those about it. See Melospiza, No. 242. Lat. bilineata, two-lined; bis, twice, and lineatus, striped; linea, a line: see Linaria, No. 207.

    This is the Posspiza bilineata of the first ed. of the Check List.
  - 259. A. běl'-lī. To J. G. Bell, of New York.
  - 260. A. b. něv-ă-dēn'-sīs. To the Territory of Nevada. It were better written nivadensis, in Latin, but is directly from the Spanish adjective nevada, snowy, white as snow; Lat. niveus, snowy, from nix, genitive, nivis, snow. The Territory was named for the snow-capped peaks of its Sierras Nevadas.
  - 261. Jūn'-cō [pronounced yoonco] hī-ĕ-mā'-līs. Lat. juncus, a reed or rush; cf. jungo, I join, junctus, joined; either, reeds growing densely together, or used as withes to bind with?

     For hiemalis, see Anorthura, No. 76.
  - 262. J. h. aī'-kēn-ī. To Charles E. Aiken, of Colorado, its discoverer. This and several other connecting forms of Junco (Nos. 264, 266, 267) are not in the orig. ed. of the Check List.

- 263. Junco hiemalis oregonus (Towns.) Coues. B 352. C 175. R 218. Oregon Snowbird.
- 264. Junco hiemalis annectens (Bd.) Coues. B —. C —. R 219. Pink-sided Snowbird.
- 265. Junco hiemalis caniceps (Woodh.) Coues. B 353. C 176. R 220. Gray-headed Snowbird.
- 266. Junco hiemalis dorsalis (Henry) Coues. B 351. C —. R 221. Red-backed Snowbird.
- 267. Junco hiemalis cinereus (Sw.) Coues. B 350. C —. R 222. Cinereous Snowbird.
- 268. Spizella monticola (Gm.) Bd. B 357. C 177. R 210. Tree Chipping Sparrow.
- 269. Spizella domestica (Bartr.) Coues. B 359. C 178. R 211. Chipping Sparrow; Hairbird.
- . 270. Spizella domestica arizonæ Coues. B —. C 178a. R 211a.

  Arizona Chipping Sparrow.
- 271. Spizella agrestis (Bartr.) Coues. B 358. C 179. R 214. Field Chipping Sparrow.
- 272. Spizella pallida (Sw.) Bp. B 360. C 180. R 212. Clay-colored Chipping Sparrow.
  - 263. J. h. ŏr-ĕ'-gŏ-nŭs. To the Territory of the Oregon. The name is much in dispute; by some derived from the name of a plant (origanum) growing there. It is probably, however, the Algonkin name of the "great river," the Columbia.
  - 264. J. h. ān-nēc'-tēns. Present participle of annecto, I join together, connect, annex; ad, to, and necto, I fasten, join. The bird is very closely related to several others.
  - 265. J. h. cā/-nǐ-cēps. Lat. canus, hoary, grayish white, and -ceps, the termination indicating head, from κεφαλή.
  - 266. J. h. dor-sa'-lis. Lat. dorsum, the back, whence the late Latin adjective, dorsalis.
  - 267. J. h. cīn-ĕr'-ĕ-ŭs. Lat. cinereus, ashy (-colored); cinis, ash.
    The true Mexican cinereus has been found in the United States (Arizona) since the orig. ed. of the Check List appeared.
  - 268. Spīz-ēl'-lă [pronounced speedzaylla] mon-tĭ'-cŏ-lă. An arbitrary diminutive, in Latin form, from Gr. σπίζα, a finch.—Lat. monticola, a mountain-dweller, from mons, genitive montis, a mountain, and colo, I dwell. Mons is from a root min, whence emineo, for example, I project; eminent, imminent, prominent, and also the deponent verb minor, to threaten, whence minatory, &c., are all allied.
- 269. S. dŏm-ēs'-tĭ-că. Lat. domestica, from domus, a house.
  This is S. socialis of the orig. ed. of the Check List.
- 270. S. d. ă-ri-zō'-naē. To the Territory of Arizona. See Peucæa, No. 253.
- 271. S. āg-rēs-tīs. Lat. agrestis, of or pertaining to a field; ager, a field, supposed by some to be related to ago, as something that may be worked; others say from the Gr. ἀγρός, land. This is S. pusilla of the orig. ed. of the Check List.
- 272. S. pāl'-lī-dă. Lat. pallidus, pale, pallid.

- 273. Spizella breweri Cass. B 361. C 180a. R 212. Brewer's Chipping Sparrow.
- 274. Spizella atrigularis (Cab.) Bd. B 362. C 181. R 215. Black-chinned Chipping Sparrow.
- 275. Zonotrichia albicollis (Gm.) Bp. B 349. C 182. R 209. White-throated Crown Sparrow.
- 276. Zonotrichia leucophrys (Forst.) Sw. B 345. C 183. R 206. White-browed Crown Sparrow.
- 277. Zonotrichia leucophrys intermedia Ridg. B 346. C 183b. R 207a. Intermediate Crown Sparrow.
- 278. Zonotrichia gambeli Nutt. B 346. C 183a. R 207. Gambel's Crown Sparrow.
- 279. Zonotrichia coronata (Pall.) Bd. B 347. C 184. R 208. Golden Crown Sparrow.
- 280. Zonotrichia querula (Nutt.) Gamb. B 348. C 185. R 205. Harris's Crown Sparrow.
- 281. Chondestes grammicus (Say) Bp. B 344. C 186. R 204, 204a.

  Lark Finch.
- 273. S. brēw'-ĕr-ī. To Thomas Mayo Brewer, of Boston, long the leading oölogist of the United States.
  - This is given in the first ed. of the Check List as a var. of pallida.
- 274. S. ā-trī-gŭī-ā'-rīs. Lat. ater, atra, atrum, black; and gularis, pertaining to the throat; gula, the throat, gullet.
- 275. Zō-nō-trīch'-ĭ-ā [pronounced Dzonotreekeya] āl-bī-cōl'-līs. Gr. ζώνη, a girdle, band, zone, and τριχάς or τριχιάς, some kind of bird; in allusion to the conspicuously banded heads of sparrows of this group. Or, the latter part of the word may be directly from τριχίας (θρίξ, genitive τριχός), hairy; i.e., having the head striped. Lat. albicollis, white-throated; albus, white, and collum, the collar, neck.
- 276. Z. leū-cō'-phrys. Gr. λευκός, white, and ὀφρύς, eyebrow.
- 277. Z. 1. in-tēr-měd'-ī-ă. Lat. intermedius, intermediate, between two things; inter, between, among, and medius, middle; related to Gr. μέσοs, of same meaning.
  Not in the orig. ed.; since discriminated both from leucophrys and from gambeli.
- 278. Z. găm'-bĕl-ī. To William Gambel, of Philadelphia, one of the pioneers in Californian ornithology.

  In the orig. ed. this is given as a var. of leucophrys; since decided to be distinct.
- 279. Z. cŏr-ō-nā'-tă. Lat. coronatus, crowned, participle of corono, I crown; corona, a crown. Coronis or Κορώνις was the name of a Thessalian princess; also, a scroll with which writers marked the end of a piece of writing—"finis coronat opus." Corone οr κορώνη was also a crow or raven, into which the princess was fabled to have been transformed by her spouse Apollo, and survives in ornithology in the term Corvus corone L.
- 280. Z. quer'-ŭ-lă. Lat. querulus or querulosus, plaintive, querulous; from queror, to complain, lament.
- 281. Chōn-dēs'-tēs grām'-mǐ-cǔs. Gr. χόνδρος, cartilage; also, a kind of grain; -estes is from the root έδω, I eat. Is not the word more properly to be written chondrestes? We suppose it to be masculine. Lat. grammicus, from gramma, a line, word, mark, in allusion to the

- 282. Passerella iliaca (Merr.) Sw. B 374. C 188. R 235. Fox Sparrow.
- 283. Passerella iliaca unalascensis (Gm.) Ridg. B 375. C 189. R 235a. Townsend's Fox Sparrow.
- 284. Passerella iliaca schistacea (Bd.) All. B 376. C 189a. R 235c. Slate-colored Fox Sparrow.
- 285. Passerella iliaca megarhyncha (Bd.) Hensh. B —. C —. R 2356. Large-billed Fox Sparrow.
- 286. Calamospiza bicolor (Towns.) Bp. B 377. C 190. R 256. Lark Bunting.
- 287. Spiza americana (Gm.) Bp. B 378. C 191. R 254. Black-throated Bunting.
- 288. Spiza townsendi (Aud.) Ridg. B 379. C 192. R 255. (?) Townsend's Bunting.
  - stripes on the head; Gr. γράμμα, γραμμικόs. Usually written grammaca or grammacus, for which there is no authority. And even the corrected form is bad enough; for grammicus does not mean lineatus, striped, marked with lines, but linearis, linear, having the quality of a line.
  - 282. Pās-sĕr-ēl'-lā ī-lī'-ā-că. An arbitrary diminutive of Lat. passer, like spizella from spiza.

     For iliaca, see Turdus iliacus, No. 4. Applicability of the name inobvious; it may be intended to note some resemblance to the thrush in mention, or refer to the conspicuous markings of the flanks.
- 283. P. i. ū-nă-lās-cēn'-sĭs. The name of the Aleutian Island for which this species is named, has no settled orthography: Unalashka, Unalaschka, Unalascha, Ouna-, Oona-, Aoona-, Aona-, &c. In the present case, Pennant wrote Unalascha Bunting, of which Gmelin made Emberiza unalaschcensis, and was nearly followed by Ridgway; but the word may be euphonized as above, just as we have alascensis as the name of a wren, No. 78.

This stands as Passerella townsendii in the orig. ed.

284. P. i. schīs-tā'-cĕ-ă. Lat. (late) schistaceus, slaty, relating to slate; in this case, in color; schistos or σχιστόs, split, cleft, or fissile, capable of easy cleavage, as slate-stone is. The same stem is seen in schism, schismatic.

This stands as P. townsendii var. schistacea in the orig. ed.

285. P. i. měg-ă-rhỹn/-chă. Gr. μέγα, great, large, and ἡύγχος, Lat. rhynchus, snout, muzzle, beak. More exactly to be written megalorhyncha.

Not in the orig. ed. of the Check List; since revived by H. W. Henshaw.

- 286. Căi-ă-mō-spīz'-ă bī'-cŏi-ŏr. Lat. calamus or Gr. κάλαμοs, a reed, rush, cane, flag; and spiza. See under Passer, No. 192, and Melospiza, No. 242.—Lat. bicolor, two-colored; bis, twice; in allusion to the black-and-white of-the-male.
- 287. Spīz'-ă [pronounced Speedzah] ăm-ĕr-ĭ-cā'-nă. See under Melospiza, No. 242.

This stands as Euspiza amer. in the orig. ed. For the change, see Ridg., Pr. Nat. Mus., ii, 1880, p. 3.

288. S. town'-sĕnd-ī. To J. K. Townsend.

Given as Euspiza towns. in the orig. ed. No second specimen of this alleged species is known, and it is not improbable that the type came from an egg laid by S. americana. But even such immediate ancestry would not forbid recognition of "specific characters;" the solitary bird having been killed, it represents a species which died at its birth.

- 289. Zamelodia ludoviciana (L.) Coues. B 380. C 193. R 244. Rose-breasted Song Grosbeak.
- 290. Zamelodia melanocephala (Sw.) Coues. B 381. C 194. R 245. Black-headed Song Grosbeak,
- 291. Guiraca cœrulea (L.) Sw. B 382. C 195. R 246. Blue Grosbeak.
- 292. Passerina ciris (L.) Gray. B 384. C 196. R 251. Painted Finch; Nonpareil.
- 293. Passerina versicolor (Bp.) Gray. B 385. C 197. R 250. Versicolor Painted Finch.
- . 294. Passerina amœna (Say) Gray. B 386. C 198. R 249. Lazuli Painted Finch.
- 295. Passerina cyanea (L.) Gray. B 387. C 199. R 248. Indigo Painted Finch; Indigo-bird.
- 296. Spermophila moreleti Pucheran. B 388. C 200. R 252. Morelet's Seed-eater.
- 297. Phonipara zena (L., 1758) Bryant. B —. C 201. R 253. (!W.I.) Black-faced Finch.
- 289. Zā-měl-ō'-dǐ-ă lū-dō-vǐ-cǐ-ā'-nă. Gr. ζά, an intensive particle, and μελφδία, singing, melody; in allusion to the strikingly rich song. — To Louisiana; see Thryothorus, No. 68. rlain, probably This is given as Goniaphea lud. in the orig. ed. For the change, see Coues, Bull. Nutt. Club, v, 1880, p. 98.
- 290. Z. měl-ăn-ŏ-cĕph/-ă-lă. Gr. μέλας, feminine μέλαινα, neuter μέλαν, black; κεφαλή, the head.
  - 291. Guĭr'-ă-că [pronounced Gweerahcah] coē-rŭl'-ĕ-ă. The generic word is barbarous, from some South American vernacular, and of uncertain meaning. It occurs, with several similar words, as guira, in Marcgrave. We mark the accent (for which there is no authority) as usually heard. — For carulea, see Polioptila, No. 36.
  - 292. Pās-sĕr-ī'-nă cī'-rĭs. Passerina, formed from Passer, as Passerella and Passerculus also are. - Ciris, Gr. κείριs, a kind of bird, into which Scylla, daughter of Nisus, is fabled to have been changed. - Nonpareil = "the incomparable." For use of Passerina, instead of Cyanospiza of the orig. ed., see Coues, Bull. Nutt.
  - Club, v, 1880, p. 96. 293. P. vēr-sī'-cŏl-ŏr. Lat. versicolor, of changing or versatile colors, many-colored, party-colored; verso, I turn about, change, am occupied with, versed in, &c.; color, color.
  - 294. P. ă-moē'-nă [ahmwaynah]. Lat. amæna, delightful, charming, dressy.
  - 295. P. cỹ-ăn'-ĕ-ă. Lat cyaneus, Gr. κυάνεος or κύανος, dark blue.
  - 296. Spēr-mo'-phǐ-lǎ more-lět'-ī. Gr. σπέρμα, genitive σπέρματος, a seed; from σπείρω, equal to the Lat. sparge, I sow seed: and  $\phi l \lambda o s$ , from  $\phi l \lambda \epsilon \omega$ , I love. The word is contracted; the full form is spermatophila. — To — Morelet, a French naturalist.
  - 297. Phō-nǐ'-pă-ră zē'-nă. Gr. φωνή, a sound, the voice; φημί, I speak; the English "phonetic" is from the same. The rest of the word appears to be from Lat. pario, I bring forth, beget, produce, having the same root as is seen in primi-para, par-turient, vivi-par-ous, &c.; if so, the word is a hybrid which would be better written sonipara or vocipara. The meaning of zena we do not know; we suppose it not to be of Greek or Latin derivation.

This is given as P. bicolor in the orig. ed. of the Check List, after Fringilla bicolor L.,

1766; but it seems that F. zena L., 1758, is the prior tenable name.

- 298. Pyrrhuloxia sinuata Bp. B 389. C 202. R 243. Texas Cardinal Grosbeak.
- 299. Cardinalis virginiana Bp. B 390. C 203. R 242. Cardinal Grosbeak; Virginia Redbird.
- 300. Cardinalis virginiana ignea (Bd.) Coues. B —. C 203a. R 242a. Fiery-red Cardinal Grosbeak.
- 301. Pipilo erythrophthalmus (L.) V. B 391. C 204. R 237. Towhee Bunting; Chewink.
- 302. Pipilo erythrophthalmus alleni Coues. B—. C 204a. R 237a. White-eyed Towhee Bunting.
- 303. Pipilo maculatus oregonus (Bell) Coues. B 392. C 205. R 2386. Oregon Towhee Bunting.
- 304. Pipilo maculatus arcticus (Sw.) Coues. B 393. C 205a. R 238. Arctic Towhee Bunting.
- 305. Pipilo maculatus megalonyx (Bd.) Coues. B 394. C 2056. R 238a. Spurred Towhee Bunting.
- 298. Pyr-rhŭ-lōx'-I-ă sĭn-ŭ-ā'-tă. A forcible combination of Pyrrhula and Loxia: see these words, Nos. 190 and 199; or may be said to be more properly compounded of pyrrhuls, πυβροs, fiery-red, and λοξίας; in which event, it should be written pyrrholoxia.— Lat. sinuatus, bent, bowed, curved, as the bill of the bird is; from sinuo, the verb; sinus, the noun, a curve, bending, bay.
- 299. Cār-dīn-ā'-līs vīr-gīn-ĭ-ā'-nā. Lat. cardinalis, pertaining to a door-hinge; cardo, genitive cardinis, a door-hinge; hence, that upon which something turns or depends; as, cardinal points of the compass; hence, any important thing or person; applied with obvious signification to the chief officials of the Pope. These ecclesiastical dignitaries wear red; hence the phrase "cardinal-red." The term is applied to the bird as descriptive of its rich red color. As a Latin word, cardinalis is only an adjective; used substantively, its gender is either masculine or feminine. We take the latter, because most words ending in is- are feminine.— Lat. virginiana, of Virginia, euphemistically named for Elizabeth, daughter of Henry VIII.
- 300. C. v. ig'-nĕ-ă. Lat. igneus, fiery, flaming; said of color as well as of other properties; ignis, fire.
- 301. Pi'-pīl-ō ĕ-ryth-rōph-thāl'-mŭs. Vieillot, in forming the word, wrote both pipilo and pipillo. It is a Latin verb, meaning, like pipio, I pip, peep, chirp. Notice the accentuation and quantity of the vowels.—Gr. ἐρνθρός, red or reddish; ἐρεύθω, I redden; ὀφθαλμός, the eye, from ὅπτομαι, a verb obsolete in the present, or ὀράω, I see; we find both words in "ophthalmic," "optic." The species is red-eyed.—The curious English words "towhee" and "chewink" are onomatopæic: that is, coined to imitate the sound of the bird's voice.
- 302. P. e. ăl'-lĕn-ī. To Joel Asaph Allen, of Cambridge, Mass., one of the leading naturalists of the United States.
- 303. P. mă-cŭi-ā'-tŭs ŏr-ĕ-gō'-nŭs. Lat. maculatus, spotted; macula, a spot.— To the Oregon River. Quantity of the penult in question, perhaps better ore'gŏnus.

  The stock species, P. maculatus, is not North American.
- 304. P. m. ārc'-tĭ-că. See Sialia, No. 29.
- 305. P. m. mě-găl'-ŏ-nȳx. Gr. μεγάλη (feminine of μέγαs), large, great, and ὅνυξ, Lat. onyx, a nail, claw, talon. The word is commonly accented on a long penult; a practice perhaps defensible on the ground that megalŏ-ŏnyx = megalōnyx.

- 306. Pipilo fuscus mesoleucus (Bd.) Ridg. B 397. C 206. R 240. Brown Towhee Bunting; Cañon Bunting.
- 307. Pipilo fuscus albigula (Bd.) Coues. B C 206a. R 240a. White-throated Towhee Bunting.
- 308. Pipilo fuscus crissalis (Vig.) Coues. B 396. C 206b. R 240b. Crissal Towhee Bunting.
- 309. Pipilo aberti Bd. B 395. C 207. R 241.

  Abert's Towhee Bunting.
- 310. Pipilo chlorurus (Towns.) Bd. B 398. C 208. R 239. Green-tailed Towhee Bunting.
- 311. Embernagra rufovirgata Lawr. B 373. C 209. R 236. Green Finch.
- 312. Dolichonyx oryzivorus (L.) Sw. B 399. C 210. R 257. Bobolink; Reed-bird; Rice-bird.
  - 306. P. fūs'-cǔs měs-ŏ-leū'-cǔs. Lat. fuscus, fuscous, dark, dusky, like furvus; both allied to Gr. δρφόs, of same meaning, from δρφνή, night or darkness? Gr. μέσος, middle, λευκός, white; in allusion to the color of the middle under parts. This word is derived from λεύσσω or γλαύσσω, I shine; this from ἀγλαία, splendor, the name of one of the Muses.

    This is given as P. fuscus in the orig. ed.; but the bird of Arizona is said to be distinguishable from the Mexican stock species.
  - 307. P. f. āl-bī'-gŭ-lā. [Not albigew'ler.] Lat. albus, white; gula, throat. This is one of numberless cases where the termination of the word is in question. Albigula may be taken as a feminine noun, and left in this form, whatever the gender of the word with which it is associated; or it may be considered an adjective in -us, -a, -um, and made masculine to agree with P. fuscus. There is ample authority and precedent for the latter course, which our taste disinclines us to take. English affords a parallel latitude of construction, as when we say indifferently "yellow-rump warbler" or "yellow-rumped warbler," "Carolina chickadee" or "Carolinian chickadee." A better form than either albigulus or albigula would be albigularis.
  - 303. P. f. crīs-sā'-līs. Late Lat. crissalis, pertaining to the crissum, or under-tail coverts, which in this bird are highly colored. There are no such classic words, they having been invented by Illiger in 1811; but there is a verb crisso, expressing a certain action of the parts.
  - 309. P. ă'-bert-î. To Lieutenant J. W. Abert, of the U. S. Army, who discovered it.
  - 310. P. chlō-rū'-rŭs. Gr. χλωρδε, green, from χλδα, green grass; οὖρα, tail.
- 311. Em-bēr-nā'-gră rū-fŏ-vīr-gā'-tă. Embernagra is a villanous word, concocted by Lesson out of Emberiza and Tanagra. Emberiza, a bunting, is a word the derivation of which is not classic. It is said, doubtless correctly, to be Latinized from the O. H. G. Embritz; "Charleton (1668) has Embryza" (Wharton's MS.); and we may add that there were various other forms of the word before it settled into the present one. There are Latin words Tanagra and Tanagra; but these are geographical proper names, having nothing to do with the present case. Tangara or Tanagra is a South American vernacular word. —Lat. rufus, rufous, reddish, and virgatus, literally, made of twigs; from virga, a rod, switch, the application being the stripes with which the bird is marked. Commonly written rufivirgata: see Lophophanes, No. 42.
- 312. Dŏl-ĭch'-ŏ-nȳx ŏ-ry-zĭ'-vŏ-rŭs. Gr. δολιχός, long, and ὅνυξ, a nail, claw, talon. The gender is in question; but the Greek ὅνυξ, Lat. onyx, is masculine, though Latin words in -yx are usually feminine. The usual pronunciation is dolichō'nyx: but see Pipilo, No. 305. Gr. ὅρυζα, or Lat. oryza, rice, and νονο, I devour.

- 313. Molothrus ater (Bodd.) Gray. B 400. C 211. R 258. Cowbird.
- 314. Molothrus ater obscurus (Gm.) Coues. B—. C 211a. R 258a. Dwarf Cow-bird.
- 315. Molothrus æneus Cab. B —, C —, R 259. Bronzed Cowbird.
- 316. Agelæus phœniceus (L.) V. B 401, C 212. R 261. Red-winged Marsh Blackbird.
- 317. Agelæus phœniceus gubernator (Wagl.) Coues. B 402. C 212a. R 261a. Red-shouldered Marsh Blackbird.
- 318. Agelæus tricolor Nutt. B 403. C 212b. R 262.

  Red-and-white-shouldered Marsh Blackbird.
- 313. Mō-lō'-thrŭs ā'-tĕr. Unde derivatur? The orthography and etymology of molothrus are alike in dispute. Swainson himself says, "μολοθρος, qui non vocatus alienas aedes intrat;" that is, an uninvited guest. There being no such Greek word as μολοθρός, but there being a good Greek word μολοβρός, meaning one who roams in quest of food, a vagabond, a beggar, a parasite, a "tramp" (as we should say now), and therefore exactly answering to Swainson's explanation of his molothrus, it has been supposed by Cabanis that Swainson meant to say molobrus, and the word has consequently been changed. Though this is very true, it is also to be observed that Swainson wrote molothrus more than once, showing it not to be a misprint or other mistake, and that, further, it is quite possible to construct the word molothrus from μῶλος and θρώσκω (θορεῖν, θόρω, θόω), and answer all the conditions of Swainson's definition; molothrus being, in this case, a bird which takes uninvited possession of other birds' nests, and there leaves an alien egg in mockery of the rightful owners. We therefore see no necessity to replace molothrus by molobrus. The first o is marked long as being Gr. ω, the second as lengthened by position.

This stands in the orig. ed. as M. pecoris, corrected in a footnote.

314. M. a. ōb-scū'-rŭs. Lat. obscurus, obscure, dark; obscuro, I darken; Gr. σκιά, shadow, shade.

This stands as M. pecoris var. obscurus in the orig. ed.

- 315. M. a. aē'-nĕ-ŭs. Lat. æneus, of brass, brassy, brazen, bronzed; from æs, genitive æris, brass.

  Not in the orig. ed.; since discovered by J. C. Merrill, in Texas.
- 316. A-gĕl-aē'-ŭs phoē-nī'-cĕ-ŭs. Gr. ἀγελαῖος, pertaining to flocks and herds, from ἀγέλη, a flock: this from ἀγείρω, I assemble, from ἄγω, I lead; in allusion to the gregariousness of these Blackbirds. Gr. φοινίκεος, or Lat. phæniceus, deep red; "a color first introduced into Greece by the Phænicians." The fabulous bird Phænix, and the name of Phænician, and the word for flame-color, are all the same, φοίνιξ. This itself is a radical word, but related through φοινός, φόνος, with φένω, φάω, I kill, slay, as if the idea of the whole set of words were that of murder, from its traditional color of blood. The obvious application is to the scarlet on the wings.
- 317. A. p. gŭb-ēr-nā'-tŏr. Lat. gubernator, Gr. κυβερνήτης (cybernetes), a pilot, helmsman; gubernum or gubernaculum, a rudder, tiller; guberno, Gr. κυβερνάω or κυβερνώ. I steer a ship; hence, to direct or govern in general. Govern, governor, are directly from guberno, and the actual Latin lingers in gubernatorial. The implication is the red shoulder-knots or epaulettes of the bird, as if signs of rank or command.
- 318. A. trī'-cŏl-ŏr. Lat. tricolor, three-colored; tres, three, becoming in composition tri
  This stands as A. phæniceus var. tricolor in the first ed., but proves to be sufficiently distinct.

- 319. Xanthocephalus icterocephalus (Bp.) Bd. B 404. C 213. R 260. Yellow-headed Swamp Blackbird.
- 320. Sturnella magna (L.) Sw. B 406. C 214. R 263. Meadow Starling; Field-lark.
- 321. Sturnella magna mexicana (Scl.) Ridg. B —. C —. R 263a. Mexican Meadow Starling.
- .322. Sturnella magna neglecta (Aud.) Allen. B 407. C 214a. R 264. Western Meadow Starling.
- 323. Icterus vulgaris Daud. B 408. C —. R 265. (! W.I.)
  Troupial.
- 324. Icterus spurius (L.) Bp. B 414. C 215. R 270. Orchard Oriole.
  - 325. Icterus spurius affinis (Lawr.) Coues. B —. C 215a. R —. (?)
    Texas Orchard Oriole.
  - 326. Icterus galbula (L., 1758) Coues. B 415. C 216. R 271.
    Baltimore Oriole.
  - 319. Xān-thŏ-cĕph'-ăl-ŭs īc-tĕr-ŏ-cĕph'-ăl-ŭs. Gr. ξανθός, bright yellow.—Gr. ἴκτερος, or Lat. icterus, see Icteria, No. 144. Related apparently to ἴκω, I attack, as disease does.
  - 320. Stūr-nēl'-lă māg'-nă. Diminutive of Lat. sturnus, a starling; as spizella from spiza.— Lat. magnus, great, large; root mag, as seen in Gr. μέγαs; whence also mactus, magnified, glorified: magi, magician, magic, are all allied.
  - 321. S. m. mēx-ĭ-cā/-nă. Latinized Mexican. See Sialia, No. 28.

    Not in the orig. ed. Since discovered in Texas by J. C. Merrill.
  - 322. S. m. nēg-lēc'-tă. Lat. neglecta, neglected, that is, not chosen, not heeded; from nec, not, and lego, I choose, select, &c. See Parus, No. 51.
  - 323. Ic'-tĕr-ŭs vūl-gā'-rĭs. See Sturnus, No. 363. Troupial or troopial, from the Fr. trouper, is simply trooper, the bird that goes in troops.

Not in the orig. ed. Said to straggle to Southern States. No late case of its so doing.

- 324. I. spŭ'-rĭ-ŭs. For Icterus, see Icteria, No. 144, and Xanthocephalus, No. 319. Lat. spurius, illegitimate, bastard, spurious; related to the Gr. σπορά, seed, generation, birth, &c., σπείρω, I sow seed. The bird was formerly called "Bastard Baltimore Oriole," whence the undeserved Linnæan name.
- 325. I. s. āf-fi'-nīs [accent the penult]. Lat. affinis, ad, and finis, allied, affined.

  This subspecies is very slightly distinguished from its stock.
- 326. I. gāl'-bŭ-lã. Lat. galbula or galgula, some small yellow bird of the ancients; doubtless derived from some word signifying yellow; there are Latin words galbus, galbanus, Germ. gelb, &c., of such meaning. The curious English word oriole, for which no derivation is given in some standard works, has evidently a similar reference to the color yellow, being equivalent to aureole; Lat. aurum or Gr. αδρον, gold: such form of the word for gold, with or-instead of aur-, is seen in the Fr. or. "Baltimore," the former specific name of the bird, is not directly from the city of that name, but from the name of Sir George Calvert, first Baron of Baltimore, the colors of the bird being chosen by him for his livery, or, as Catesby has it (N. H. Car., i, 1731, p. 48), the bird being named from its resemblance in color to the Lord's coat of arms "which are Paly of six Topaz and Diamond, a Bend, interchang'd." The name baltimore, L., 1766, as given in the orig. ed. of the Check List, is antedated by Coracias galbula L., 1758; see Coues, Bull. Nutt. Club, April, 1880, p. 98.

- 327. Icterus bullocki (Sw.) Bp. B 416. C 217. R 272. Bullock's Oriole.
- 328. Icterus cucullatus Sw. B 413. C 218. R 269. Hooded Oriole.
- 329. Icterus parisiorum Bp. B 411. C 219. R 268. Scott's Oriole.
- 330. Icterus melanocephalus auduboni (Gir.) Coues. B 409. C 220. R 266. Audubon's Black-headed Oriole.
- 331. Scolecophagus ferrugineus (Gm.) Sw. B 417. C 221. R 273. Rusty Grackle.
- 332. Scolecophagus cyanocephalus (Wagl.) Cab. B 418. C 222. R 274. Blue-headed Grackle.
- 333. Quiscalus macrurus Sw. B 419. C 223. R 275. Great-tailed Crow Blackbird.
- 327. I. bŭl'-lŏck-ī. To William Bullock, sometime a collector in Mexico, and proprietor of a famous museum in London.
- 328. I. cŭ-cūl-lā'-tŭs. Lat. cucullatus, hooded; cuculla, a kind of hood or cowl fastened to a garment, to be drawn over the head.
- 329. I. păr-īs-ī-ō'-rŭrn. Lat. Parisiorum, of the Parisians. The Parisii were a people of Gaul, settled on the river Senones, now the Seine; their chief city, Lutetia, called also Lutetia Parisiorum and Parisii, is now Paris. There is no applicability of the name to the bird: Bonaparte probably so called it from national vanity, or because he found a specimen in a museum in Paris. The name is commonly but wrongly written parisorum.
- 330. I. měl-ăn-ŏ-cěph/-ăl-ŭs aūd/-ŭ-bŏn-ī. Gr. μέλας, feminine μέλαινα, black; and κεφαλή, head. Το J. J. Audubon.
- 331. Scō-lē-cŏ'-phǎ-gùs fēr-rū-gǐn'-ĕ-ùs. Gr. σκωληκοφάγοs, a worm-eater; σκώληξ, genitive σκώληκοs, a worm, and φάγω, I eat. It is also a Latin word, scolex, worm. Lat. ferrugineus, rusty-red, color of iron-rust; from ferrugo, iron-rust; ferrum, iron. The curious English word grackle or grakle is anglicized from Lat. graculus or graculus, a very uncertain
  - bird, by some supposed to be the jackdaw, by others the cormorant or sea-crow; and the Latin word itself is supposed to be merely in imitation of a hoarse croak, gra, gra. See what is said under Querquedula, No. 714.
- 332. S. cy-ăn-ŏ-ceph/-ăl-ŭs. Gr. κύανος, or Lat. cyaneus, blue; and κεφαλή, head.
- 333. Quīs'-că-lŭs māc-rū'-rŭs. Unde derivatur quiscalus? We have no proof whence it comes or what it means: it varies in form, as quiscala, quiscula. Mr. W. C. Avery asks: "Is quiscalus an onomatopæon? I can find no Latin or Greek word like it." Mr. H. T. Wharton observes: "Quiscalus seems a native name; if it is, the termination-us only obscures its origin without Latinizing it." Professor A. Newton remarks at greater length: "Quiscalus was doubtless taken by Vieillot from the Gracula quiscula of Linnæus (S. N., ed. 10, p. 109). I cannot find this word or any thing like it in any older author; but I have an instinctive conviction that it must occur somewhere; for, as far as my studies of Linnæus's work go, they show me that he did not invent names. From his printing the word in both eds. (10th and 12th) with a capital initial letter, it is obvious that he regarded it as a substantive, and I should think he must have found it in some book of travels as the local name of a bird. The word seems to me Spanish or quasi-Spanish—say Creole—and the regular Castilian quisquilla, which dictionaries explain to be a trifling dispute, suggests a meaning, especially when one reads of the noisy and fussy bickerings of your Boat-tails." If, as seems highly probable, we are here on the

- 334. Quiscalus major V. B 420. C 224. R 277.

  Boat-tailed Crow Blackbird; Jackdaw.
- 335. Quiscalus purpureus (Bartr.) Licht. B 421. C 225. R 278.

  Purple Crow Blackbird; Purple Grackle.
- 336. Quiscalus purpureus æneus Ridg. B —. C —. R 278b. Bronzed Crow Blackbird.
- 337. Quiscalus purpureus aglæus (Bd.) Coues. B 422. C 225a. R 278a. Florida Crow Blackbird.
- 338. Corvus corax L. B 423. 424. C 226. R 280.
- 339. Corvus cryptoleucus Couch. B 425. C 227. R 281.
  White-necked Raven.
- 340. Corvus frugivorus Bartr. B 426. C 228. R 282. Common American Crow.
- 341. Corvus frugivorus floridanus (Bd.) Coues. B 427. C 228a. R 282a. Florida Crow.
- 342. Corvus caurinus Bd. B 428, C 228b, R 282b.
  Northwestern Crow.

right track of the word, we may perhaps go a step further, and trace the undoubtedly barbarous word quisculus through quisquilla to the similar Lat. quisquillæ, which the lexicons give as meaning refuse, dregs, or other trifling worthless matters; as we might say, riff-raff, rag-tag; and such would not be wholly inappropriate to these vagabond troopers, so common everywhere as to come under the contempt of familiarity. — Gr. μακρός, long, large, and οὖρα, tail.

- 334. Q. mā'-jŏr. Lat. major, greater, comparative of magnus.
- 335. Q. pūr-pŭr'-ĕ-ŭs. See Carpodacus, No. 194.
- 336. Q. p. aē'-nĕ-ŭs. See Molothrus, No. 315.

  Not in the orig. ed. of the Check List: since recognized.
- 337 Q. p. āg-laē'-ŭs. Gr. ἀγλαίος or ἀγλαίος, shining, from ἀγλαία, splendor; also the name of one of the Muses. Obs.— Not to be confounded with agelæus, which see, No. 316.
- 338. Cōr'-vǔs cŏr'-āx. Lat. corvus, a crow. Lat. corax or Gr. κόραξ, a raven. Corvus is by some considered an onomatopæon, and referred through the Gr. κράζω, κράζω, to croak, back to a Sanscrit root of same signification. Corax is more obviously a word of similar formation, as may also be the English crow.
- 339. C. cryp-to-leū'-cūs. Gr. κρυπτόs, hidden (with which compare Eng. crypt), and λευκόs, white; the allusion being to the concealed white at the bases of the feathers of the neck.
- 340. C. frū-gǐ'-vŏ-rŭs. Lat. frugivorus, fruit-eating; frux, genitive frugis, fruit, and voro, I devour. Frux is from fruor, fruitus, fructus, as it is something that may be enjoyed. Voro is rooted in βορ, as seen in βορά, food, and βόσκω (βόω), I eat.

  This is given as C. americanus in the orig. ed. of the Check List.
- 341. C. f. flo-ri-da'-nus. To Florida. Flora, Goddess of flowers; flos, a flower.
- 342. C. caū-rī'-nūs. There is no such Latin word. Caurinus has been supposed to be equivalent to corvinus, crow-like, but is directly derived from caurus, the North-west wind, the species having been discovered on the North-west coast of the United States.

This stands as *C. americanus* var. *caurinus* in the orig. ed.: it has been redetermined to be distinct, as originally described by Baird.

- 343. Corvus maritimus Bartr. B 429. C 229. R 283. Fish Crow.
- S44. Picicorvus columbianus (Wils.) Bp. B 430. C 230. R 284. Clarke's Nutcracker.
- 345. Gymnocitta cyanocephala Maxim. B 431. C 231. R 285. Blue Nutcracker.
- 346. Psilorhinus morio (Wagl.) Cab. B 444. C 232. R 288. Brown Jay.
- 347. Pica rustica hudsonica (Cab.) Ridg. B 432. C 233. R 286. American Magpie.
- 348. Pica rustica nuttalli (Aud.) Coues. B 433. C 233a. R 287. Yellow-billed Magpie.
- 349. Cyanocitta cristata (L.) Strickl. B<sub>.</sub>434. C<sub>.</sub>234. R<sub>.</sub>289. Blue Jay.
- 350. Cyanocitta stelleri (Gm.) Strickl. B 435. C 235. R 290. Steller's Jay.
- 351. Cyanocitta stelleri annectens (Bd.) Ridg. B —. C —. R 290b. (?)
  Connective Jay.
- **343. C.** măr-ĭt'-ĭ-mŭs. See *Ammodramus*, No. 238. This stands as *C. ossifragus* in the orig. ed.
- 344. Pī-cĭ-cōr'-vŭs cŏ-lūm-bĭ-ā'-nŭs. The generic name is compounded of *pica* and *corvus*: see these words, Nos. 347 and 338.—The specific name refers to the Columbia River, whence Lewis and Clarke first brought specimens.
- 345. Gym-nŏ-cīt'-tă cy-ăn-ŏ-cĕph'-ā-lā. Gr. γυμνός, naked; in allusion to the nostrils being exposed, as is unusual in this family; κίττα οι κίσσα, a jay.— See Scolecophagus, No. 332.
- 346. Psǐ-lŏ-rhī'-nŭs mŏr'-ĭ-ō. Gr. ψιλόs, smooth, bare, bald, in allusion to the uncovered nostrils, from ψίω; and βίs, genitive βινόs, the nose.—The specific name is morio, "a dark brown gem," in allusion to the color, which is remarkable in this group of birds.
- 347. Pī'-că rūs'-tī-că hǔd-sŏn'-ī-că. Lat. pica, a magpie. It is supposed by some to be for piga, that equivalent to pigta or picta, from pingo, I paint; hence signifying painted, speckled, pied. The same dubious etymology is ascribed to the masculine form of the word, picus, which see, No. 433. Lat. rusticus, rustic, rural, from rus, the country as distinguished from the city. —To Hudson's Bay, named after Henry Hudson, the explorer. This stands as P. melanoleuca hudsonica in the orig. ed.; but rustica has long priority.
- 348. P. r. nŭt'-tăl-lī. To Thomas Nuttall, the botanist and ornithologist.

  This stands as P. melanoleuca nuttalli in the orig. ed.
- 349. Cỹ-ăn-ŏ-cīt'-tă crīs-tā'-tă. Gr. κυανός, cyaneus, blue, and κίττα, a jay.—Lat. cristatus, crested; crista, a crest; related to cresco, I grow, and crinis, hair, through a common root.

  For use of Cyanocitta instead of Cyanurus, as in the orig. ed., see Coues, Bull. Nutt. Club, v, 1880, p. 98.
- 350. C. stěl'-ler-i. To G. W. Steller, surgeon and naturalist.
- 351. C. s. ān-nēc/-tēns. Lat. annectens (ad and necto, to bind), annexing, annectant, connecting, tying together; because this subspecies is intermediate between others of the same stock, serving to link them to each other.

Not in the orig. ed. of the Check List.

- 2352. Cyanocitta stelleri macrolopha (Bd.) Coues. B 436. C 235a. R 290c. Long-crested Jay.
- 353. Cyanocitta stelleri frontalis (Ridg.) Coues. B —. C 235b. R 290a. Blue-fronted Jay.
- 354. Aphelocoma floridana (Bartr.) Cab. B 439. C 236. R 291. Florida Jay.
- 355. Aphelocoma fioridana woodhousii (Bd.) Allen. B 438. C 236a, R 292. Woodhouse's Jay.
- 356. Aphelocoma floridana californica (Vig.) Coues. B 437, C 236b, R 293, Californian Jay.
- 357. Aphelocoma ultramarina arizonæ Ridg. B 440. C 237. R 295. Arizona Jay.
- 358. Xanthura luxuriosa (Less.) Bp. B 442. C 238. R 296. Rio Grande Jay.
- 359. Perisoreus canadensis (L.) Bp. B 443. C 239. R 297. Canada Jay.
- 360. Perisoreus canadensis fumifrons Ridg. B —. C —. R 297b. Alaskan Jay.
- 352. C. s. māc-rō'-lō-phā. Gr. μακρός, long, and λόφος, a mane, crest, comb, from λέπω, as is also λεπίς, λέπος, a scale, and many similar words. Usually pronounced macrolō'pha.
- 353. C. s. fron-ta'-lis. Lat. frontalis, relating to frons, the forehead, front.
- 354. A-phĕ-lŏ'-cŏ-mă flō-rǐ-dā'-nă. Gr. ἀφελής, smooth, sleek, and κόμη, Lat. coma, hair; in allusion to the lack of crest. The word primarily means smooth, even in the sense of free from stones; α privative, and φελός or φελλός, a stone; φελλεύς, rocky soil, &c.
- 355. A. f. wood-hous'-ĭ-ī. To S. W. Woodhouse, M. D., of Philadelphia, who explored in New Mexico and Arizona.
- 356. A. f. căl-ĭ-fōr'-nĭ-că. To the State of California.
- 357. A. ūl-tră-măr-ī'-nă ă-rī-zō'-naē. Lat. ultra, beyond, from the adverb uls, beyond, opposed to cis, on this side; and marina, marine, relating to the sea, mare; in allusion to the deep blue color, as of the high sea; "ultramarine" blue. See Peucæa, No. 253.

  This stands in the orig. ed. as A. sordida, "Sieber's Jay."
- 358. Xān-thū'-ră lūx-ŭ-rǐ-ō'-să. Gr. ξανθός, yellow, and οὖρα, tail. Luxuosa was doubtless intended by Lesson for Lat. luxuriosa, luxurious, in allusion to the elegant coloration. This stands in the orig. ed. as X. yncas var. luxuosa, but proves to be distinct from the Peruvian yncas.
- 359. Pěr-ĭ-sō'-rě-ŭs căn-ă-dēn'-sĭs. Unde derivatur? One of the dictionaries gives a sorix, defined as a bird dedicated to Saturn; whence Perisoreus might be derived as an adjectival form, intensified by the preposition peri. This would accord in idea with the term infaustus bestowed by Linnæus on the European species, and also with Dysornithia, the generic term invented by Swainson; there being some superstition attaching to the jays of this genus. But we advance this etymology as mere conjecture. We may note also the Gr. σορόs, a tomb or sepulchre.
- 330. P. c. fū/-mī-frōns. Lat. fumus, smoke, and frons, forehead; related to Gr. 6ύω, I offer incense.

Described since the orig. ed.; Ridg., Pr. Nat. Mus., ii, 1880, p. 5.

- 361. Perisoreus canadensis obscurus Ridg. B —. C 239a. R 298. Oregon Jay.
- 362. Perisoreus canadensis capitalis Bd. B —. C 239b. R 297a. Rocky Mountain Jay.
- 363. Sturnus vulgaris L. B —. C —. R 279. (G. ! E.) European Starling.
- 364. Pitangus derbianus (Kaup) Scl. B —. C —. R 308. Lord Derby's Flycatcher.
- 365. Myiodynastes luteiventris Scl. B —, C —, R 310. Sulphur-bellied Flycatcher.
- 366. Milvulus tyrannus (L.) Bp. B 122. C 240. R 302. (! S. A.) Fork-tailed Flycatcher.
- 367. Milvulus forficatus (Gm.) Sw. B 123. C 241. R 301. Swallow-tailed Flycatcher; Scissor-tail.
- 368. Tyrannus carolinensis (L.) Bd. B 124. C 242. R 304. Tyrant Flycatcher; King-bird; Bee-martin.
- 369. Tyrannus dominicensis (Gm.) Rich. B 125. C 243. R 303. Gray Tyrant Flycatcher; Gray King-bird.
  - 361. P. c. ōb-scū'-rŭs. See Molothrus, No. 314.
  - 362. P. c. căp-ĭ-tā'-lĭs. Lat. capitalis, capital, relating to the head, caput, the color of which distinguishes the race from the stock species.
- 363. Stūr'-nus vūl-gā'-rīs. Lat. sturnus, a stare or starling.—Lat. vulgaris, vulgar, common; vulgus, or volgus, the people or folk, is digammated Gr. Fολχοs, with transposition of letters from δχλοs, a crowd.

Not in the orig. ed. Only American as occurring in Greenland, and there only accidentally, in one known instance.

364. Pǐt-ān'-gŭs dēr-bǐ-ā'-nŭs. *Pitangus* is a barbarous word, of some South American vernacular; it occurs, in several forms, in Marcgrave. — The species is dedicated to the Earl of Derby.

Not in the orig. ed. of the Check List. Since discovered in Texas by G. B. Sennett. See Coues, The Country, i, p. 184, July 13, 1878.

365. Mỹi-ŏ-dyn-ās'-tēs lūt-ĕi-vēn'-trīs. Gr. μυῖα, a fly, and δυναστής, a sovereign, ruler, &c.; δύναμις, power, from δύναμαι, I can, I am able.—Lat. lūteus, luteous, yellow, from lūtum, a plant used for yellow dye, and venter, genitive ventris, the belly; said to be digammated from Gr. ἔντερον, the entrails.

Not in the orig. ed. of the Check List: since discovered in Arizona by H. W. Henshaw. See Hensh., Rep. Expl. W. 100 Merid., v, 1875, p. 346, pl. xiv.

- 366. Mīl'-vŭ-lŭs tyr-ān'-nŭs. Lat. milvulus, diminutive of milvus, a kite. Lat. tyrannus, Gr. τύραννος, a ruler, despot, "tyrant;" well applied to a bird of this genus.
- 367. M. fōr-fi-cā'-tŭs. Lat. forficatus, a participial adjective, as if from a verb forfico; forfex, a pair of shears, scissors, which the deeply forked tail resembles.
- 368. Tyr-ān'-nŭs că-rŏ-lĭn-ēn'-sĭs. See Milvulus, No. 366.—Named after the State of Carolina: the direct adjective from Carolus, Charles. See Mimus, No. 16.
- 369. T. dŏm-ĭn-ĭ-cēn'-sĭs. Named after the island of Hayti, or St. Domingo; dominicus, dominus, domus. See Dendrαca, No. 129.

- 370. Tyrannus verticalis Say. B 126. C 244. R 306.
  Arkansas Tyrant Flycatcher.
- 371. Tyrannus vociferans Sw. B 127. C 245. R 307. Cassin's Tyrant Flycatcher.
- 372. Tyrannus melancholicus couchi (Bd.) Coues. B 128, 129. C 246. R 305. Couch's Tyrant Flycatcher.
- 373. Myiarchus crinitus (L.) Cab. B 130. C 247. R 312.

  Great Crested Flycatcher. [See Addenda, No. 880.
- 374. Myiarchus erythrocercus Scl. and Salv.? B 132? C —, R 311. (?)
  Rufous-tailed Crested Flycatcher.
- 375. Myiarchus cinerescens (Lawr.) Scl. B 131. C 248. R 313.
  Ash-throated Crested Flycatcher.
- 376. Myiarchus lawrencii (Gir.) Bd. B. 133. C 249. R 314. (! M.)
  Lawrence's Crested Flycatcher.
- 377. Sayiornis sayi (Bp.) Bd. B 136. C 250. R 316. Say's Pewit Flycatcher.
  - 370. T. vēr-tī-cā'-līs. Lat. verticalis, vertical, i. e., relating to the vertex, top or crown of the head, which has a flame-colored patch. The etymological meaning of vertex is vortex, the turning or whirling thing, from verto, I turn.
  - 371. T. vō-cī'-fĕr-āns. Lat. present participle vociferans, vociferating, vociferous, from vocifero; vox, genitive vocis, voice, and fero, I bear.
  - 372. T. měl-ān-chöl'-ĭ-cŭs. Gr. μελαγχολικόs, melancholy, from μέλαι, feminine μέλαινα, black, and χόλοs, gall, bile; Lat. melancholicus, atrabilious. The ancients had some notions on this subject which make the term not wholly inapplicable to a bird of splenetic, irritable disposition, as all of this genus are. To Lt. D. N. Couch, U. S. A., who collected extensively in Matamoras and Texas.
  - 373. Mỹĩ-ār'-chùs crī-nī'-tùs [not "crinnytus," as usually heard]. Gr. μωῖα, a fly, and ἀρχός, a ruler, leader, chief, from ἄρχω, I am first, lead, rule, or ἀρχή, the beginning. This theme is seen in our prefix arch-, as arch-bishop, &c. Lat. crinitus, haired, i. e., crested, from crinis, hair of the head. See Myiodioctes, No. 146.
  - 374. M. ĕ-rÿth-rŏ-cēr'-cŭs. Gr. ἐρυθρόs, reddish, and κέρκοs, tail.

    Not in the Check List, orig. ed. Since discovered in Texas by G. B. Sennett. The proper name of the species is much in question. The bird is the M. crinitus erythrocercus of Coues, Bull. U. S. Geol. Surv. Terr., iv, no. 2, p. 32, and v, no. 3, p. 402; the M. erythrocercus var. cooperi of Ridgway, Pr. Nat. Mus., i, p. 138; and the M. mexicanus of Ridg., Pr. Nat. Mus., ii, p. 14.
  - 375. M. cīn-ĕr-ēs/-cēns. Lat present participle of an inceptive verb cineresco, I grow ashy; in the sense of being somewhat ashy; cinereus, ashy, from cinis, ash. N. B.—The name has always been written cinerascens, for which we find no authority; while there is actually a verb cineresco: we therefore emend as above.
  - 376. M. law-ren'-ci-i. To George Newbold Lawrence, of New York.
  - 377. Sāy-ī-ōr'-nīs sāy'-ī. "Sayornis" is a violent combination of the name of Mr. Thomas Say, of Philadelphia, with the Greek word for bird, ὅρνις. It may be somewhat improved as above, when the combination of vowels becomes no more unusual than is seen in myio-dioctes, myia-rchus, &c. In equally loose style, Bonaparte made the specific name sayus,—a direct Latinization of the same person's name; but it must either be put in

- -378. Sayiornis nigricans (Sw.) Bp. B 134. C 251. R 317.
  Black Pewit Flycatcher.
- . 379. Sayiornis fusca (Gm.) Bd. B 135. C 252, R 315. Pewit Flycatcher; Phobe-bird.
  - 380. Contopus borealis (Sw.) Bd. B 137. C 253. R 318. Olive-sided Pewee Flycatcher.
  - 381. Contopus pertinax Cab. B —. C 254. R 319. Coues's Pewee Flycatcher.
- 382. Contopus virens (L.) Cab. B 139. C 255. R 320. Pewee Flycatcher; Wood Pewee.
- 383. Contopus virens richardsoni (Sw.) Coues. B 138. C 255a. R 321. Western Pewee Flycatcher.
- 384. Empidonax acadicus (Gm.) Bd. B 143. C 256. R 324. Acadian Flycatcher.
- 385. Empidonax trailli (Aud.) Bd. B 140. C 257. R 325a.
  Traill's Flycatcher.
- 386. Empidonax trailli pusillus (Bd.) Coues. B 141. C 257a. R 325.
  Little Western Flycatcher.
  - the genitive, sayi or saii, or in adjectival form, sayana or saiana; it must in the latter case be feminine to agree with sayiornis. The above emendation of both generic and specific names is respectfully submitted. (See Coues, Bull. Nutt. Club, v, 1880, p. 99.)
  - 378. S. nig'-ri-cans. Present participle of nigrico, I am blackish; niger, black.
  - 379. S. fūs'-că. Lat. fuscus, dark, dusky, swarthy. See Pipilo, No. 306.
- 380. Cōn'-tŏ-pūs bŏr-ĕ-ā'-līs. Gr. κόντος, in some sense unknown to us, and ποῦς, foot. Lat. borealis, northern; boreas, the northwind. "Pewee," like "pewit," is an onomatopæon.

  N. B. Many words ending in -opus, from the Gr. ποῦς and a connecting vowel o, are habitually accented on the lengthened penult, and the last syllable is made short. But as -pus here stands for Gr. ποῦς, and the connecting vowel is invariably short, we should throw the accent back to the antepenult, and dwell on the last syllable. Thus, not Contō'-pūs, Hæmatō'-pūs, Phalarō'-pūs, but Cont'ŏ-pūs, Hæma'tŏ-pūs, Phalar'ŏ-pūs.
- 381. C. pēr'-tī-nāx. Lat. pertinax, pertinacious, holding fast on to; from per and tenax, tenacious, from teneo, I hold; this species closely resembling C. borealis.
- 382. C. vĭr'-ēns. See Dendræca virens, No. 112.
- 383. C. v. rīch'-ārd-sŏn-ī. To Dr. John Richardson, an author of the Fauna Boreali-Americana, &c.
- 384. Em-pĭd-ō'-nāx ă-căd'-ī-cŭs. Gr. ἐμπίς, genitive ἐμπίδος, a small kind of insect, gnat; and ἀναξ or ἄναξ, king. Acadicus, Latinized adjective for Acadian; from Acadia or Acadie.

  N. B. This species has never been found, and probably does not occur, in the region formerly called Acadia; the name is therefore geographically false. The name "Acadian Flycatcher," whence Muscicapa acadica Gm., no doubt actually refers to Traill's or the Least Flycatcher, the proper name of the present species being probably Empidonax subviridis (Bartr.) Coues. Lat. subviridis, somewhat green, greenish.
- 385. E. traīl'-lī. To Thomas Stewart Traill, a Scottish naturalist. He was professor of medical jurisprudence in the University of Edinburgh, and editor of one of the later editions of the "Encyclopædia Britannica."
- 386. E. t. pŭ-sil/-lŭs. See Sitta, No. 60.

- 287. Empidonax minimus Bd. B 142. C 258. R 326. Least Flycatcher.
- -388. Empidonax flaviventris Bd. B 144. C 259. R 322. Yellow-bellied Flycatcher.
- 389. Empidonax flaviventris difficilis Bd. B 144a. C —. R 323. (?)
  Western Yellow-bellied Flycatcher.
- 390. Empidonax hammondi (Xant.) Bd. B 145. C 260. R 327. Hammond's Flycatcher.
- 391. Empidonax obscurus (Sw.) Bd. B 146. C 261. R 328. Wright's Flycatcher.
- 392. Mitrephorus fulvifrons pallescens Coues. B —. C 262. R 329a. Buff-breasted Flycatcher.
- 393. Ornithium imberbe Scl. B., C., R 331.

  Beardless Flycatcher.
- 394. Pyrocephalus rubineus mexicanus (Sel.) Coues. B 147. C 263. R 330. Mexican Vermilion Flycatcher.
- 395. Nyetidromus albicollis (Gm.) Burm. B —, C —, R 356. Pauraque.
  - 387. E. min'-i-mus. Lat. minimus, least, smallest, superlative degree of parvus, little.
- 388. E. flā-vǐ-vēn'-trīs. Lat. flavus, yellow; ventris, pertaining to the belly, venter, belly; probably digammated from Gr. ἔντερον, the entrails.
- 389. E. f. dif-fi'-ci-lis. Lat. difficilis, difficult, not facile; dis-facilis, not easily do-able; facio, I do; like agilis, active, or utilis, useful, from ago and utor. It is applied to the bird as the French would call a person difficile, that is, hard to get at, manage, understand, impracticable; the subspecies not being readily distinguished from E. flaviventris.
- 390. E. hăm'-mŏnd-ī. To Dr. W. A. Hammond, sometime Surgeon General, U. S. Army.
- 391. E. ōb-scū'-rŭs. See *Molothrus*, No. 314. To C. Wright, the discoverer. Swainson's bird is very uncertain, and our species might be called E. wrighti.
- 392. Mī-trē'-phŏr-ŭs fūl'-vī-frōns pāl-lēs'-cēns. Gr. μίτρα or μίτρη, a mitre or other headdress, and φορόs, bearing, from φέρω, same as Lat. fero, I bear. We believe either mitrephorus or mitrophorus to be admissible; the former has currency, though the latter may be preferable. Lat. fulvus, yellowish, fulvous, and frons, forehead. Lat. pallescens, somewhat pale, from pallesco, I grow pale; palleo, of same meaning. The allusion is to the pale coloration in comparison with the stock-form fulvifrons. [See Index, p. 137.]
- 393. Ōr-nīth'-I-ŭm īm-bēr'-bĕ. Gr. δρνίθιον, a little bird; diminutive of ὕρνις, a bird. Lat. imberbe, beardless, from in, negation, and barba, beard. The genus is commonly written Ornithion, but it is customary to change -ον of the Greek into -um in Latin.

  Not in the orig. ed.; since discovered in Texas by G. B. Sennett. See Coues, The Country, i, p. 184, July 13, 1878.
- 394. Py-rŏ-cĕph'-ă-lŭs rŭb-ĭn'-ĕ-ŭs mēx-ĭ-cã'-nŭs. Gr. πῦρ, genitive πυρός, fire, κεφαλή, head. Lat. rubineus (not classic), equivalent to rubens, ruby-colored, rose-red.
- 395. Nyc-ti'-drö-mus āl-bi-cōl'-līs. Gr. νύξ, genitive νυκτός, night, and δρόμος, act of running: in allusion to the nocturnal activity of the bird. See Ammodramus, No. 238. Lat. albus, white, and collis, neck.

Not in the orig. ed.; since discovered in Texas by G. B. Sennett, and J. C. Merrill.

- 396. Antrostomus carolinensis (Gm.) Gould. B 111. C 264. R 353. Chuck-will's-widow.
- 397. Antrostomus vociferus (Wils.) Bp. B 112. C 265. R 354.
  Whip-poor-will. [See Addenda, No. 881.
- 398. Phalænoptilus nuttalli (Aud.) Ridg. B 113. C 266. R 355. Nuttall's Whip-poor-will.
- /399. Chordediles popetue (V.) Bd. B 114. C 267. R 357. Night-hawk.
- 400. Chordediles popetue henryi (Cass.) All. B 115. C 267a. R 357a. Western Night-hawk.
- 401. Chordediles popetue minor (Cab.) Ridg. B —. C —. R 357b. (!W.I.) Cuban Night-hawk.
  - 402. Chordediles acutipennis texensis (Lawr.) Ridg. B 116. C 268. R 358. Texan Night-hawk.
    - 296. An-trō'-stŏ-mus că-rŏ-lın-ēn'-sıs. Gr. ἄντρον, Lat. antrum, a cave, στόμα, mouth; in allusion to the cavernous capacity of this fissirost.—The curious English name, like "whip-poor-will," is an onomatopœon, being an attempt to express the bird's cry in words.
  - 397. A. vō-cĭ'-fĕr-ŭs. Lat. vociferus, vociferous, clamorous, from vox, genitive, vocis, voice, and fero, I bear; vox is said to be digammated from Gr. ωψ.
  - 398. Phǎl-aē-nōp'-tīl-ŭs nǔt-tǎl-lī. Gr. φάλαινα, a moth, and πτίλον, plumage; in allusion to the peculiar velvety plumage, like the furriness of a moth's wing. To Thomas Nuttall.

    This is given as Antrostomus nuttalli in the orig. ed. The genus has since been established by Ridgway, Pr. Nat. Mus., ii, 1880, p. 5.
  - 399. Chōr-dē-dī'-lēs popetue. Gr. χορδή, a chord, a stringed instrument, and δείλη, contracted from δείελος, root εἴλω, the evening, here apparently meaning to close in, as evening does. The allusion is to the crepuscular habits of the bird, its curious notes being oftenest heard at evening. Swainson originally wrote chordeiles, an inadmissible contraction, and further erroneous in retaining Gr. ει instead of changing to long Lat. i. The word has sometimes been written chordiles. Cabanis properly emends as above. Swainson was very negligent in these matters: for instance, he made a genus aipunemia, the proper form of which is æpycnemis. The word popetue is barbarous, of meaning and pronunciation alike unknown to us. We have heard it as three and as four syllables, accented in each case on the antepenult.

This stands as Chordeiles virginianus in the orig. ed.

- 400. C. p. hen'-ry-i. To Dr. T. Charlton Henry, who collected and observed in the West.
- 401. C. p. min'-or. Lat. minor, minor, less, smaller, this form holding such relation to the stock species.

Not in the orig. ed. of the Check List. Stated to have occurred in Florida. See Ridg., Pr. Nat. Mus., iii, 1880, p. 219.

402. C. ă-cū-tǐ-pēn'-nīs tēx-ēn'-sīs. Lat. acutus, acute, sharp, pointed, and penna, wing or feather, in allusion to the long wings. — Texensis, adjective formed from Texas. Texas is properly a plural noun, singular Texa, meaning the Texas; as we should say now, the Texans, a race of the Caddos. Tachies and Taxus are also found.

This stands as C. texensis of the orig. ed.

- 403. Panyptila saxatilis (Woodh.) Coues. B 107. C 269. R 349. White-throated Rock Swift.
- .404. Nephœcetes niger borealis (Kenn.) Coues. B 108. C 270. R 350. Black Rock Swift.
- 405. Chætura pelasgica (L.) Steph. B 109. C 271. R 351. Chimney Swift.
- 406. Chætura vauxi (Towns.) De Kay. B 110. C 272. R 352. Vaux's Chimney Swift.
- 407. Basilinna xantusi (Lawr.) Elliot. B —. C 273. R 347. Xantus Humming-bird.
- 408. Eugenes fulgens (Sw.) Gld. B —. C 274bis. R 334. Refulgent Humming-bird.
- 409. Trochilus colubris L. B 101. C 275. R 335. Ruby-throated Humming-bird.
- Păn-yp/-ti-lă sāx-āt/-i-līs. Gr. πάνν, much, very, from πῶς, πῶσα, πῶν, all, and πτίλον, wing: in allusion to the great length of this member. Lat. saxatilis, rock-inhabiting: saxum, a rock.
  - 404. Něph-oē'-cĕ-tēs nīg'-ĕr bŏr-ĕ-ā'-līs. Gr. νέφος, a cloud, and οἰκέτης, an inhabitant; well applied to this bird of great wing and high flight. See Poacetes, No. 232. Lat. niger, black. Lat. borealis, northern.
  - 405. Chaë-tū'-ră pěl-ās'-gĭ-că. Gr.  $\chi\alpha(i\tau\eta)$ , a stiff hair, a bristle, and  $o\partial\rho\alpha$ , tail, in allusion to the spines which project from the ends of the tail-feathers.

The specific word was written pelagica by Linnæus in 1758, and pelasgia by him in 1766. The word has occasioned much conjecture as to its orthography, derivation, and applicability. We cannot suppose it to be pelagica, pelagic, relating to the high seas, like marine. It is apparently one of Linnæus's whims of nomenclature, by which he likened this migratory species to a Pelasgian, one of the nomadic tribes of Greece, the Pelasgi, Πελασγοί. There is indeed a geographical name pelasgia, but such would hardly be used in this form, and would be geographically false, moreover. Excluding pelasgia or pelagica as out of the question, and supposing the allusion to be to the nomadic Pelasgi, we conclude that the proper form of the word is as above given, pelasgica, the adjective meaning Pelasgian, i.e., in a tropical sense, nomadic, migratory.

- 406. C. vaūx'-ī. To William S. Vaux, of Philadelphia.
- 407. Băs-11-în'-nă xăn'-tŭs-î. Gr. βασίλιννα, a queen, feminine form of βασίλεύs, a king. To Louis John Xantus de Vesey, who later called himself John Xantus, an energetic and successful collector in South-western United States, and Mexico. We suppose the name originally meant yellow, ξανθός, xanthus, and in fact it is written xanthusi sometimes.

  This is given as Heliopædica xantusi in the orig. ed. of the Check List.

408. Eū'-gĕn-ēs fūl'-gēns. Gr. εὐγενήs, well-born; from εἶ, well, and γένοs, birth; γίγνομαι, I am born. — Lat. fulgens, glittering, refulgent, from fulgeo, I shine, flash, gleam, glitter. Not in the orig. ed.: since discovered in Arizona by H. W. Henshaw.

409. Trŏch'-ĭ-lŭs cŏl'-ŭ-brĭs. Gr. τρόχιλος or τροχίλος, Lat. trochilus, a kind of bird; from τροχός, a runner. The bird originally so called by Herodotus was an Egyptian species of plover, of the genus Ægialitis, which was so named from its habit of coursing the banks of streams. The name was also applied by the ancients to some small bird, species uncertain, perhaps a warbler, wren, or kinglet. Very curiously, the name was afterward transferred to the American humming-birds, becoming fixed in modern nomen-

- 410. Trochilus alexandri Boure. and Muls. B 102. C 276. R 336. Alexander Humming-bird.
- 411. Selasphorus rufus (Gm.) Sw. B 103. C 277. R 340. Rufous Humming-bird.
- 412. Selasphorus alleni Hensh. B —. C —. R 341.
  Allen Humming-bird.
- 413. Selasphorus platycercus (Sw.) Gld. B 104. C 278. R 339. Broad-tailed Humming-bird.
- 414. Calypte annæ (Less.) Gld. B 105. C 279. R 338.
  Anna Humming-bird.
- 415. Calypte costæ (Bourc.) Gld. B 106. C 280. R 337. Costa Humming-bird.
- 416. Atthis heloisæ (Less. and De Lat.) Reich. B —. C 281. R 342. Heloise Humming-bird.
- 2417. Stellula calliope Gld. B —. C 282. R 343.
  Calliope Humming-bird.
  - 418. Calothorax lucifer (Sw.) Gray. B —. C —. R 349. Lucifer Humming-bird.

clature as a genus in that family in consequence of such usage on the part of Linnæus. — The name colubris might be an adjective formed from coluber, a snake, in allusion to the scales on the hummer's throat; but this is unlikely. There are old treatises on birds in which the terms colibri, kolibri, colibry occur, and the word is doubtless barbarous.

- 410. T. ă-lĕx-ān/-drī. To Alexandre.
- 411. Sĕ-lās'-phŏr-ŭs rū'-fūs. Gr. σέλας, σέλας, σίλας, depops, bearing, φέρω, I bear; euphoniously compounded, at the expense of strict propriety. Lat. rufus, rufous, reddish.
- 412. S. ăl'-lĕn-ī. To C. A. Allen, of Nicasio, California. Not in the orig. ed.; since distinguished from S. rufus by Mr. Henshaw: see Bull. Nutt. Club, ii, 1877, p. 54.
- 413. S. plat-y-cer'-cus. Gr. πλατύς, broad, wide; κέρκος, tail.
- 414. C. ān'-naē. Dedicated to the Duchess of Rivoli.

  This is Selasphorus anna in the orig. ed.
- 415. Că-lỹp'-tē cōs'-taē. Gr. Καλυπτή, a proper name; καλύπτω, I conceal. Το —— Costa.

  This is Selasphorus costa in the orig. ed.
- 416. At'-thĭs hĕl-ŏ-ī'-saē. Gr. 'Aτθίs, Attic, Athenian; probably in allusion to some peculiar charm of the bird. Attic was ne plus ultra Greek, as Parisian is par excellence French. This is Selasphorus heloisæ of the orig. ed.
- 417. Stēl'-lǔ-lǔ cāl-lī'-ŏ-pē. Lat. stellula, a little star, diminutive of stella, a star. Gr. Καλλιόπη, Calliope, one of the Muses; καλός, feminine καλλή, beautiful, &c., and όψ, voice. The application of the word to a voiceless bird is not obvious, unless it be simply dedicatory.
- 418. Căl-ŏ-thō'-rāx lū'-cĭ-fĕr. Gr. καλός, beautiful, and θώραξ, thorax, chest. Lat. Lucifer, Lucifer, the light-bearer, from lux, lucis, light, and fero, I bear. Both words note the glittering plumage.

Not in the orig. ed.; since discovered in Arizona by H. W. Henshaw; and first announced from that locality under the erroneous name of "Doricha enicura," in Am. Sportsm., v, p. 328, Feb. 20, 1875. See Lawr., Bull. Nutt. Club, ii, 1877, p. 108.

- 419. Amazilia fuscocaudata (Fras.) Elliot. B. —. C —. R 345. Dusky-tailed Humming-bird.
- 420. Amazilia cerviniventris Gld. B —. C —. R 346. Buff-bellied Humming-bird.
- 421. Iache latirostris (Sw.) Elliot. B —, C —, R 348. Circe Humming-bird.
- 422. Trogon ambiguus Gld. B 65. C 284. R 384. (!M)
  Copper-tailed Trogon.
- /423. Ceryle alcyon (L.) Boie. B 117. C 286. R 382. Belted Kingfisher.
  - 424. Ceryle americana cabanisi (Reich.) Coues. B 118. C 287. R 383. Texas Kingfisher.
  - 425. Crotophaga ani L. B 66, 67. C 288. R 389. Black Ani.
  - 419. Am-ā-zīl'-ĭ-ā fūs-cŏ-caūd-ā'-tă. The word amazilia is apparently Latinized from Lesson's word amazili, used in the plural form amazilis for a group of hummers. We do not know what it means. Lat. fuscus, dark, and caudata, tailed; cauda, tail.

    Not in the orig. ed.; since discovered in Texas by J. C. Merrill. This has been

Not in the orig. ed.; since discovered in Texas by J. C. Merrill. This has been called *Pyrrhophæna riefferi* in papers relating to the Texas specimens. See Merrill, Bull. Nutt. Club, i, 1876, p. 88, and Ridg., Pr. Nat. Mus., i, 1878, p. 147.

- 420. A. cer-vī-nĭ-vēn'-trĭs. Lat. cervinus, relating to a deer, cervus; and ventris, pertaining to the belly, venter. The allusion is to the fuwn-colored under parts.

  Not in the orig. ed.; since discovered in Texas by J. C. Merrill. See Bull. Nutt. Club, ii, 1877, p. 26, and Pr. Nat. Mus., i, 1878, p. 148.
- 421. I'-ă-chē lā-tǐ-rōs'-trĭs. Gr. laχή, a battle-cry; also a proper name, whence derived. Lat. latirostris, broad-billed; latus, wide, like Gr. πλατύs, of same meaning; and rostrum, beak.

  Not in the orig. ed. of the Check List; since discovered in Arizona by H. W. Henshaw. See Amer. Sportsm., Feb. 20, 1875.
- 422. Trō'-gōn ām-bi'-gŭ-ŭs. Gr. τρώγων, a gnawer, rodent, from τρώγω, I gnaw, eat away, corrode; from the stout, dentate bill; see Troglodytes, No. 74. The word was applied by Moehring in 1752 to the Brazilian Trogon, called curucui by the natives, and made generic by Brisson in 1760. Lat. ambiguus, ambiguous, equivocal, of more than one meaning, in a double sense; hence, doubtful, uncertain; from ambo, both, on two sides, and ago, to act or do. Ambiguity is literally a double-dealing, "with double sense deluding;" compare Fr. double entendre, and such homely expressions as "back and fill," "blow hot and cold," "on the fence," "hedge" (to bet on both sides). It was badly applied to this fine species when considered doubtfully distinct from T. mexicanus.

This stands as *T. mexicanus* in the orig. ed. For its actual occurrence in Texas, see Pr. Nat. Mus., i, 1878, p. 118.

- 423. Cē'-ry-lē āl'-cy-ōn. Gr. κήρυλος, a kingfisher.— Gr. ἀλκυών, Lat. haleyon or aleyon, a kingfisher. 'Αλκυώνη or Aleyone was a mythical character, daughter of Æolus, fabled to have been transformed into a kingfisher when, out of love for her shipwrecked husband Ceyx, she threw herself into the sea. The kingfisher was also believed to nest on the water, at a time the waves were stilled; hence the term "haleyon days."
- 424. C. ăm-ĕr-ĭ-cā/-nă căb-ăn/-ĭs-ī. To Dr. Jean Cabanis, long time one of the leaders of German ornithology, and editor of the Journal für Ornithologie.
- 425. Crŏ-tō'-phă-gă ā'-nī. Gr. κροτών, a bug, tick, plant-louse; and φάγοs, from φάγομαι, I eat.

- 426. Crotophaga sulcirostris Sw. B —, C —, R 390. Groove-billed Ani.
- 427. Geococcyx californianus (Less.) Bd. B 68. C 289. R 385. Ground Cuckoo; Chaparral Cock; Road-runner.
- 428. Coccygus erythrophthalmus (Wils.) Bd. B 70. C 290. R 388. Black-billed Cuckoo.
  - 429. Coccygus americanus (L.) Bp. B 69. C 291. R 387. Yellow-billed Cuckoo.
  - 430. Coccygus seniculus (Lath.) V. B 71. C 292. R 386. Mangrove Cuckoo.
  - 431. Campephilus principalis (L.) Gr. B 72. C 293. R 359. Iyory-billed Woodpecker.
  - 432. Hylotomus pileatus (L.) Bd. B 90. C 294. R 371. Pileated Woodpecker.
  - .433. Picus borealis V. B 80. C 296. R 362. Red-cockaded Woodpecker.
    - 426. C. sūl-cĭ-rōs'-trĭs. Lat. sulcus, a groove, furrow, channel; a word sibilated from Gr. δλκόs, a trace, track, trail; and rostris, pertaining to the beak, rostrum.

      Not in the orig. ed.; since discovered in Texas by G. B. Sennett. See Coues, The Country, i, July 13, 1878, p. 184.
    - 427. Gĕ-ŏ-cōc'-cȳx căl-ĭ-fŏr-nĭ-ān'-ŭs. Gr. γη̂ or γέα, the earth, and κόκκυξ, a cuckoo. The latter word is onomatopœic, and runs in similar forms through many languages, the idea being always to express the cuckoo's voice in a word: Lat. cuculus; Fr. coucou; Eng. cuckoo, cuckow; Germ. Rufuf, &c. See Coccygus, No. 428.
    - 428. Cōc'-cy-gŭs č-ryth-rōph-thāl'-mŭs. The generic name is modified from κόκκυξ, a cuckoo. Its orthography has given rise to much variance of opinion. It was originally written by Vieillot coccyzus; such spelling has been accepted by Sclater and others, and is perhaps defensible on the ground that there is a Greek verb κοκκύζω, I make a noise like a cuckoo, whence a noun κοκκύζος, becoming coccyzus in Latin, might be formed. Boie first emended Vieillot's name to coccygus, in which he was followed by Cabanis and many others. Other forms of the word found in ornithological writings are: coccyzon, coccygius, coccysus, coccyzius, coccygon. We adopt Boie's form coccygus, being directly from the genitive of κόκκυξ, not wishing to unnecessarily interfere. For erythrophthalmus, see Pipilo, No. 301.
    - 429. C. ăm-ĕr-ĭ-cā/-nŭs. To America. See Parula, No. 93.
    - 430. C. sen-I'-cu-lus. Lat. seniculus, a little old man; diminutive of senex, an old man. The allusion is probably to the gray on the head, a sign of senility.
    - 431. Cām-pē'-phīl-ŭs prīn-cǐ-pā'-līs. Gr. κάμπη, a caterpillar, from its bending; well-illustrated in the way a "measuring-worm" bends. The word primarily means a bending: καμπτόs, bent; κάμπτω, I bend; the same word is seen in Campylorhynchus, for example. φίλος, φιλέω, I love. Lat. principalis, principal, chief, from the great size of the bird.
    - 432. Hy-lŏ'-tŏ-mŭs pī-lĕ-ā'-tŭs. Gr. ὁλοτόμος, cutting wood, i.e., a woodcutter: ὕλη, wood, and τέμνειν, to cut.—Lat. pileatus, capped, i.e., crested; from pileus or pileum, a cap; related to pilus, a hair; the same root is seen in depilatory, pile, as of velvet, &c.
    - 433. Pī'-cŭs bŏr-ĕ-ā'-lïs. Lat. Picus, a mythical person, and also a woodpecker, because the former, one of the victims of Circe, whose love he had scorned, was transformed into a woodpecker. The etymology of picus is doubtful; the word is said by some to be prob-

- 434. Picus scalaris Wagl. B 79. C 297. R 363. Texas Woodpecker.
- 435. Picus scalaris nuttalli (Gamb.) Coues. B 78. C 297a. R 364. Nuttall's Woodpecker.
- 436. Picus scalaris lucasanus (Xant.) Coues. B —. C 297b. R 363a. St. Lucas Woodpecker.
- 437. Picus stricklandi Malh. B —. C —. R 365. Strickland's Woodpecker.
- 438. Picus villosus L. B 74. C 298. R 360, 360a. Hairy Woodpecker.
- 439. Pieus villosus harrisi (Aud.) All. B 75. C 298a, R 360b. Harris's Woodpecker.
- 440. Picus pubescens L. B 76. C 299. R 361. Downy Woodpecker.
- 441. Picus pubescens gairdneri (Aud.) Coues. B 77. C 299a. R 361a. Gairdner's Woodpecker.

ably for pigus, from pingo, I paint, and hence to mean pigus or picus, painted, spotted; if so, it is well applied to the woodpecker, a bird of variegated colors, a much pied bird: compare Pica, No. 347. Others hold, however, that picus is from the same root as the Gr.  $\pi i \pi \omega$  or  $\pi i \pi os$ , a little bird, a peeper, chirper; just as Gr.  $i \pi \pi os$  or  $i \kappa \kappa os$  and Lat. equus (which was formerly spelled very differently, and with c instead of q) are cognate. This would make it an onomatopoon, like pipit, pipilo, &c. — Lat. borealis, northern; boreas, the north-wind.

Note. — According to Professor Newton (Ibis, 3d ser., vi, 1876, p. 94 seq.), the type of the Linnæan genus Picus is P. martius. The same author adds, in a private note addressed to Dr. Coues, that "the adjective in any other combination loses its classical allusion, which all naturalists, including Linnæus, until comparatively recent times, recognized." It would also appear that our H. pileatus, No. 432, is congeneric with P. martius. On these premises, No. 432 should stand as Picus pileatus, and some other generic name be found for Nos. 433–441. It is regretted, that, as the untoward circumstances (tent-life in unbookish Arizona) under which these proof-sheets are being corrected do not permit us to follow up the matter at present, we are obliged to let the current nomenclature pass with this explanation.

- 434. P. scā-lā'-rīs. Lat. scalaris, ladder-like; scala, a flight of stairs, a ladder, scale, shortened from scandla, from scando, I climb. The idea in Wagler's mind may have been the climbing or scaling of trees by the bird; more likely the bars on the back, resembling the rounds of a ladder.
- 435. P. s. nut'-tal-li. To Thomas Nuttall. Perhaps entirely distinct from No. 434.
- 436. P. s. lū-căs-ā'-nŭs. To Cape St. Lucas, S. Cala., where discovered.
- 437. P. strick'-lănd-i. To Hugh E. Strickland, the eminent English ornithologist. Not in the orig. ed. Since discovered in Arizona by H. W. Henshaw. See Amer. Sportsm., v, p. 328, Feb. 20, 1875.
- 438. P. vīl-lō'-sus. Lat. villosus, shaggy, hairy, villous; from villus, a hair, tuft of hair.
- 439. P. v. hăr'-rīs-ī. To Edward Harris, companion and friend of Audubon.
- 440. P. pū-bē'-scēns. Lat. pubescens, present participle of pubesco, I come to puberty, i. e., the time when the hair grows on the genitals; pubes, the parts on which such hair grows; hence, pubescent, hairy, downy.
- 441. P. p. gaīrd'-nĕr-ī. To Dr. Gairdner, a Scottish naturalist.

- 442. Xenopicus albolarvatus (Cass.) Bd. B 81. C 295. R 366. White-headed Woodpecker.
- 443. Picoïdes arcticus (Sw.) Gray. B 82. C 300. R 367. Black-backed Three-toed Woodpecker.
- 444. Picoïdes americanus Brehm. B 83. C 301. R 368. Banded-backed Three-toed Woodpecker.
- 445. Picoïdes americanus dorsalis (Bd.) Allen. B 84. C 301a. R 368a. Striped-backed Three-toed Woodpecker.
- 446. Sphyropicus varius (L.) Bd. B 85. C 302. R 369. Yellow-bellied Woodpecker.
- 447. Sphyropicus varius nuchalis Bd. B 86. C 302a. R 369a. Nuchal Woodpecker.
- 448. Sphyropicus varius ruber (Gm.) Ridg. B 87. C 302b, or 303. R 369b. Red-breasted Woodpecker.
- 449. Sphyropicus thyroïdes (Cass.) Bd. B 88, 89. C 304, 305. R 370. Brown-headed Woodpecker.
- 442. Xĕn-ŏ-pī'-cŭs āl-bŏ-lār-vā'-tŭs. Gr. ξένος, a guest, stranger; ξένος, rare, foreign, &c. Lat. albolarvatus, white-masked; albus, white, and larca, a mask. The same word is used for insects in their early stage, when the characters of the imago, or perfect insect, are masked or hidden in the caterpillar.

Given as *Picus a.* in the orig. ed. For generic characters, see Ridgw., Pr. Nat. Mus., ii, 1880, p. 6.

- 443. Pi-cŏ-ī'-dēs ārc'-tĭ-cŭs. Lat. picus, a woodpecker, and Gr. εἶδοs, resemblance. The word is one of the numerous bastards in the genera of Picidæ, which authors seem bent on producing; there is no such word as Picus in Greek, yet they have constantly compounded it with Greek adjectives. The εἶ becomes long i; the o is the connecting vowel; the word should have the diæresis over the i, and be pronounced in four syllables, with accent on the penult. All such hybrid words are so far wrong as to be past praying for, and scarcely worth the trouble of trying to twist into some decent shape.
- 444. P. ăm-ĕr-ĭ-cā'-nŭs. To America. See Parula, No. 93.
- 445. P. a. dor-sa'-lis. Lat. dorsalis, pertaining to dorsum, the back.
- 446. Sphy-rŏ-pī'-cŭs văr'-ĭ-ŭs. Gr. σφῦρον, a hammer, and Lat. picus. It was originally written sphyrapicus by Baird; but the connecting vowel should be o in this case. It is usually accented on the antepenult, with shortening of the i in picus, for which we see no reason, beyond our extreme tendency to throw the accent always backward. The word is a hopeless hybrid, even when emended as above; sphyrocopus (σφυροκόπος) would have been classic for a hammerer. Lat. varius, various, varied, variegated; referring to the coloration in this case.
- 447. S. v. nū-chā-lis. Quasi-Lat. nuchalis, relating to the nape, nucha, which is red in this bird, not in S. varius. See Leucosticte, No. 205.
- 448. S. v. r\u00e4b'-\u00e4r. Lat. ruber, red.

  This stands as S. ruber in the body of the orig. ed. of the Check List: as above in the appendix.
- 449. S. thỹ-rō-ī/-dēs. Gr. θυρεοειδήs, resembling a certain kind of shield; in allusion to the shield-shaped black spot on the breast; θυρεόs, a shield, είδοs, resemblance. The fuller form of the word would be thyreoïdes, in five syllables. It has always been wrongly written thyroideus. See especially Picoïdes, No. 443.

Note. — S. williamsoni, No. 305 of the orig. ed., is the male of the same species.

- 450. Centurus carolinus (L.) Bp. B 91. C 306. R 372. Red-bellied Woodpecker.
- .451. Centurus aurifrons Wagl. B 92. C 307. R 373. Yellow-fronted Woodpecker.
- 452. Centurus uropygialis Bd. B 93. C 308. R 374. Gila Woodpecker.
- 453. Melanerpes erythrocephalus (L.) Sw. B 94. C 309. R 375. Red-headed Woodpecker.
- 454. Melanerpes formicivorus bairdi Ridg. B 95. C 310. R 377. Californian Woodpecker.
- 455. Melanerpes formicivorus angustifrons Bd. B C 310a. R 377a. Narrow-fronted Woodpecker.
- -456. Asyndesmus torquatus (Wils.) Coues. B 96. C 311. R 376. Lewis's Woodpecker.
- 457. Colaptes auratus (L.) Sw. B 97. C 312. R 378. Golden-winged Woodpecker; Flicker.
  - 458. Colaptes chrysoïdes Malh. B 99. C 313. R 379. Gilded Woodpecker.
  - 450. Cēn-tū'-rūs că-rŏ-lī'-nūs. Gr. κέντρον, a point, prickle, and οδρα, tail; spine-tailed. The full form would appear to be Centrurus (like Centrocercus, for example), but there is a way of getting Centurus from κέντη-; κεντέω is the verb to prick, goad, &c. Carolinus is badly syncopated from carolinianus; carolinensis would have been better still.
  - 451. C. aur'-ĭ-frons. Lat. aurifrons, golden-forehead; aurum, gold (yellow), and frons, forehead.
  - 452. C. ū-rŏ-pȳ-gĭ-ā'-līs. There is a very late Latin word uropygium, the rump, from which the above is derived as an adjective. But this is merely a modern Latinizing of the good Gr. οὐροπύγιον or οἰροπύγιον, the rump; from οὖρα, tail, and πυγή, the buttocks. The allusion in this case is to the conspicuously white rump of the bird, which a Greek would have called πύγαργος (pygargus).
  - 453. Měl-ăn-ēr'-pēs ĕ-rÿth-rŏ-cĕph'-ā-lŭs. Gr. μέλαs, genitive μέλανοs, black, and ἔρπηs, a creeper; ἔρπω, I creep, crawl. See Catherpes, No. 66. The full form would be melanoherpes. Gr. ἐρυθρόs, red, and κεφαλή, head.
  - 454. M. fōr-mī-cī'-vŏr-ŭs baīrd'-ī. Lat. formica, an ant, and voro, I devour, in allusion to a habit of the species.—To Prof. Spencer Fullerton Baird. See Ridg., Bull. U. S. Nat. Mus., No. 21, 1881, p. 85. Given in the orig. ed. as M. formicivorus.
  - 455. M. f. ān-gūs'-tĭ-frōns. Lat. angustus, narrow, straitened, from ango, I press upon, draw together, &c.; Gr. ἄγχω, I squeeze, strangle, distress, &c.; the same root and idea is seen in anxious, anxiety, &c.; frons, forehead. The allusion is to the narrowness of the yellow frontal band.
  - 456. Ă-syn-dēs/-mus tor-quā/-tus. Gr. à privative, σόν, together, with, δεσμός, a bond; in allusion to the loosened texture of the feathers of the under parts. Lat. torquatus, collared; torquis, a necklace, collar; torqueo, I twist, twine around; tortus, twisted, distorted, contortion; so also torture, as of one wrenched or racked. The allusion is to the ashy collar on the neck of the bird. The English name is that of Merriwether Lewis, the explorer in company with Clark (Clark's Crow, Picicorvus).
  - 457. Cŏl-āp'-tēs aūr-ā'-tŭs. Gr. κολαπτής, a chisel, hammer; κολάπτω, I use such an instrument; very appropriate to a woodpecker.—Lat. auratus, gilded, golden (colored); aurum, gold; also very apt to this bird.
  - **458.** C. chry-sŏ-ī'-dĕs. Gr. χρύσεος, χρυσοῦς, golden, of the color of gold, χρυσός; εἶδος, resemblance.

- 459. Colaptes mexicanus Sw. B 98. C 314. R 378a.
  Red-shafted Woodpecker.
- 460. Conurus carolinensis (L.) Kuhl. B 63. C 315. R 392. Carolina Parrot; Paroquet.
- 461. Aluco flammeus pratincola (Bp.) Coues. B 47. C 316. R 394. American Barn Owl.
- 462. Bubo virginianus (Gm.) Bp. B 48. C 317. R 405. Great Horned Owl.
- 463. Bubo virginianus arcticus (Sw.) Cass. B —. C 317a. R 405a, 405b. Arctic Horned Owl.
- 464. Bubo virginianus pacificus Cass. B —, C 317b, R 405c.
  Pacific Horned Owl.
- 459. C. mēx-ĭ-cā/-nŭs. To Mexico.
- 460. Cō-nū-rŭs că-rŏ-līn-ēn'-sĭs. Gr. κῶνος, Lat. conus, a cone, pine-cone, whence our word for a figure of that kind; οδρα, tail; in allusion to the wedged or cuneate tail.

Note. — The nomenclature of our owls, Nos. 461–488, must be considered still unsettled in several instances, though we have endeavored to approximate toward a fixed terminology in this difficult group, where the species and subspecies are not readily determined, and where authors have bandied about the generic and specific names so indiscriminately as to produce great confusion. The names here provisionally adopted are in the main according to results reached by Mr. Ridgway, who has given special attention to these birds.

461. Ăl-ū'-cō flām'-mě-ŭs prāt-īn'-cŏ-lã. The meaning of Aluco we do not know, further than that it has long been used for some kind of owl; perhaps related to ἐλεόs, which occurs in Aristotle as the name of some owl, and is enumerated by Brisson among the synonyms of the European barn owl. Numberless names of owls in very many languages are doubtless more nearly related than their diverse orthography would show at first sight, and mostly appear to be onomatopæic, in imitation of the hooting, howling cries of these inauspicious birds of the night: Eng. owl, owlet, howlet; A. S. ul, eul, ule; Dutch, uil; Dan. ugle; Sw. uggla; Germ. cule; Fr. hulotte; Ital. alocho (compare aluco); Sansk. uluka, &c. — Lat. flammeus, flaming, fiery-red; flamma (flag-ma), a flame, blaze; the root is seen in flagrant, flagitious, deflagrate; flagro, I flare up, am inflamed; and many kindred words. The allusion, rather strong, is to the flagrant colors of this species in comparison with most owls. —Lat. pratincola, an inhabitant of fields; pratum, a meadow, incola, an inhabitant (in and colo, I cultivate).

This stands as *Strix flammea americana* in the orig. ed., and Ridgway has *A. flammea americana*; but *pratincola* Bp. (1838) antedates *americana* Aud. (1839); and, on the generic nomenclature of owls, especially on the type of *Strix* L., see Newton, Yarr. Br. B., 4th ed., i, p. 150, and Ibis, 3d ser., vi, 1876, p. 94.

- 462. Bū'-bō vĭr-gĭn-ī-ā'-nŭs. Lat. bubo, the horned owl; perhaps related to bubulus or bubalus; bos, Gr. βοῦς, a bull, horned cattle; there is a similar Greek word βύας, for a horned owl. So, also, the verb bubo or bubalo, to low, hoot; the word for the bittern, butor, botaurus (bos, taurus), and others, are related, all being onomatopæic, with reference to the lowing or bellowing of cattle. Virginianus, see Cardinalis, No. 299.
- 463. B. v. ārc/-tĭ-cŭs. See Sialia, No. 29.
- 464. B. v. pā-cĭ'-fī-cŭs. Lat. pacificus, pacific, peaceable, peace-making; pax, peace, facio, I do, make; "the stilly sea." The reference is to the habitat of the bird.

We retain the three forms of Bubo as given in the orig. ed. Mr. Ridgway, after dismissing Mr. Cassin's var. pacificus, has four: B. v., and B. v. arcticus, as we have them;

465. Scops asio (L.) Bp. B 49. C 318. B 402. Screech Owl; Mottled Owl; Red Owl.

- 466. Scops asio kennicotti (Elliot) Coues. B —. C 318a. R 402d. Kennicott's Screech Owl.
- 467. Scops asio maxwellæ Ridg. B —. C —. R 402c. Rocky Mountain Screech Owl.
- 468. Scops asio maccalli (Cass.) Coues. B 50. C 318b. R 402b.

  McCall's Screech Owl.
- · 469. Scops asio floridanus Ridg. B —. C 318c. R 402a. Florida Screech Owl.
- 470. Scops trichopsis Wagl. B —. C —. R 403. (?)
  Mexican Screech Owl.
- 471. Scops flammeolus (Licht.) Scl. B —. C 319. R 404. Flammulated Screech Owl.
- 472. Asio wilsonianus (Less.) Coues. B 51. C 320. R 395 Long-eared Owl.

with B. v. subarcticus, after Hoy, and B. v. saturatus, Ridg., from the North-west coast, the latter being var. pacificus of Hist. N. A. B., iii, p. 65.

- 465. Scops ăs'-ĭ-ō. Lat. scopes or scops, Gr. σκώψ, a kind of owl. Here we have a name for owl which regards the bird in an entirely different sense from that implied in any of the onomatopæic names. The etymology is disputed. Some say from σκώπτω, I mock, scoff, deride, which would make scops the same as σκώπτης, a mocker, mimic; the actions of an owl seeming to travesty the beholder. Others have it from σκοπέω, I look out, survey, contemplate, the root of this being seen in scope, telescope, &c.; or from σκέπτομαι, I examine, scrutinize, am sceptical about any thing; the reference being to the great staring eyes of the bird, or its air of contemplation. Lat. asio, a horned owl; occurring in Pliny; apparently a word of Hebrew extraction, the significance of which is unknown to us.
  - 466. S. a. kĕn-nĭ-cŏt'-tī. To Robert Kennicott, of Illinois, an ardent and able naturalist, who sadly lost his life on the Yukon River, in Alaska, where the variety was procured.
  - 467. S. a. măx'-wĕl-laē. To Mrs. M. A. Maxwell, of Boulder, Colorado, the discoverer. Not in the orig. ed. of the Check List; since described. See Field and Forest, June, 1877, pp. 210, 213.
  - 468. S. a. măc-căl'-lī. To Colonel G. A. McCall, U. S. A., of Philadelphia, who studied ornithology in Texas.

    The S. a. enano, recently attributed to Texas by Coues and Sennett, has been identified with this by Ridgway.
  - 469. S. a. flo-ri-da'-nus. To Florida.
  - 470. S. trīch-ōp'-sīs. Gr. θρίξ, genitive τριχόs, hair, and ἄψ, aspect, countenance; i. q., hairy-faced, bristly about the bill? or general plumage of that character?

    Not in the orig. ed. of the Check List. If not the species itself, then its identification with any United States specimens, would appear to be dubious. The name is inserted upon Mr. Ridgway's authority.
  - 471. S. flām-mě'-ŏ-lŭs. Lat. flammeolus, diminutive of flammeus: see Aluco, No. 461.
  - 472. As'-ĭ-ō wĭl-sŏn-ĭ-ā'-nŭs. For asio, see Scops, No. 465.—Latinized Wilsonian; to Alexander Wilson, "father of American ornithology."

This stands as Otus vulgaris var. wilsonianus in the orig. ed., but is now regarded as

- 473. Asio accipitrinus (Pall.) Newt. B 52. C 321. R 395. Short-eared Owl.
- 474. Strix cinerea Gm. B 53. C 322. R 399. Great Gray Owl.
- 475. Strix cinerea lapponica (Retz.) Coues. B —. C —. R 399a. (! A.)
  Lapland Great Gray Owl.
- 476. Strix nebulosa Forst. B 54. C 323. R 397. Barred Owl.
- 477. Strix nebulosa alleni Ridg. B—. C—. R 397a. Florida Barred Owl.
- 478. Strix occidentalis (Xant.) Ridg. B —. C 324. R 398. Western Barred Owl.
- 479. Nyctea scandiaca (L.) Newt. B 61. C 325. R 406. Snowy Owl.

sufficiently distinct from the European bird. — The genus Otus is from the Lat. otus, Gr.  $\delta \tau \sigma s$  or  $\delta \tau \delta s$ , the eared owl; Gr.  $o\delta s$  or  $\delta s$ , genitive  $\delta \tau \delta s$ , an ear; from  $o\delta \alpha s$ , a handle. (See Bubo, No. 462, and compare  $\beta \delta \alpha s$  and  $\beta o\delta s$ .) — The genus Asio would appear to be eligible for the group of long-eared owls commonly called Otus of late years. — It is quite likely that the most available specific name for our bird is americanus (Steph.), as Ridgway has it.

- 473. A. āc-cĭp-ĭt-rī'-nŭs. Lat. accipitrinus, accipitrine, hawk-like; see Accipiter, No. 494.

  This stands as Brachyotus palustris in the orig. ed. But both the eared owls may well be put in one genus, and the name accipitrinus has priority over brachyotus. This last word is literal Greek for "short-eared."
- 474. Strīx cĭn-ĕr'-ĕ-ă. Lat. strix, stryx, or strynx, or Gr. στρίγξ, a screech-owl; from strido, I screech, utter shrill strident sounds of any kind; Gr. στρίζω; sibilated from τρίζω. The same root is seen in the English strident, stridulous.—Lat. cinereus, ashy; cinis, ashes. See Harporhynchus, No. 22.

This stands as Syrnium lapponicum var. cinereum in the orig. ed., by a blunder; for the latter name has priority over the former. The late rectifications made by Newton in the genera of owls cause Strix to be referred to the common Brown Owl of Europe, strictly congeneric with our Barred Owl. If the great Gray Owls be considered generically distinct, they may be called Scotiaptex. Mr. Ridgway uses the genus Ulula for this group, which he separates from Strix proper.

475. S. c. lāp-pŏn'-ĭ-că. To Lapland.

This European conspecies of the great Gray Owl has lately been attributed to North America by Ridgway: see Bull. Nutt. Club, iii, 1878, p. 37; Alaska. Not in orig. ed.

- 476. S. něb-ŭ-lō'-să. Lat. nebulosus, nebulous, misty, foggy, in the sense here of dark clouded color; from the Gr. νεφέλη (νέφος), a cloud. So, also, Lat. nubes, a cloud; nubo, I marry, nubilis, marriageable; the bride being veiled (nupta) for the nuptials.

  This is Syrnium nebulosum of the orig. ed.
- 477. S. n. ăl'-lĕn-ī. To J. A. Allen, of Cambridge, Mass. See *Pipilo*, No. 302.

  Not in the orig. ed.; since described. See Ridg., Pr. Nat. Mus., ii, 1880, p. 8.
- 478. S. ōc-cĭ-dēn-tā'-lĭs. Lat. occidentalis, occidental, western, where the sun sets; occĭdo, I fall down (ob and cado, not occīdo, I slay).

  This is Syrnium occidentale of the orig. ed.
- 479. Nyc'-tĕ-ă scān-dĭ-ā'-că. Gr. Nυκτεύs, Lat. Nycteus, a proper name; as an adjective, nocturnal; Lat. nox, Gr. νύξ, night. There are very many derivatives, of which Nyctala is one.—Lat. Scandiaca, Skandinavian, relating to Scandia or Scandinavia.

- 480. Surnia funerea (L.) Rich. & Sw. B 62. C 326. R 407.

  American Hawk Owl.
- 481. Surnia funerea ulula (L.) Ridg. B —. C —. R 407. (?) (! A.) European Hawk Owl.
- 482. Nyctala tengmalmi richardsoni (Bp.) Ridg. B 55. C 327. R 400. Richardson's Owl.
- 483. Nyctala acadica (Gm.) Bp. B 56, 57. C 328. R 401.

  Acadian Owl; Saw-whet Owl.
- 484. Glaucidium gnoma Wagl. B 60. C 329. R 409. Pygmy Owl.
- 480. Sūr'-nī-ā fū-nĕ'-rī-ā. Surnia and Syrnium are forms of the same word, the meaning and derivation of which are alike unknown to us; we follow Newton in using the former; see Sund., Tent., p. 104. Lat. fimereus, funereal; from finus, a funeral, burial procession. Applicable to an owl, either regarded as a bird of ill omen, or with reference to its dismal cry, as if wailing the dead.

This stands Surnia ulula hudsonica in the orig. ed. Names of owls are "confusion worse confounded." See Ridg. Pr. Nat. Mus., ii, 1880, p. 8.

481. S. f. ŭ1-ŭ1-ă. Lat. ulula, a Plinian name of the screech-owl; ululo, Gr. ὀλολύζω, I howl, halloo, make a "hullaballoo"; all onomatopœic. Compare also the Hebrew, בְּבֶּק, whence hallelujah.

Not in the orig. ed. The old world Hawk Owl, at best hardly distinguishable from the American, is stated to occur in Alaska as a straggler from Asia; and all the Hawk Owls of Great Britain are said to be of the American variety. The case itself is as perplexing as its nomenclature is involved.

- 482. Nyc'-tă-lă těng'-măl-mī rīch'-ārd-sŏn-ī. Gr. νύκταλος or νύσταλος, drowsy, sleepy. See Nyctea, No. 479, for basis of the word.—To P. G. Tengmalm, a Swedish naturalist.—To Sir John Richardson, the English naturalist.
- 483. N. ă-căd'-ĭ-că. To Acadia, or Acadie, a locale now in Maine, scene of Longfellow's "Evangeline."
- 484. Glaū-cīd'-ĭ-ŭm gnō'-mā. There is a Greek word γλαυκίδιον, but that is some kind of fish, not a bird. It is, however, related to γλαθξ, which means an owl. There is also an adjective γλαυκώδης, from γλαῦξ and είδος, from which Glaucidium may be modified. The allusion in all these cases is to the eyes of the bird; if not in color, then in the general aspect and expression of these remarkable organs of vision. There being actually no owls with blue eyes, as γλαυκός, glaucus, is commonly translated, the direct implication is probably to the owl as the bird of wisdom, sacred to Minerva, γλαυκῶπις being one of the most familiar Homeric epithets of the "blue-eyed" goddess. Such may therefore be the meaning of  $\gamma \lambda \alpha \hat{v} \xi$ , without reference to the color of the bird's own eyes. —The word gnoma is very pat for an owl, and especially interesting in such application. Gr. γνώμα, an opinion, decision ; γνώμη, reason ; γνώμων, a judge, arbiter ; all from γιγνώσκω, I know; whence also gnostic, and the very English word know, with countless related forms, all rooted in the idea of knowledge. Hence gnoma is apt for the bird of Minerva, goddess of wisdom, and is given just as Athene was made a similar epithet. Furthermore, the English word gnome, by which we may directly translate gnoma in this case, is from the same root, meaning etymologically "the knowing one," "one who arbitrates certain destinies": by metonymy, a kind of sprite or elf presiding over mines. Gnoma is thus an eligible epithet of a bird which combines a reputation for wisdom with certain superstitions connected with the gnome-like or goblin-like quality of its knowingness.

- 485. Glaucidium ferrugineum (Maxim.) Kaup. B —. C 330. R 410. Ferrugineous Owl.
- 486. Micrathene whitneyi (Coop.) Coues. B —, C 331. R 411. Elf Owl.
- 487. Speotyto cunicularia hypogæa (Bp.) Coues. B 58, 59. C 332. R 408. Burrowing Owl.
- 488. Speotyto cunicularia floridana Ridg. B —. C —. R 408a. Florida Burrowing Owl.
- 489. Circus cyaneus hudsonius (L.) Coues. B 38. C 333. R 430. Marsh Hawk; Harrier.
- 490. Rostrhamus sociabilis plumbeus (—) Ridg. B 37. C 334. R 429. Everglade Kite.
- 491. Ictinia subcœrulea (Bartr.) Coues. B 36. C 335. R 428. Mississippi Kite.
- 492. Elanus glaucus (Bartr.) Coues. B 35. C 336. R 427. White-tailed or Black-shouldered Kite.
- 485. G. fēr-rū-gin'-ĕ-ŭm. Lat. ferrugineum, rusty-red; ferrugo, iron-rust; ferrum, iron.
- 486. Mī-cră-thēn'-ē whĭt'-nĕy-ī. Gr. μικρόs, small; 'Αθήνή or 'Αθηνά or 'Αθηνά, the Greek goddess of wisdom, to whom the owl was sacred. There was already a genus Athene, when Dr. Coues constructed the above. The genus Atthis, No. 416, is rooted with the same, as are Attic, Athens, Athenian, Athenœum, &c. To Professor J. D. Whitney, Director of the Geological Survey of California.
- 487. Spě-δ'-ty-tō cǔn-ī-cǔ-lā'-rī-ă hy-pŏ-gaē'-ă. Gr. σπέος, a cave, excavation; τυτώ, a kind of owl. The first refers to the burrowing of this species; the last, like ulula, is onomatopæic, in imitation of an owl's hooting or "tooting"; tyto, a "tooter."—Lat. cunicularius, a miner, burrower; cuniculus, a mine, pit, hole.—Lat. hypogæum, a vault, cellar; Gr. ὑπόγειος, under ground, subterranean; ὑπό, under, γέα, γη, the ground. Thus all three words refer to the same thing.
- 488. S. c. flōr-ĭd-ā'-nă. To Florida, "land of flowers."

  Not in the orig. ed.; since described; Ridg., Am. Sportsman, July 4, 1874, p. 216.
- 489. Cīr-cŭs cȳ-ăn'-ĕ-ŭs hŭd-sŏn'-ī-ŭs. Gr. κίρκοs, Lat. circus, a kind of hawk, so called from its circling in the air. Gr. κύανοs, Lat. cyaneus, blue; the color of the old male. To Hudson's Bay.
- 490. Röstr-hām'-ŭs sŏ-cĭ-ā'-bĭ-līs plūm'-bĕ-ŭs. Lat. rostrum, beak, and hamus, Gr. χαμόs, a hook, from the greatly decurved form of the upper mandible. It is a queerly compounded word, meaning literally bill-hook, though the person who invented it meant to say hook-bill, hamirostrum. It is very bad form as it stands, but we hardly know how to emend without entirely changing it.—Lat. sociabilis, sociable, gregarious; socius, a companion.— Lat. plumbeus, plumbeous, lead-colored.
- 491. Ic-tīn'-ī-ă sūb-coē-rŭl'-ĕ-ă. Gr. lκτίν or lκτίνοs, a kite; probably rooted same as ἴκτεροs, a disease, in the idea of attacking; Lat. ictus, a blow, &c.— Lat. sub, a prefix of diminishing force, and cαruleus, blue; bluish, pale blue. See Dendrαca, No. 117.
  - This stands as *I. mississippiensis* in the orig. ed. See Coues, Pr. Phila. Acad., 1875, p. 345.
- 492. Ĕl'-ăn-ŭs glaū'-cŭs. Lat. elanus, a kite; derived from the Gr. ἐλαύνω, I drive on, urge forward, press upon, harass, &c.; a good name for a bird of prey which exhibits what the French would call elan.—Lat. glaucus, Gr. γλαυκόs, bluish, glaucous; from λεύω, λεύσσω, I shine. See Glaucidium, No. 484.

This is Elanus leucurus in the orig. ed. See Coues, Pr. Phila. Acad., 1875, p. 345.

- 493. Elanoïdes forficatus (L., 1758) Coues. B 34. C 337. R 426. Swallow-tailed Kite.
- ✓494. Accipiter fuscus (Gm.) Bp. B 17. C 338. R 432. Sharp-shinned Hawk; Pigeon Hawk.
- 495. Accipiter cooperi Bp. B 15, 16. C 339. R 431. Cooper's Hawk; Chicken Hawk.
  - 496. Astur atricapillus (Wils.) Bp. B 14. C 340. R 433.

    American Goshawk.
  - 497. Astur atricapillus striatulus Ridg. B —. C —. R 433a. (?)
    Western Goshawk.
  - 498. Falco sacer Forst. B —. C 341. R 412b.

    American Continental Gyrfalcon.
  - 493. Ĕl-ăn-ŏ-ī'-dēs for-fī-cā'-tŭs. Lat. elanus (see No. 492) and Gr. εἶδοs, resemblance. For forficatus, see Milvulus, No. 367.

    This is Nauclerus furcatus of the orig. ed. See Coues, Bull. U. S. Geol. Surv. Terr., iv, No. 1, 1878, p. 42.
  - 494. Ac-cīp'-ĭ-tĕr fūs'-cŭs. Lat. accipiter, a general name for a hawk; accipio, I take, seize; from ad and capio: Gr. κάπτω, of similar meaning. Some, however, derive the word (as it seems to us, fancifully) from acuo and peto, i. e., the swift flyer. The root cap- is a very general one for words denoting this idea of taking; as in English accept, except, captive, capable, capacious, &c. Lat. fuscus, fuscous, dark-colored.
  - 495. A. coop'-ĕr-i. To William Cooper, of New York.
  - 496. As'-tūr ā-trī-că-pīl'-lūs. Lat. astur, a hawk; evidently related to aster, a star; asterias, starry, i. e., speckled; French autour is the same. The European Goshawk was called Asterias and "Star-hawk" by some of the old ornithologists, and the term ἀστερίας ἰέραξ is classic. The Italian is astore or asturo, and some dialectic form of this is said to give the name to the Açores or Azores Islands, from the abundance of hawks there. For atricapillus, see Parus, No. 44. The word gos- prefixed to hawk is Anglo-Saxon; goshafoc is goose-hawk; hafoc, and many similar words, are related to faucon, falcon, falco, which see, No. 498.
  - 497. A. a. strī-ā'-tŭ-lŭs. Lat. striatulus, diminutive of striatus, striate, streaked, striped; implying not the smallness of the streaked object, but the fineness of the stripes themselves.

    Not in the orig. ed. of the Check List. Since described by Ridg., Hist. N. A. B., iii, 1874, p. 240.
  - 493. Fāl'-cō să'-cĕr. Gr. φάλκων, Lat. filco, a falcon, from the falx, falcis, a sickle, scythe: in allusion to the falcate form of the hooked beak. The English is directly from falco, and the word reappears in many languages: Fr. faucon; Ital. falcone; Span. halcon, &c. The word Gyrfalcon or Jerfalcon has much exercised the ingenuity of the dictionaries. To us the etymology seems clear and indisputable. It is found in many forms, as ger-, gir-, gyr-, gir-, ier-, iier-, and this leads directly to lepós, divine, sacred, noble, auspicious, chief, &c.; lepeós, a priest; whence lépaţ, the actual Greek word for a hawk, as used in divination, and therefore sacred. The idea is the same as that in hierarch, &c. The English Gyrfalcon or Jerfalcon is therefore a mere transliteration of Hierofalco. In the same spirit, Steenstrup recently made a genus Gyralca for the principal bird of the auk tribe, already known in many vernaculars by a corresponding epithet. Speculations respecting gyr- as meaning gyrus, a whirl, from the hawk's gyrations, are superfluous. Lat. sacer, sacred, consecrated, sanctified, &c.; the root sac- is the Greek root άγ, as seen in ἄγιος, άγνός.

By the above name we indicate the continental Gyrfalcon of Arctic America, corre-

- 499. Falco sacer obsoletus (Gm.) Ridg. B —. C —. R 412c. Labrador Gyrfalcon.
- 500. Falco islandicus Gm. B 12. C —. R 412a. Iceland Gyrfalcon.
- 501. Falco candicans Gm. B 11. C 341a. R 412. (G.) Greenland Gyrfalcon.
- 502. Falco mexicanus Licht. B 10. C 342. R 413.

  American Lanier Falcon.
- 503. Falco peregrinus Tunstall. B 5, 6. C 343. R 414.

  Peregrine Falcon; Duck Hawk.
- 504. Falco peregrinus pealii (Ridg.) Coues. B —. C 343a. R 414a. (?)
  Peale's Peregrine Falcon.
- , 505. Falco columbarius L. B 7. C 344. R 417. Pigeon Hawk.
  - 506. Falco columbarius suckleyi Ridg. B —. C 344a. R 417a. (?) Suckley's Pigeon Hawk.
    - sponding to *F. gyrfalco* of Continental Europe, without raising the much-vexed question of their identity. We give the dark Labrador bird as a variety of this, and the Icelandic and Greenlandic as both specifically distinct; though we suppose all the northern *Hierofalcones* to be but geographical races of a single species.
  - 499. F. s. ōb-sŏl-ē'-tŭs. Lat. obsoletus, unaccustomed, unwonted, disused, obsolete; here referring simply to the ill-defined character of the markings; ob and soleo, I am accustomed.

    Not in orig. ed. This is Falvo labradorus of Audubon, lately accredited by Mr. Ridgway with varietal distinction, and identified with F. obsoletus Gm.
  - 500. F. is-lānd'-ī-cŭs. [ees-]. Latinized directly from the native name of Ice-land (Îsland, otherwise known as Eisland and Ijsland), and thus meaning Icelandic,—not "insular."
  - 501. F. cān'-dĭ-cans. Lat candico, I am white; present participle of the verb; candidus, white; candeo, I am shining, &c. Candid is pure, clean, hence truthful; candescent, brilliantly glowing; candidates were so called because clothed in white; candles give light; canescent hairs grow white; in all these, and countless words, the same root is seen.

    In the orig. ed. as Falco sacer var. candicans; see above, No. 498.
  - 502. F. mēx-ī-cā'-nūs. To Mexico, whence Lichtenstein described it. It has been identified with F. polyagrus of Cassin. "Lanier" or "Lanner" is the name applied in ornithology and falconry to certain Old World species; it is from laniarius, of a butcher, laniator, a butcher, from lanio, I lacerate, mangle; lanius (which see, No. 186) is the same thing.
  - 503. F. pĕr-ĕ-grī'-nŭs. See Helminthophaga, No. 109.
    This stands as F. communis in the orig. ed. It is well to stretch a point in favor of Tunstall, 1779, to be able to restore this well-known name.
  - 504. F. p. pēal'-ī-ī [in three syllables]. To Titian R. Peale, of United States Exploring Expedition fame. Of doubtful standing.
  - 505. F. cŏl-ūm-bā/-rĭ-ŭs. Post-classic Lat. columbarius, pertaining to a pigeon, columba; or, a pigeon-fancier, as this spirited little falcon is.
  - 506. F. c.  $s\bar{u}ck'-l\bar{e}y-\bar{i}$ . To George Suckley, known in ornithology for his researches in Oregon and Washington Territories. The first syllable is long, and pronounced with the full Latin force of  $\bar{u}$ , like oo in moon. A very dubious bird.

- 507. Falco columbarius richardsoni Ridg. B —. C 345. R 418. Richardson's Pigeon Hawk.
- 508. Falco sparverius L. B 13. C 346, R 420. Sparrow Hawk.
- 509. Falco sparverius isabellinus (Sw.) Ridg. B —. C 346a. R 420a. Isabel Sparrow Hawk.
- 510. Falco sparverioïdes Vig. B —. C —. R 421. (! W. I.) Cuban Sparrow Hawk.
- 511. Falco fuscicærulescens V. B 9. C 347. R 419. Femoral Falcon.
- 512. Buteo unicinctus harrisi (Aud.) Ridg. B 46. C 348. R 434. Harris's Buzzard Hawk.
- 513. Buteo albocaudatus V. B —, C —, R 441.
  White-tailed Buzzard Hawk.
- 507. F. e. rĭch'-ārd-sŏn-ī. To Sir John Richardson, the species having been described and figured in the Fauna Boreali-Americana.
- 508. F. spār-vě'-rĭ-ŭs. Post-classic Latin, meaning, relating to a sparrow, as columbarius from columba. There is a quasi-Latin word sparvius, from which sparverius is directly formed. The word sparrow in some of its forms doubtless antedates any corresponding word in the South European languages. We have not traced the Latin sparvius or sparverius back of Gesner, 1555. See Passer, No. 192.
- 509 **F. s.** i-să-bēl-li'-nŭs. The Lady Isabel, having confidence in her husband's prowess, vowed not to change her chemise until that warrior had taken a certain town. He was longer about it than she expected, and she wore the garment until it assumed a peculiar brown tint: hence the term "isabel-color"; whence quasi-Latin isabellinus.
- 510. F. spār-vĕ-rī-ŏ-ī'-dēs. This is an aggravated case of bastardy. Anglo-Saxon and Gothic sparwa or sparra, Latinized as sparvius, a sparrow, whence sparverius, a sparrower, so to speak, or sparrow-catcher, as this hawk is; with the Gr. εἶδοs, to denote the resemblance of the West Indian to the North American bird.

Not in the orig. ed. of the Check List. Lately said to have occurred in Florida. See Ridg., Pr. Nat. Mus., iii, 1880, p. 220.

- 511. F. fūs-cī-coē-rǔl-ēs'-cēns. Lat. fuscus, dusky, and cœrulescens, growing blue; i. e., being bluish: cœruleus, blue. This was written fuscocærulescens by Vieillot, but the above is preferable. "Femoral" relates to the color of the thigh; femur, the thigh-bone.

  This is F. femoralis of the orig. ed. See Sharpe, Cat. Accip. Br. Mus., i. p. 400.
- 512. Bū'-tě-ō ū-nǐ-cīnc'-tǔs hǎr'-rǐs-ī. Lat. buteo, a buzzard-hawk; of doubtful etymology; the word occurs in Pliny. Lat. uni, once, and cinctus, girded; unus, one, and cingo, I gird, bind about; with reference to the single zone of white color on the tail. To Edward Harris, of Philadelphia.
- 513. B. āl-bŏ-caūd-ā'-tŭs. Lat. albus, white, caudatus, tailed; cauda, tail. The latter part of the word being a participial adjective of a supposed verb caudo, permits albus to be in the "ablative of instrument," "white" being that wherewith the bird is "tailed." In another form, it would be albicauda, like albicilla for instance. See No 42.

Not in the orig. ed.; since discovered in Texas both by G. B. Sennett and J. C. Merrill. See Coues, The Country, July 13, 1878, p. 184; and Ridg., Pr. Nat. Mus., i, Oct. 2, 1878, p. 154.

- 514. Buteo cooperi Cass. B 29. C 349. R 437. (?)
  Cooper's Buzzard Hawk.
- 515. Buteo harlani (Aud.) Bp. B 22. C 350. R 438. Harlan's Buzzard Hawk.
- 516. Buteo borealis (Gm.) V. B 23. C 351. R 436. Red-tailed Buzzard Hawk; Hen Hawk.
- 517. Buteo borealis calurus (Cass.) Ridg. B 20, 24. C 351a. R 436b.
  Western Red-tailed Buzzard Hawk.
- 518. Buteo borealis lucasanus Ridg. B —. C 351b. R 436c. St. Lucas Buzzard Hawk.
- 519. Buteo borealis krideri Hoopes. B —, C 351c. R 436a. (?) Krider's Buzzard Hawk.
- 520. Buteo lineatus (Gm.) Jard. B 25. C 352. R 439. Red-shouldered Buzzard Hawk.
- 521. Buteo lineatus elegans (Cass.) Ridg. B 26. C 352a. R 439a. Western Red-shouldered Buzzard Hawk.
- 522. Buteo abbreviatus Cab. B —. C 353. R 440.
  Band-tailed Hawk.
- 523. Buteo swainsoni Bp. B 18, 19, 21, 28. C 354. R 442. Swainson's Buzzard Hawk.
- 514. B. coop'-ĕr-ī. To Dr. James G. Cooper, of California, well known for his studies of the birds of that country. Doubtful species: only one specimen known.
- 515. B. hār'-lăn-î. To Dr. Richard Harlan, of Philadelphia, author of Medical and Physical Researches, Fauna Americana, etc.
- 516. B. bor-ĕ-ā'-līs. Lat. borcalis, northern; boreas, the north wind.
- 517. B. b. căl-ū'-rŭs. Gr. καλός, beautiful, and οὖρα, tail.
- 518. B. b. lū-căs-ā'-nŭs. Named after Cape St. Lucas, Lower California.
- 519. B. b. krī'-dĕr-ī. To John Krider, the veteran taxidermist of Philadelphia. Dubious.
- 520. B. lī-nē-ā/-tŭs. Lat. lineatus, lineated, limned, from linio; linea, a line. In reference to the streaking of the plumage.
- 521. B. 1. ē'-lĕ-gāns. Lat. elegans, elegant, because select, chosen: e and ligo, I pick out.
- 522. B. āb-brēv-Ĭ-ā'-tŭs. Lat. abbreviatus, shortened; ab and brevio, I abridge, contract; brevis, short; Gr. βραχύs. Applicability unknown to us.

  This stands as B. zonocercus in the orig. ed. See Ridg., Pr. Nat. Mus., iii, 1880, p. 220.
  - 523. B. swaīn'-sŏn-ī. To William Swainson, Esq., the celebrated English naturalist.
    Mr. Sharpe has lately called this B. obsoletus (Gm.), but very erroneously, Gmelin's bird of that name being a Gyrfalcon. B. insignatus of Cassin is simply a melanism.
     B. bairdi of Cassin is the young. This bird is the nearest form we have to the European B. vulgaris, which latter has been attributed to Michigan: see Maynard, Bull. Nutt. Club, i, No. 1, 1876, pp. 2-6.

The meaning of the word "buzzard" is unknown to us. It runs through several languages, as buzhard, buzard, busard, buse. Some think it onomatopæic, related to buzz; that seems doubtful; more likely related to the Latin buteo. Butes is a Latin proper name, but of no obvious connection.

- 524. Buteo pennsylvanicus (Wils.) Bp. B 27. C 355. R 443.

  Broad-winged Buzzard Hawk. See Addenda, Nos. 882, 883.
- 525. Archibuteo lagopus sancti-johannis (Gm.) Ridg. B 30, 31. C 356. R 447. American Rough-legged Buzzard.
- 526. Archibuteo ferrugineus (Licht.) Gr. B 32. C 357. R 448. Ferrugineous Rough-legged Buzzard.
- 527. Asturina plagata Schl. B 33. C 358. R 445. Gray Hawk.
- 528. Urubitinga anthracina (Licht.) Lafr. B —. C —. R 444.

  Anthracite Hawk.
- 529. Onychotes gruberi Ridg. B —, C 359. R 446. Gruber's Hawk.
- 530. Pandion haliaëtus (L.) Sav. B 44. C 360. R 425. Fish Hawk; Osprey.
- 531. Thrasyaëtus harpyia (L.) Gr. B —, C —, R 450. (! M.) Harpy Eagle.
  - 524. B. pēnn-syl-văn'-ĭ-cŭs. See Dendræca, No. 124.
  - 525. Arch-ĭ-bū'-tĕ-ō lਬg-ō'-pūs sānc-tĭ-jŏ-hān'-nīs. Lat. archi-, equivalent to Gr. ἀρχόs, a leader, a chief; ἄρχω, I rule, I am first; the word simply means "arch-buzzard," like archbishop, archetype, architect, &c. Lat. lagōpus, Gr. λαγώπουs, hare-footed, from λαγώs, a hare, and ποῦs, a foot: in allusion to the feathering of the tarsi. The penult here remains long in Latin as it is in Greek; but words in -opus, where the o is simply a connecting vowel, shorten the penult. Lat. sancti-johannis, of Saint John, alluding to the place in Newfoundland so called.
  - 526. A. fēr-rū-gĭn'-ĕ-ŭs. See Scolecophagus, No. 331.
  - 527. As-tūr-ī'-nă plā-gā'-tă. Asturina is simply formed from Lat. astur, which see, No. 496, without any difference of meaning. Lat. plagata, striped, from plago, I strike; plaga, a blow, stroke, stripe; Gr. πληγή, a blow, wound, from πλήσσω or πλήττω, I strike. Commonly written plagiata, for which we see no good reason.
  - 528. U-rū-bǐ-tīn'-gặ ān-thrặ-cī'-nặ. Urubitinga is a barbarous word, of some South American dialect; urubu means a vulture; we do not know what the rest of the word is, nor the quantity of the first two vowels; we hear them long and leave them so. Lat. anthracinus, Gr. ἀνθράκινος, carbuncular; ἄνθραξ, genitive ἄνθρακος, a carbuncle; also a live coal, a coal. The application in the present case is not to a glowing coal, like a carbuncle, but to a dead coal, coal-black; the glossy black of anthracite coal, as the bird is.
  - 529. Ŏ-nych'-ŏ-tēs grū'-bĕr-ī. Gr. ὅνυξ, genitive ὅνυχος, a claw; the rest of the word is the regular suffix -της, -tes, making the whole signify "the clawed one." Notice the accent.

     To F. Gruber, a taxidermist of San Francisco.
     This bird is questionably North American; but distinct from any Hawk in this list.
  - 530. Pān-dī'-ōn hāl-i-ā-ē'-tūs. Lat. Pandion, Gr. Πανδίων, was the alleged father of Progne and Philomela: see Coues, B. Col. Vall., i, 1878, p. 371. Observe quantity and accent of the penult.—Gr. äλs, genitive άλδs, salt, the sea, and ἀητόs, an eagle; "sea-eagle." See Haliaëtus, No. 533.
  - 531. Thră-sỹ-ā-ē'-tũs hār-pỹ'-ă or hār-pỹ'-1-ă [either three or four syllables; in either case pronounced harpwee'ah]. Gr. θρασύs, bold, audacious, and ἀητόs, eagle; see No. 533. Generally written Thrasaëtus, as originally by Gray: but the above is preferable; compare Thrasyas, Thrasybulus, Thrasymachus, &c., all retaining the y (v).—The "Αρπυιαι.

- 532. Aquila chrysaëtus (L.) Cuv. B 39. C 361. R 449. Golden Eagle.
- 533. Haliaëtus albicilla (L.) Leach. B 42. C —. R 452. (G.) White-tailed Eagle; Sea Eagle.
- 534. Haliaëtus leucocephalus (L.) Savig. B 41, 43. C 362. R 451. White-headed Eagle; Bald Eagle.
- 535. Polyborus cheriway (Jacq.) Cab. B 45. C 363. R 423. Caracara Eagle.

Harpyie or Harpies were fabulous monsters, embodying the idea of female rapacity as birds of prey, with crooked talons and beak  $(\tilde{a}_{\rho}\pi\eta)$ .

Not in the orig. ed. of the Check List; lately ascertained to occur in Texas. See Oswald, Am. Nat., 1878, p. 151; and Ridg., Pr. Nat. Mus., iii, 1880, p. 221.

- 532. A'-quīl-ă chrys-ā-ē'-tūs. Lat. aquila, an eagle. The etymology is disputed. It is given by some, without qualification, as from Gr. ἀκύς, Lat. acer, ocior, sharp, swift, from √ac or √ak. Some say from aquilus, dark, swarthy; others, as related to aquilo, the north wind; others from Gr. ἀγκύλος, crooked, hooked, as the bird's beak is: this would correspond to the derivation of gryps, γρύψ, a griffin, from γρυπός, bent, hook-nosed. It is conjectured, also, from ἀγκύλη, the curve of the limb, or the curved limb, with which the bird, as Jove's lightning-bearer, grasped the thunder-bolts. Some allied forms of the word, in which g appears instead of the q, as aguila, aigle, eagle, favor the supposition that the name has something to do with the great wings of the bird. Gr. χρυσαίετος or χρυσάετος, golden eagle; χρυσός, golden, ἀετός, eagle. See Haliaĕtus, No. 533.
- 533. Hăl-ĭ-ā-ē'-từs āl-bĭ-cīl'-lă. Gr. άλς, genitive άλός, salt; the (salt) sea; and ἀετός or ἀητός or αἰετός, an eagle; there is also the actual Greek ἁλιάετος or ἀλιαίετος, for the "seaeagle," that is, the osprey. There is also the actual Latin transliteration "haliæetos," for the same bird. So many vowels coming together, with such variation in the original Greek, has kept the orthography incessantly fluctuating. Savigny, who was a classical scholar, as well as an ornithologist, originally spelled the genus he founded Haliæëtus. This is perfectly correct, in fact, the poetic form, as transliterated from άλιαlετος, with only the usual and proper change of Greek  $\alpha_i$  into Latin  $\alpha$ . Many purists keep to this spelling, which is perfectly defensible, and has the advantage of being that used by the founder of the genus. But, as Haldeman remarks, however desirable Haliwëtus may be in poetical writing, it is more consonant with a strict scientific spirit to simplify the word into Haliaëtus, deriving it in this case from ἀετός or ἀητός. We accept and adopt this form upon such understanding. Having settled this, the next question arises respecting the quantity of the vowels, and accentuation of the syllables. If derived from ἀετός, the word would be  $H\bar{a}li\bar{a}'\bar{e}t\bar{u}s$ ; if from  $a\eta\tau\delta s$ , it would be  $H\bar{a}li\bar{a}\bar{e}'t\bar{u}s$ . We prefer the latter. In any event, the form "Haliætus," in four syllables, is inadmissible: the word must have at least five syllables. But ornithologists may be forgiven for anything in this case, seeing that the grammarians have disputed it for some centuries. - Lat. albicilla, white-tailed. See Motacilla, No. 86.

This species, though frequently attributed to North America, has of late years been dropped. It is now restored, on the strength of its occurrence in Greenland, though not elsewhere in North America that we know of. Not in the orig. ed. of the Check List.

- 534. H. leū-cŏ-cĕph'-ă-lŭs. Gr. λευκός, white, and κεφαλή, head.
- 535. Pŏl-y̆'-bŏr-ŭs cheriway. Gr. πολυβόρος, eating a great deal, very voracious. Cheriway and Caracara are both barbarous words, the meaning of which we know not: from some South American dialect.

This stands in the orig. ed. as P. tharus var. auduboni.

- 536. Pseudogryphus californianus (Shaw) Ridg. B 2. C 364. R 453. Californian Vulture.
- 537. Cathartes aura (L.) Ill. B 1. C 365. R 454. Turkey Buzzard.
- 538. Catharista atrata (Bartr.) Less. B 3. C 366. R 455. Carrion Crow.
- 539. Columba fasciata Say. B 445. C. 367. R 456.
  Band-tailed Pigeon.
- 540. Columba erythrina Licht. B 446. C 368. R 457. Red-billed Pigeon.
- 541. Columba leucocephala L. B 447. C 369. R 458. White-crowned Pigeon.
- 542. Engyptila albifrons (Bp.) Coues. B —. C —. R 463. White-fronted Pigeon.
- 536. Pseū-dŏ-gry/-phùs căl-ĭ-fōr-nĭ-ā'-nùs. Gr. ψεῦδος, false, from ψεύδω, I deceive, and Lat. gryphus, for gryps, genitive gryphis, a griffin, a fabulous bird; from Gr. γρύψ, the same, from γρυπός, bent, hook-nosed. The word is badly formed in two languages: had better have been Pseudogryps. Gryphus is a name early transferred by ornithologists from its fabulous prototype to the condor of the Andes; and Mr. Ridgway made Pseudogryphus from the resemblance of the Californian vulture to the latter.

This stands as Cathartes cal. in the orig. ed. See Ridg., Bull. Nutt. Club, v, 1880, p. 79.

- 537. Căth-ār'-tēs aū'-ră [ow-rah, not or-ah]. Gr. καθαρτής, a purifier, from καθαίρω, I cleanse, purify, purge; from the good offices of the bird as a scavenger in warm countries.—
  Aura is a name applied to this bird by the oldest writers who speak of it, and, in all its various forms, as rendered by De Laet and others who treat of tropical American Cathartidæ, it is of South American or Mexican origin, and apparently related to urubu or ourubu. It early crystallized in its present orthography, and was soon Latinized, or at least declined as a Latin word; as, rex aurarum, or regina aurarum (genitive plural), "king of the vultures." That it has any connection with Lat. aura, Gr. aδρα, air, atmosphere, may well be doubted.
- 538. Căth-ăr-is'-tă ā-trā'-tă. Badly framed from καθαρίζω, only another form of καθαίρω, of same meaning; see No. 537.—Lat. atrata, participial adjective, blackened; ater, black. This stands as Cathartes atratus in the orig. ed. See Ridg., Bull. Nutt. Club. v, 1880, p. 80.
- 539. Cŏl-ūm'-bă fās-cĭ-ā'-tă. Lat. columba, a pigeon; etymology unknown.—See Chamæa, No. 39.
- 540. C. ĕ-ryth-rī'-nā. Lat. erythrina, Gr. ἐρυθρῖνος, reddish; from ἐρυθρός, red. This is C. flavirostris of the orig. ed. As the bill is not at all yellow, another name is desirable. See Ridg., Pr. Nat. Mus., ii, 1880, p. 9.
- 541. C. leū-cŏ-cĕph/-ă-lă. Gr. λευκός, white, and κεφαλή, head.
- 542. En-gyp'-tı-lā āl'-bı-frōns. Gr. έγγύς, narrow, slender, contracted, and πλίλον, a feather; from the attenuated outer primaries. Lat. albus, white; from, forehead.

Not in the orig. ed.; since discovered in Texas by G. B. Sennett. See Coues, Bull. U. S. Geol. Surv. Terr., iv, 1878, p. 48, and Bull. Nutt. Club, v, 1880, p. 100; Ridg., Pr. Nat. Mus., i, 1878, p. 158.

- 543. Ectopistes migratorius (L.) Sw. B 448. C 370. R 459. Wild Pigeon; Passenger Pigeon.
- 544. Zenaidura carolinensis (L.) Bp. B 451. C 371. R 460. Carolina Dove.
  - 545. Zenaida amabilis Bp. B 449. C 372. R 462. Zenaida Dove.
  - 546. Melopelia leucoptera (L.) Bp. B 450. C 373. R 464. White-winged Dove.
- 547. Chamæpelia passerina (L.) Sw. B 453. C 374. R 465. Ground Dove.
- 548. Chamæpelia passerina pallescens (Bd.) Coues. B—. C 374a. R—. (?) St. Lucas Ground Dove.
- 549. Scardafella inca (Less.) Bp. B 452. C 375. R 466. Scaled Dove.
- 550. Geotrygon martinica (Gm.) Reich. B 454. C 376. R 467. Key West Pigeon.
- 543. Ec-tŏ-pīs'-tēs mī-gră-tō'-rĭ-ŭs. Gr. ἐκτοπιστής, a wanderer, passenger; ἐκτοπίζω, I wander, change place; from ἐκ, out of, and τόπος, place; "out of place."—Lat. migratorius, of same meaning; migro, I migrate.
- 544. Zĕn-ă-ī-dū'-ră cā-rŏ-līn-ēn'-sīs. We think zenaida is a barbarous word. Its meaning we do not know. See Phonipara, No. 297, and compare zena there given. The rest of the word is formed by adding the Greek οὐρά. Bonaparte originally wrote zenaidura, which has usually, of late, following Dr. Coues' lead, been turned to zenædura; but if the word is not classic, there is no occasion for the modification.
- 545. Zěn-š-ī'-dă ăm-ā'-bīl-īs. Zenaïda, a proper name, perhaps Spanish; meaning unknown to us: see No. 544. Lat. amabilis, lovable, lovely; amo, I love.
- 546. Měl-ŏ-pěl-i'-ă leū-cōp'-těr-ă. Gr. μέλος, melody, and πέλεια, a dove. Name derived from πέλλος, the peculiar dark slaty-blue color, so characteristic of pigeons; we say to-day in sporting parlance "blue-rocks" for the ordinary domestic pigeon. The word, like many others ending in -pelia, is often wrong-written -peleia. Observe that the Greek ει becomes long i in Latin, giving us -pelia, accented on the penult. Gr. λευκός, white, and πτερόν, a wing.
- 547. Chăm-aē-pĕl-ī'-ă pās-sĕr-ī'-nă. Gr. χαμαl, an adverb, on the ground, and πέλεια, a dove. See No. 546. See Chamæa, No. 39. This word is spelled about a dozen different ways, by writers or printers who are careless or ignorant. Lat. passerina, sparrow-like, in allusion to the diminutive size: passer, a sparrow. See No. 192.
- 548. C. p. pāl-lēs'-cēns. See Mitrephorus, No. 392. Scarcely distinguishable from No. 547.
- 549. Scār-dă-fēl'-lă īn'-că. Scardafella is an Italian word, thus accounted for by Bonaparte, who founded the genus, in his "Coup d'Œil sur l'Ordre des Pigeons" (p. 43 of the separate copies): "une expression du Dante m'a inspiré le nom de scardafella, qui peint l'apparence écailleuse de notre treizième genre." The "scaly appearance" is due to the coloration, not the texture, of the feathers. Inca is a barbarous word; the incas or yncas were Peruvian chiefs.

This is S. squamosa var. inca in the orig. ed.; later determined to be distinct.

550. Gĕ-ŏ-trȳ'-gōn mār-tĭn'-ĭ-că. Gr. γέα, the earth, the ground, and τρυγών, a pigeon; from τρύζω, to coo; onomatopœic, like turtur. There seems to be reason for keeping the penult long, and accenting it.—Lat. martinica, Latinized adjective from Martinique, one of the West Indies.

- 551. Starnænas cyanocephalus (L.) Bp. B 455. C 377. R 468. Blue-headed Pigeon.
- 552. Ortalis vetula maccalli (Bd.) B 456. C 378. R 469.

  Texan Guan.
- 553. Meleagris gallipavo L. B 458. C 379. R 470. Domestic Turkey; Mexican Turkey.
- 554. Meleagris gallipavo americana (Bartr.) Coues. B 457. C 379a. R 470a. Common Wild Turkey of the United States.
- -555. Canace canadensis (L.) Bp. B 460. C 380. R 472.
  Canada Grouse; Spruce Partridge.
  - 556. Canace canadensis franklini (Dougl.) Coues. B 461. C 380a. R 472a. Franklin's Spruce Partridge.
  - 557. Canace obscura (Say) Bp. B 459. C 381. R 471. Dusky Grouse.
  - 558. Canace obscura richardsoni (Dougl.) Coues. B—. C 381a. R 471b. Richardson's Dusky Grouse.
  - 551. Stār-noē'-năs cy-ăn-ŏ-ceph'-ă-lŭs. From ——? (probably Italian; Agassiz gives Starna as a proper name), and Gr. olvás, Lat. œnas, the vine: also, a kind of pigeon; αnas seems to have been transferred to the pigeon, as œnanthe was to some other bird; see Saxicola, No. 26. The olvás of Aristotle is Columba livia L. Gr. κυανόs, cyanus, blue, and κεφαλή, head.
  - 552. Or'-tăl-ĭs vĕt'-ŭ-lă măc-căl'-lī. Gr. ὀρταλls, a pullet, a kind of quail. This word was universally written ortalida, until Mr. Wharton showed that the way Merrem, writing Latin, constructed the sentence in which the word first occurs made it the accusative case; arguing hence that Merrem meant to found a genus ortalis, not ortalida. See Ibis, October, 1879, p. 450. The Rev. Mr. Avery's MS. in our possession makes the same correction, though without comment. Lat. vetula, a little old woman; derisive diminutive from vetus, old, veteran; digammated from Gr. ĕτos, a year. To General George A. McCall, U. S. Army.
  - 553. Měl-ĕ-āg'-rīs gāl-lī-pā'-vō. Gr. μελεαγρίς, Lat. meleagris, a guinea-hen; literally, a field-tender, farmer; from μέλει, relating to the care of a thing, and ἄγρος, a field. The word not transferred from the African Numida to the American Turkey until near the middle of the 16th century, and occasionally confounded for many years after that. Meleager or Μελεαγρός was a mythical person who suffered a cruel fate: his sisters, the Meleagrides, who bitterly lamented his death, were changed into guinea-hens; the profusely-spotted plumage of which gives evidence of the tears they shed for him. Lat. gallipavo, usually written gallopavo, a very late combination of gallus, a cock, and paro, a pea-fowl, bird of Juno; the latter word from the Gr. ταώς or ταῶν, a pea-fowl.
  - 554. M. g. ăm-ĕr-ĭ-cā'-nă. Of America.
  - 555. Căn'-ă-cē că-nă-dēn'-sis. Canace, a proper name; she lived in incest with her brother; application not obvious, unless referring in a general way to the polygamy of gallinaceous birds.

This and following species are given as *Tetrao* in the orig. ed.; but may be properly separated generically from *Tetrao wogallus*.

- 556. C. c. frank'-lin-i. To Sir John Franklin, of Arctic fame and sorrow.
- 557. C. ōb-scū'-rus. Lat. obscurus, obscure, i. e., dark-colored.
- 558. C. o. rīch/-ārd-sŏn-ī. To Sir John Richardson, often already mentioned in this List.

- 559. Canace obscura fuliginosa Ridg. B —. C 381b. R 471a. Fuliginous Dusky Grouse.
- 560. Centrocercus urophasianus (Bp.) Sw. B 462. C 382. R 479. Sage-cock; Cock-of-the-Plains.
- 561. Pediœcetes phasianellus (L.) Elliot. B —. C 383. R 478.

  Northern Sharp-tailed Grouse.
- 562. Pediœcetes phasianellus columbianus (Ord) Coues. B 463, C 283a. R478a. Common Sharp-tailed Grouse; Prairie Hen of the Northwest.
- 563. Cupidonia cupido (L.) Bd. B 464. C 384. R 477.

  Pinnated Grouse; Prairie Hen.
- 564. Cupidonia cupido pallidicineta Ridg. B —. C 384a. R 477a. Pale Pinnated Grouse.
- 565. Bonasa umbella (L.) Steph. B 465. C 385. R 473.
   Ruffed Grouse; "Pheasant" in the Middle and Southern States.
  - 559. C. o. fū-lī-gĭn-ō'-să. Lat., post-classic, fuliginosa, of a dark sooty color; fuligo, soot; fulica, or fulix, a coot; so called from its color.
  - 560. Cēn-trŏ-cēr'-cŭs ū-rŏ-phā-sĭ-ā'-nŭs. Gr. κέντρον, a spine, and κέρκοs, tail; "sharptailed."—Gr. οὖρα, tail, and φασιανόs, Lat. phasianus, Fr. faisan, Engl. pheasant, pertaining to the river Phasis in Colchis. The scientific name of the English pheasant is Phasianus colchicus.\* The name "pheasant" has been ignorantly transferred to various American birds of this family.
  - 561. Pĕd-ĭ-oē'-cĕ-tēs phā-sǐ-ăn-ēl'-lŭs. Gr. πεδίον, a plain; as we should say, prairie; from πέδον, the ground; and οἰκέτης, an inhabitant; see Poæcetes, No. 232. The word was originally written Pediocætes. Lat. phasianellus, diminutive of phasianus; see Centrocercus, No. 560.
  - 562. P. p. cŏl-ūm-bĭ-ā/-nŭs. To the Columbia river, whence the birds were brought by Lewis and Clarke.
  - 563. Cŭ-pī-dŏ'-nĭ-ă cŭ-pī'-dō. The bird was named by Linnæus Tetrao cupido, after the "blind bow-boy," son of Venus, not with any allusion to erotic concerns, but because the little wings on the bird's neck were likened to "Cupid's wings." The same idea is repeated in the English "pinnated" grouse. Professor Reichenbach formed his genus Cupidonia by merely adding a suffix. If he had written cupidinea, he would have had a classic word, directly formed, like cupidus, from cupido, exactly expressing the sense intended by Linnæus to be conveyed.—The Latin tetrao, from the Gr. τετράων, and tetrix, from the Gr. τέτριξ, were certain gallinaceous birds, so called from their wont to cackle, τετράζειν: all onomatopœic.
  - 564. C. c. pāl-lī-dī-cīnc'-tă. Lat. pallidus, pallid, pale; and cinctus, begirt, encircled; cingo, I bind.
  - 565. Bŏn-ā'-să ūm-bēl'-lŭs. Gr. βόνασος, Lat. bonasus, a wild bull. The allusion here is to the "drumming" noise made by the bird, likened to the bellowing of a bull; see Bubo, No. 462, and Botaurus, No. 666. Also written Bonasia.—Lat. umbellus, or umbella, an umbel, umbrella; from umbra, shade, shadow, whence penumbra, umbrageous, &c. The allusion is to the tuft of feathers on the side of the neck, as in the case of cupido, which see, No. 563. Linnæus wrote Tetrao umbellus, masculine; but we see no reason why umbella, the noun feminine, should not be used with Bonasa; it is equally good Latin. The adjective umbellata would be preferable to either.

- 566. Bonasa umbella umbelloïdes (Dougl.) Bd. B 465\*. C 385a. R 473a. Gray Ruffed Grouse.
- 567. Bonasa umbella sabinii (Dougl.) Coues. B 466. C 385b. R 473b. Oregon Ruffed Grouse.
- 568. Lagopus albus (Gm.) Aud. B 467, 470?. C 386. R 474. Willow Ptarmigan.
- 569. Lagopus rupestris (Gm.) Leach. B 468. C 387. R 475. Rock Ptarmigan.
- 570. Lagopus leucurus Sw. B 469. C 388. R 476. White-tailed Ptarmigan.
- •571. Ortyx virginiana (L.) Bp. B 471. C 389. R 480. Virginia Partridge; Quail; Bob-white.
- 572. Ortyx virginiana floridana Coues. B —. C 389a. R 480a. Florida Partridge.
- 573. Ortyx virginiana texana (Lawr.) Coues. B 472. C 389b. R 480b. Texas Partridge.
- 574. Orortyx picta (Dougl.) Bd. B 473. C 390. R 481.

  Plumed Partridge; California Mountain Quail.
- 575. Lophortyx californica (Shaw) Bp. B 474. C 391. R 482. Crested Partridge; California Valley Quail.
- 576. Lophortyx gambeli Nutt. B 475. C 392. R 483. Gambel's Crested Partridge; Arizona Quail.
- 577. Callipepla squamata (Vig.) Gr. B 476. C 393. R 484. Scaled Blue Partridge.
- 566. B. u. ūm-bēl-lŏ-ī'-dēs. Lat. umbellus, which see, next above, and είδος.
- 567. B. u. sā-bī'-nĭ-ī. To J. Sabine.
- 568. Lăg-ō'-pǔs āl'-bǔs. Gr. λαγώπους, Lat. lagopus, hare-foot; λαγώς, a hare, and ποῦς, foot.— Lat. albus, white. For the length of the accented penult, see Archibuteo, No. 525.
- 569. L. rū-pēs/-tris. Late Lat. rupestris, pertaining to, or inhabiting, rocks; rupes, a rock.
- 570. L. leū-cū'-rŭs. Gr. λευκός, white, οὖρα, tail.
- 571. Or'-tyx vīr-gǐn-ĭ-ā'-nă. Gr. ὅρτυξ, a quail; related to ὁρταλίς; both are akin to ὅρτυς, a bird. The word is masculine in Greek, but in transliteration into Latin becomes feminine, like other nouns of same termination.—The English word partridge, Scot. patrick, Fr. perdrix, Span. perdiz, Ital. perdice, Lat. perdix, Gr. πέρδιξ, are all the same.
- 572. O. v. flo-ri-da/-na. To Florida.
- 573. O. v. tēx-ā'-nă. To Texas.
- 574. Ōr-ōr-tyx pic'-tă. Gr. ὅρος, a mountain, and ὅρτυξ; see Oroscoptes, No. 14.—Lat. pictus, painted, depicted; pingo, I paint; in allusion to the beautiful colors.
- 575. Loph-or'-tyx cal-ĭ-for'-nĭ-ca. Gr. λόφος, a crest, helmet, and ὅρτυξ.
- 576. L. găm'-běl-ī. To William Gambel, of Philadelphia. See Zonotrichia, No. 278.
- 577. Cāl-lǐ-pēp'-lă squā-mā'-tă. Gr. καλός, feminine καλλή, and πέπλος, a certain robe of state; καλλιπέπλος, beautifully robed, as this quail is.— Lat. squamata, squamous, scaled, covered with scales, the peculiar colors presenting such an appearance; squama, a scale.

- 578. Cyrtonyx massena (Less.) Gould. B 477. C 394. R 485.

  Massena Partridge.
- 579. Coturnix dactylisonans Meyer. B —. C —. R —. Migratory Quail (imported).
- 580. Squatarola helvetica (L.) Cuv. B 510. C 395. R 513. Black-bellied Plover; Bull-head.
- 581. Charadrius dominicus Müll. B 503. C 396. R 515.

  American Golden Plover.
- 582. Charadrius dominicus fulvus (Gm.) Ridg. B —. C —. R 515a. (! A.)
  Asiatic Golden Plover.
- 583. Charadrius pluvialis L. B —. C —. R 514. (G.) European Golden Plover.
- 578. Cÿr-tō'-nÿx mās-sē'-nă. Gr. κυρτόs, bent, curved, crooked, and ὅνυξ, a claw, nail; related to Lat. uncūs, a hook.— To the French Marshal André Massena, Prince d'Essling.
- 579. Cō-tūr'-nĭx dāc-tỹl-ĭ'-sŏn-āns. Lat. coturnix, a quail; onomatopœic, a sono vocis, from the sound of the voice, just as we have invented "bob-white" and "whip-poorwill."—Lat. dactylisonans, sounding a dactyle. The dactyle, in poetry, is a foot consisting of a long and two short syllables; from δάκτυλος, the finger, which has a long and two short joints. Sono, I sound; sonorous, &c.

This bird, lately imported, has become naturalized, with the same right to a place in the list that *Passer domesticus* has acquired.

- 580. Squă-tă-rō'-lă hēl-vē'-tī-că. Of squatarola the authors learned little, until a note from Professor Newton supplied the desired information, in substance as follows: As a generic term it is of course from the Linnæan Tringa squatarola, and Linnæus obviously got his trivial name from Willughby, who says (Ornith., ed. 1676, p. 229), "Pluvialis cinerea. Squatarola Venetiis dicta, ubi frequens est. The Gray Plover." The word is not to be found in the best Italian dictionaries; but Salvadori, in his Fauna d'Italia Uccelli, seems to acknowledge it as a genuine word; though probably it is only local in its application. It may possibly have to do with the regular Italian squartare, "to quarter." Lat. helvetica, from ancient Helvetia, now Switzerland; the bird is still often called "Swiss plover." The Helvetians were probably so called from their fairness, with flaxen or auburn hair; helves, helveolus (related to gilvus), meaning some such color.
- 581. Chăr-ād'-rǐ-ŭs dŏm-ĭn'-ĭ-cŭs. [Ch- hard; second syllable long.] Gr. χαράδριος, some kind of a bird, supposed to be a plover, and the same as τρόχιλος; from χαράδρα, the watery places inhabited by such birds. As used by Aristotle, the word apparently refers to Oedicnemus crepitans. Lat. dominicus, see Dendraca, No. 129.

This stands as *C. fulvus* var. *virginicus* in the orig. ed., but Müller's name has priority over Gmelin's. See Ridg., Pr. Nat. Mus., ii, 1880, p. 9; and Cassin, Pr. Phila. Acad., 1864, p. 246.

- 582. C. d. fül'-vŭs. Lat. fulvus, fulvous, yellow. Not in the orig. ed. Since discovered in Alaska. See Cours, in Elliot's Prybilov Report, 1875, 179; and Birds N. W., 1874, p. 450, note.
- 583. C. pluv-i-ā/-līs. Lat. pluvialis, rainy, pertaining to rain, bringing rain; pluvia, rain; pluo, to rain: the bird was supposed in some way related to rain or the rainy season: "plover" is the same.

Not in the orig. ed.; ascertained to occur in Greenland; see Newt., Man. N. H. Greenl., 1875, p. 101; Freke, Zoölogist, September, 1881, p. 374.

- .584. Ægialites vociferus (L.) Cass. B 504. C 397. R 516. Kildeer Ring Plover.
- .585. Ægialites wilsonius (Ord) Cass. B 506. C 398. R 522. Wilson's Ring Plover.
- 586. Ægialites semipalmatus (Bp.) Cab. B 507. C 399. R 517. Semipalmated Ring Plover; Ring-neck.
- 587. Ægialites melodus (Ord) Cab. B 508. C 400, 400a. R 520. Piping Ring Plover; Ring-neck.
  - 588. Ægialites melodus circumcinctus Ridg. B —. C 400a. R 520a. (?)
    Belted Piping Plover.
  - 589. Ægialites hiaticula (L.) Boie. B —. C —. R 518. European Ring Plover.
  - 590. Ægialites curonicus (Gm.) Gray. B —. C 400bis. R 519. European Lesser Ring Plover.
  - 591. Ægialites cantianus nivosus (Cass.) Coues. B 509. C 401. R 521. Snowy Ring Plover.
  - 534. Aēg-ĭ-ă'-lĭ-tēs vō-cĭ'-fĕr-ŭs. Gr. alγιαλίτης, masculine, or alγιαλίτις, feminine, or alγιαλός, an inhabitant of the seashore; alγιαλός, the coast, from the breaking of the waves upon it (ἄγννμι). The name is very appropriate to these beach-birds. Both forms, ægialites, masculine, and ægialitis, feminine, are in common use; either is perfectly correct; but as Boie wrote ægialites originally, this form should be preserved.—Lat. vociferus, vociferous; vox, genitive vocis, voice, and fero, I bear; vox digammated from ωψ.
  - 585. A. wĭl-sŏn'-ĭ-ŭs. To Alexander Wilson.
  - 536. A. sĕm-ĭ-pāl-mā'-tŭs. Lat. semi, half; sibilated from Gr. ἡμι, hemi-, a contraction of ἥμισυs, half, and palmatus, palmated, web-footed; palma, the palm of the hand, the hand itself; from Gr. παλάμη, of same meaning. The bird is conspicuously webbed between the toes, in comparison with its allies.
  - 587. **A.** měl-5'-dŭs. Lat. melodus, Gr. μελφδόs, melodious, sweetly singing; μέλοs, melody, and ἀδή, a song, an ode. (Notice the long o, being in place of the Gr. omega with iota subscript.)
  - 588. A. m. cīr-cūm-cīnc'-tŭs. Lat. circum, around; cinctus, belted, girded. See Parus, No. 52. The black is said to form a complete necklace.
  - 589. A. hī-ā-tī'-cŭ-lā. Of this word we can give no satisfactory account. It is "classic" in ornithology, going back for over two centuries; in form, it is a diminutive of hiatus, from hio, I yawn, gape.

Not in the orig. ed. Since ascertained to inhabit Continental North America, as well as long known in Greenland. See Brewer, Bull. Nutt. Club, iii, 1878, p. 49 seq.

- 590. A. cū-rŏn'-i-cŭs. Lat. Curonicus, Curonian, of the region formerly called Curonia.

  The bird described as £g. microrhynchus, Ridg., Am. Nat., viii, 1874, p. 109, has since been identified with the above. See Pr. Nat. Mus., ii, 1880, p. 10; 1881, p. 67. The bird is very questionably North American.
- 591. A. cān-tǐ-ā'-nŭs nǐv-ō'-sŭs. Lat. Cantianus, Kentish. Lat. nivosus, snowy, in allusion to the color; nix, genitive nivis, snow; Gr. νίψ, νιφόs, snow.

- 592. Podasocys montanus (Towns.) Coues. B 505. C 402. R 523.

  Mountain Plover.
- 593. Vanellus cristatus Meyer. B —. C —. R 512. (G.) Lapwing.
- 594. Aphriza virgata (Gm.) Gray. B 511. C 403. R 511. Surf Bird.
- 595. Hæmatopus ostrilegus L. B —, C —, R 506. (G.) European Oyster-catcher.
- 596. Hæmatopus palliatus Temm. B 512. C 404. R 507. American Oyster-catcher.
- 597. Hæmatopus niger Pall. B 513. C 405. R 508. Black Oyster-catcher.
- 598. Strepsilas interpres (L.) Ill. B 515. C 406. R 509. Turnstone.
- 599. Strepsilas interpres melanocephalus (Vig.) Coues. B 516. C 406a. R 510. Black-headed Turnstone.
- 592. Pŏd-ăs-ō'-cys mōn-tā'-nŭs. The word *Podasocys* is simply the transliteration of the familiar Homeric epithet of Achilles, "swift as to his feet" πόδας ἀκὺς 'Αχιλλεύς. Lat. montanus, pertaining to mountains.
- 593. Vā-nēl'-lūs crīs-tā'-tūs. Lat. vanus, empty, void, vain, whence vanellus, as a diminutive, for the restless, idle, and noisy bird. "In the spring the wanton lapwing gets himself another crest." (Tennyson.) Lat. cristatus, crested.

  Not in the orig. ed. Only North American as occurring in Greenland. See Reinh.,
- 594. Aph-rī'-ză vīr-gā'-tă. Gr. ἀφρόs, surf, sca-foam, and ζάω, I live; badly formed, but euphonious. Compare Aphrodite, the Greek Venus, foam-formed. Audubon, who invented the word, gives the above etymology; but Wharton's MS. suggests more direct derivation from ἀφρίζω, I foam. Lat. virgata, striped, streaked; virga, a rod, green sprout, osier; from vireo, I am green.
- 595. Haēm-at'-ŏ-pūs ōs-trĭ'-lĕ-gŭs. Gr. aiματοποῦs, red-footed; aîμα, genitive a'lματοs, blood, and ποῦs, foot. The word is commonly but wrongly accented on the penult; but that would be aiματωπόs, meaning red-eyed.—Lat. astrea, an oyster, and lego, I collect, gather. Commonly written ostralegus; but the above seems to be the correct form, agreeable with frugilegus, for example, and conformable with the actual word ostriferus in the following lines:—

Quàm quibus in patriam ventosa per æquora vectis, Pontus et *ostriferi* fauces tentantur Abydi. — Verg., Georg., i, 206, 207.

Not in orig. ed. Only North American as occurring in Greenland. See Ibis, 1861, p. 9.

- 596. H. pāl-lī-ā'-tŭs. Lat. palliatus, wearing the pallium, a kind of cloak; to "palliate" is literally to hide, cover up as with a cloak. The allusion here is to the particular coloration of the bird. See Contopus, No. 380.
- 597. H. nig'-ĕr. Lat. niger, black.

Ibis, 1861, p. 9.

- 598. Strēp'-sĭ-lās īn-tēr'-prēs. Gr. στρέφω, future στρέψω, I turn; στρέψις, a turning over; and λâς, a stone; literally "turn-stone." Lat. interpres, a go-between, factor, broker, agent; literally, an interpreter, that is, inter-prætor; prætor, a Roman magistrate, from præ and εο, I go before.
- 599. S. i. měl-ăn-ŏ-ceph'-ăl-ŭs. Gr. μέλας, genitive μέλανος, black, and κεφαλή, head.

- 600. Recurvirostra americana Gm. B 517. C 407. R 566.

  American Avocet.
- 601. Himantopus mexicanus (Müll.) Ord. B 518. C 408. R 567. Black-necked Stilt.
- 602. Steganopus wilsoni (Sab.) Coues. B 519. C 409. R 565. Wilson's Phalarope.
- 603. Lobipes hyperboreus (L.) Cuv. B 520. C 410. R 564.
  Northern Phalarope; Red-necked Phalarope.
- 604. Phalaropus fulicarius (L.) Bp. B 521. C 411. R 563. Red Phalarope; Gray Phalarope.
- 605. Philohela minor (Gm.) Gr. B 522. C 412. R 525. American Woodcock.
- 606. Scolopax rusticula L. B —, C 413. R 524. (! E.) European Woodcock,
- 600. Rě-cūr-vǐ-rōs'-tră ăm-ĕr-I-cā'-nă. Lat. recurvus, bent upward, recurved, and rostrum, beak: as the bill of the avocet notably is. The English word is either avocet or avoset, the meaning of which we know not.
- 601. Him-ān'-tŏ-pūs mēx-ī-cā'-nūs. Gr. ἱμαντόπουs, Lat. himantopus, the stilt, from ἱμάς, genitive ἱμάντος, and ποῦς, foot. The former word means a thong or strap; applied to this bird on account of its very long leathery legs like straps. Commonly accented on the penult; see Contopus, No. 380.

This stands as H. nigricollis of the orig. ed.; see Cassin, Pr. Phila. Acad., 1864, p. 246.

- 602. Stěg-ăn'-ŏ-pūs wil'-sŏn-ī. Gr. στεγανόπουs, web-footed; στεγανόs, webbed; στεγάνη, a web; στέγω, I cover, roof in, and ποῦs, foot. Commonly accented on the penult; see Contopus, No. 380.
- 603. Löb'-ĭ-pēs hy-pēr-bŏr'-ĕ-ūs. Gr. λοβόs, Lat. lobus, a lobe, flap, and Lat. pēs, foot; "lobe-foot," in allusion to the flaps on the toes.—Lat. hyperboreus, Gr. ὑπερβόρεοs, hyperborean, in the extreme north, "beyond the north wind," in the sense of where the north wind comes from.
- 604. Phăl-ār'-ŏ-pūs fŭl-ĭ-cā'-rĭ-ŭs. Gr. φαλαρίs, the coot, so called from the conspicuous white of the bill, φαλαρόs meaning white, bright, clear, &c.; and ποῦs, foot; phalaropus is "coot-foot;" the phalarope was early called "coot-footed tringa," from the flaps on the toes, like those of a coot. The full form of the word would be phalaridopus. Lat. fulicarius, relating to a coot; the specific name being derived, like the generic, from the lobate feet. See also Fulica, No. 686. See Contopus, No. 380.
- 605. Phil-ŏ'-hĕl-ă min'-ōr. Gr. φίλος, loving, or a lover, and εκος, a swamp. Commonly accented on a wrongly lengthened penult.—Lat. minor, comparative degree of parvus, smaller (than the European woodcock).
- 606. Scŏl'-ŏ-pāx rūs-tī'-cũ-lã. Gr. σκολόπαξ, Lat. scolopax, a snipe; the name of this very species. The dictionaries give it as a theme, and any possible derivation is open to conjecture. cf. σκόλοψ, from the shape of the bill (most likely); σκώληξ, a worm; σκάλλω, I scratch. Lat. rusticus, a rustic, a countryman; diminutive rusticulus; from rus, the country, as opposed to the city. The word occurs as rusticola in Linnæus, and has so almost universally been written; but as Wharton shows (Ibis, 1879, p. 453), this is erroneous. The word would be ruricola, if from rus and colo, I inhabit. Rusticula is good Latin, and the epithet of "little countryman" is very appropriate to the bird.

- 607. Gallinago media Leach. B —. C —. R 526. (G.) European Snipe.
- 608. Gallinago wilsoni (Temm.) Bp. B 523. C 414. R 526a.

  American Snipe; Wilson's Snipe.
- 609. Macrorhamphus griseus (Gm.) Leach. B 524. C 415. R 527. Red-breasted Snipe; Gray-back Snipe; Dowitcher.
- 610. Macrorhamphus griseus scolopaceus (Say) Coues. B 525. C 415a. R 527a. Western Red-breasted Snipe.
- 611. Micropalama himantopus (Bp.) Bd. B 536. C 416. R 528. Stilt Sandpiper.
- 612. Ereunetes pusillus (L.) Cass. B 535. C 417. R 541. Semipalmated Sandpiper.
- 613. Ereunetes pusillus occidentalis (Lawr.) Coues. B —. C 417a. R 541a. (?)
  Western Semipalmated Sandpiper.
- 614. Actodromas minutilla (V.) Coues. B 532. C 418. R 538. Least Sandpiper.
- 615. Actodromas bairdi Coues. B —. C 419. R 537. Baird's Sandpiper.
- 607. Gāl-līn-ā/-gō mĕd'-ĭ-ă. Lat. gallus, a cock, gallina, a hen, gallinula, a chicken, gallinarius or gallinaceus, relating to poultry; the present word is an arbitrary derivative, as a Latin word, though the forms gallinago, gallinazo, and others are found in different languages. It is formed from gallina like fringillago from fringilla, or like virago from vir.—Lat. medius, median, medium, in the middle (in size, between certain other species). Not in the orig. ed.; only North American as occurring in Greenland.
- 608. G. wil'-son-i. To Alexander Wilson.
- 609. Māc-rŏ-rhām'-phūs grīs'-ĕ-ŭs. Gr. μακρόs, great, large, long; and βάμφοs, beak, bill. Notice that the β is aspirated, requiring to be followed by h, as many writers forget. Griseus, gray, grisly, grizzly; not classic; a late Latinizing of an Anglo-Saxon word; compare Fr. gris and Gr. γραῦς οι γρηῦς, γεραιός, γέρας οι γῦρας all these relate to age, when people grow gray. The word "grous" or "grouse," "the gray bird," may be related. See Leucosticte, No. 205.
- 610. M. g. scŏl-ŏ-pā'-cĕ-ŭs. The word is formed as an adjective from scolopax, which see, No. 606; scolopaceous, scolopacine, snipe-like.
- 611. Mīc-rŏ-păl'-ă-mă hĭm-ān'-tŏ-pūs. Gr. μικρόs, small, and παλάμη, the palm, the hand; same as the Lat. palma; referring to the webbing between the toes. Himantopus, see No. 601.
- 612. E-reū-nē'-tēs pŭs-īl'-lŭs. Gr. ἐρευνητήs, a searcher; from the way in which the bird probes with its bill.— Lat. pusillus, puerile; see Sitta, No. 60.
- 613. E. p. ōc-cĭ-dēn-tā'-lĭs. Lat. occidentalis, western. See Dendræca, No. 113.
- 614. Ac-tŏ'-drŏm-ăs mĭn-ū-tīl'-lã. Gr. ἀκτή, the seashore; from ἄγννμι, ἄγω, I break, as the waves do there; δρομάς, rapidly running see Ammodramus, No. 238, and Eudromias, No. 591. Lat. minutus, small, minute, diminutive, of which minutilla is an arbitrary diminutive; minuo, I lessen, diminish; it ought to have been minutula.
- 615. A. baīr'dī. To S. F. Baird.

- .616. Actodromas maculata (V.) Coues. B 531. C 420. R 534. Pectoral Sandpiper.
- 617. Actodromas bonapartii (Schl.) Coues. B 533. C 421. R 536. White-rumped Sandpiper.
- 618. Actodromas cooperi (Bd.) Coues. B 527. C 422. R 535. (?)
  Cooper's Sandpiper.
- 619. Actodromas acuminata (Horsf.) Ridg. B —. C —. R 533. (! A.)
  Sharp-tailed Sandpiper.
- 620. Arquatella maritima (Brünn.) Bd. B 528. C 423. R 530.

  Purple Sandpiper.
  - 621. Arquatella couesi Ridg. B —. C —. R 531.
    Aleutian Sandpiper.
  - 622. Arquatella ptilocnemis (Coues) Ridg. B —. C 426bis. R 532. Prybilov Sandpiper.
  - 623. Pelidna alpina (L.) Boie. B —. C —. R 539. (G.) European Dunlin.
  - 616. A. mă-cŭl-ā'-tă. Lat. maculatus, spotted; macula, a spot.
  - 617. A. bō-nă-pār'-tǐ-ī. To Charles Lucien Bonaparte, Prince of Musignano and Canino.
  - 618. A. coop'-ĕr-ī. To William Cooper, Esq. Only one specimen known.
  - 619. A. ăc-ū-mĭ-nā'-tă. Lat. acuminata, acuminate, sharpened, from acumino; like aculeata from aculeus. See Sitta, No. 58.

    Not in the orige ed. Since observed at St. Michael's Alaska. See Pr. Nat. Mus.

Not in the orig. ed. Since observed at St. Michael's, Alaska. See Pr. Nat. Mus., iii, 1880, p. 222.

- 620. Ar-quă-tēl'-lă măr-ĭt'-ĭ-mă. Arquatella, for arcuatula, is an arbitrary diminutive of arquatus, bent, bowed: this is poor Latin for arcuatus, curved, arcuate; arcuo, I bend; arcus, a bow, an arc. It refers to the slightly curved bill.—Lat. maritimus, maritime; mare, the sea.
- 621. A. m. couēs'-ī. To Dr. Elliott Coues, U. S. A. The name of this person is Norman-French, and is still not infrequently found in the north of France, pronounced in two syllables, with the grave accent on the last: Cou-ès Coo-ayz. On the removal of his ancestors to the Isle of Wight, the pronunciation naturally became corrupted into Cowz.

  The original spelling, though sometimes changed to Cowes, has been preserved in the family, no grown male members of which are known to be living in the United States excepting the person here in mention and his brother, Dr. S. F. Coues, U. S. N. The meaning of the word is unknown to us.

Not in the orig. ed. Since described, from Alaska. Bull. Nutt. Club, v, 1880, p. 160.

622. A. m. ptīl-ōc-nē'-mīs. Gr.  $\pi\tau i\lambda o\nu$ , a feather, and  $\kappa\nu\eta\mu is$ , a greave, boot; the crus being feathered to the heel.

This is the *Tringa crassirostris* of the orig. ed., very wrongly so named; also, it is *T. gracilis*, Harting. See Coues, Elliott's Prybilov Islands, 1875.

623. Pĕl-īd'-nă āl-pī'-nă. ? Gr. πελιδνόs, gray; from πέλοs, some dark color. — Lat. Alpina, Alpine; Alpes, Alps. See Eremophila, No. 82.

Not in the orig. ed. of the Check List. Only North American as occurring in Greenland. See Newton, Man. Nat. Hist. Greenland, 1875, p. 103, where the Dunlin of Greenland is recognized as distinct from var. americana.

- 624. Pelidna alpina americana (Cass.) Allen. B 530. C 424. R 539α.

  American Dunlin.
- 625. Ancylochilus subarquatus (Güld.) Kaup. B 529. C 425. R 540. Curlew Sandpiper.
- . 626. Tringa canutus L. B 526. C 426. R 529.

  Red-breasted Sandpiper; Robin Snipe; Knot. [See Addenda, No. 884.
- 627. Calidris arenaria (L.) Ill. B 534. C 427. R 542. Sanderling.
- 628. Limosa fœda (L.) Ord. B 547. C 428. R 543. Great Marbled Godwit.
- 629. Limosa hæmastica (L., 1758) Coues. B 548. C 429. R 545. Hudsonian Godwit.
  - 624. P. a. ăm-ĕr-ĭ-cā/-nă. See Parula, No. 93.
- 625. An-cy-lŏ-chī'-lŭs sŭb-ār-quā'-tŭs. Gr. ἀγκυλόχειλοs, having a curved bill: ἀγκύλοs, crooked, bent, from ἀγκών, the bent elbow, and χείλοs, the mouth, from a word signifying to open, to gape. Lat. subarquatus, slightly curved; see Arquatella, No. 620.
- 626. Trīn'-gă căn-ū'-tūs. Lat. tringa, or trynga, or tryngas, a sandpiper; not classic. Derived from Gr. τρύγγαs, an obscure and obsolete word, occurring in Aristotle as the name of some unknown bird. The species was very aptly named by Linnæus after old King Canute, who, it is said, sat on the seashore and allowed the waves to reach him, to rebuke his toadying courtiers who had declared the sea would obey his majesty, a myth according well with the habits of sandpipers. Canutus, if it has any relation with, or is of same meaning as canus, gray, hoary, πολιόs, is well suited either to the old king, or to this sandpiper in its winter dress.
- 627. Căl-id'-rĭs ă-rē-nā'-rĭ-ă. Gr. σκαλίδριs or καλίδριs, Lat. scalidris or calidris, an obscure Aristotelian bird, by some supposed to be the modern totanus calidris. The word is apparently from σκαλίs, some digging instrument, from σκάλλω, I scrape, rake, &c., and refers to the same probing habits of this sandpiper that ereunetes signalizes. But the form Chalidris also occurs, as in Belon for example; whence some refer the word to the Gr. χάλιξ, Lat. calx, calculus, &c., considering that it alludes to the pebbly or shingly beaches which the bird frequents.—Lat. arenarius, relating to sand; arena, sand, or a sandy place, as the arena was, where gladiatorial and other sports were witnessed by the Roman brutes.
- 628. Lī-mō'-să foē'-dă. Lat. limosus, miry, muddy; limus, mud, slime. We can learn nothing of any such word as fedoa, and take it to be a misprint or other mistake for fædus, -a, -um, ugly, unseemly, &c. It might be supposed to have some relation to fædus, a compact, treaty, the sense of which is seen in federal, confederate, &c., and the application of which would be to the gregariousness of the bird. But fædus, in the latter sense, is not an adjective; it is fædus, fæderis, and the adjectival form would be federatus; while there is an adjective fædus, ugly, as well as a verb fædo, to defile, the participial of which is fædatus. In view of these facts, we propose to substitute fæda for fedoa, until some satisfactory explanation of the latter can be given. Fedoa occurs at least as far back as Edwards as the name of this species, and has since passed unchallenged.
- 629. L. haēm-ās'-tĭ-că. Gr. αἰμαστικός or αἰματικός, hæmastic or hæmatic, of a bloody-red color; αἰμάσσω, I make bloody; αἶμα, blood; referring to the red under parts, so conspicuous in this species.

This stands as L. hudsonica in the orig. ed. See Coues, Bull. Nutt. Club, v, 1880, p. 100.

- 630. Limosa ægocephala (L.) Leach. B —. C —. R 546. (G.)
  Black-tailed Godwit.
- 631. Limosa uropygialis Gould. B —, C 430. R 544. (!A.)
  White-rumped Godwit.
- 632. Symphemia semipalmata (Gm.) Hartl. B 537. C 431. R 552. Semipalmated Tattler; Willet.
- 633. Totanus melanoleucus (Gm.) V. B 539. C 432. R 548. Greater Tattler; Stone Snipe.
- 634. Totanus flavipes (Gm.) V. B 540. C 433. R 549. Lesser Tattler; Yellowshanks.
- 635. Totanus glottis (L.) Bechst. B 538. C 434. R 547. (! E.) Greenshanks.
- 636. Rhyacophilus ochropus (L.) Ridg. B —. C —. R 551. (! E.) Green Sandpiper.
- 637. Rhyacophilus solitarius (Wils.) Bp. B 541. C 435. R 550. Solitary Tattler.
- 638. Tringoïdes macularius (L.) Gr. B 543. C 436. R 557 Spotted Tattler; Spotted Sandpiper.
  - 630. L. aēg-ŏ-cĕph'-ă-lă. Gr. αἰγοκέφαλος, an Aristotelian epithet of some unknown bird; it literally means "goat-headed," but what application? About the middle of the sixteenth century it was applied by Belon to a species of Limosa, perhaps from the cry of the bird being fancied like the bleating of a goat; "bleating" is a term in every-day use now to express the peculiar sounds made by some snipes. The curious English word godwit is derived by Johnson from Anglo-Saxon god, good, and wiht, animal: by others from god, and weide, game; latter not unlikely.

Not in the orig. ed. Only North American as a straggler to Greenland.

- 631. L. ū-rŏ-py-gĭ-ā'-lĭs. See Centurus, No. 452.
- 632. Sym-phē'-mī-ā sĕm-ī-pāl-mā'-ta. Gr. σύμφημι; σύν, with, and φημί, I speak; alluding to the noisy concerts of the birds. Lat. semipalmata, half-webbed: see Ægialites, No. 584. "Willet" is derived from the sound of the bird's voice; sometimes written "pilwillet."
- 633. Tŏ-tā/-nŭs mĕl-ăn-ŏ-leū/-cŭs. Totanus is Latinized from the Italian totano, a name of some bird of the kind. We suppose it should be accented on a lengthened penult. Gr. μέλαs, genitive μέλανος, black, and λευκός, white.
- 634. T. fla'-vi-pes. Lat. flavus, yellow; pes, foot.
- 635. T. glōt'-tīs. Gr. γλῶσσα or γλῶττα, the tongue; referring to the noisiness of the bird. This is given in the orig. ed. as Totanus chloropus.
- 636. R. ōch'-rŏ-pūs. Gr. ἀχρός, pale, sallow, wan, and ποῦς, foot. From this word come Lat. ochra, and our ochre, ochreous, ochraceous, as names of some dull yellowish color. Linnæus had originally ocrophus by misprint.

Not in the orig. ed. Since found in Nova Scotia as a straggler from Europe. See Bull. Nutt. Club, iii, 1878, p. 49.

- 637. Rhy-ă-cŏ'-phīl-ŭs sŏl-ī-tā'-rī-ŭs. Gr. βύαξ, genitive βύακος, a stream, brook; βέω or βύω, I flow; and φίλος, loving, loved, a lover. Lat. solitarius, solitary; solus, alone.
- 638. Trīn-gŏ-ī'-dēs măc-ŭl-ā'-rī-ŭs. See Tringa, No. 62¶ and add εἶδοs, resemblance. Note that the word is in four syllables, accented on the penult. Lat. macularius; not classic; like maculatus and maculosus, spotted; macula, a spot.

- 639. Machetes pugnax (L.) Cuv. B 544. C 437. R 554. (!E.)
  Ruff (3); Reeve (9).
- 640. Bartramia longicauda (Bechst.) Coues. B 545. C 438. R 555.

  Bartramian Tattler.
- 641. Tryngites rufescens (V.) Cab. B 546. C 439. R 556. Buff-breasted Sandpiper.
- 642. Heteroscelus incanus (Gm.) Coues. B 542. C 440. R 553. Wandering Tattler.
- 643. Numenius longirostris Wils. B 549. C 441. R 558. Long-billed Curlew.
- 644. Numenius phæopus (L.) Lath. B —, C —, R 561. (G.) European Whimbrel.
- 639. Māch-ē'-tēs pūg'-nāx. Gr. μαχητήs, a fighter, combatant, in allusion to the pugnacity of the male in the breeding season; μάχομαι, I fight; μάχη, a battle. Lat. pugnax, pugnacious, combative; pugna, I fight; pugna, a battle; properly, fisticuffs, as the primitive mode of fighting; pugnum, the fist; root pug, whence come the whole set of words, and others, as pygmy, &c.
- 640. Bār-trăm'-ĭ-ă lōn-gĭ-caūd'-ă. To William Bartram, "grandfather of American ornithology."—The usual generic name, actiturus, is from the Gr. ἀκτίτηs, a doer by the sea, a beach-inhabiter, a "longshoreman," from ἀκτή, the seashore, and οὖρα, tail.—Lat. longus, long, and cauda, tail.

This is Actiturus bartramius of the orig. ed. See Coues, Bull. Nutt. Club, v, 1880, p. 100.

- 641. Tryn'-gĭ-tēs rū-fēs'-cēns. See Tringa, No. 626. Here we have another form of the word, nearer the original Gr. τρύγγαs, with the termination -τηs, -tes; this suffix commonly denoting active agency, as the English -er, for example, makes work-er from work. Lat. rufescens, present participle of rufesco, I grow reddish.
- 642. Hět-ě-rở'-scěl-ŭs īn-cān'-ŭs. Gr. ἕτερος, opposite, different, otherwise, and σκέλος, the leg, shin; from the peculiar scutellation of the leg.—Lat. incanus, very gray, quite hoary, as the bird is: in and canus.
- 643. Nū-mē'-nī-ŭs lōn-gī-rōs'-trīs. A curious etymology is this, if the derivation assigned be true. Gr. νέοs, new, young, and μήν, a month, μήνη, the moon; the narrow arcuate bill being likened to the new crescent moon. The same word is seen in meniscus, a kind of lens, but primarily and literally a little moon. But numenius might also be derived directly from numen, a nod, a bending of the head downward and forward (hence assent, command, and hence a divinity, who nods assent or expresses its will by such gesture); Gr. νεῦμα, a nod, νεύω, I nod; very applicable to the attitude of the bird. Whichever of these derivations we approve, they amount to practically the same thing; for numenius certainly refers to the shape of the bill, being used by the ornithologists of the heroic age as synonymous with arquata or arcuata. Lat. longinostris, long-billed; longus and rostrum. "Curlew" is not an imitation of the bird's voice, but a mangling of the French name cour-lieu, "run-place," from the coursing of the birds: compare courlis, courly, courlan, cocorli, &c.
- 644. N. phaē/-ŏ-pūs. Gr. φαιός, dark colored, dusky, gray, swarthy; its exact meaning is expressed when we say "gray of the morning:" related to φαίνω, I appear; ποῦς, foot. "Whimbrel" is apparently Anglo-Saxon; related to whim, whimsical, in the sense of flighty, a gad-about.

Not in the orig. ed. Only North American as a bird of Greenland.

- 645. Numenius hudsonicus Lath. B 550, C 442, R 559, Hudsonian Curlew.
- 646. Numenius borealis (Forst.) Lath. B 551. C 443. R 560. Eskimo Curlew.
- 647. Numenius taïtensis (Gm.) Lath. B —. C 442bis. R 562. (!A.)
  Otahiti Curlew.
- 648. Tantalus loculator L. B 497. C 444. R 500.
  Wood Ibis.
- 649. Plegadis falcinellus (L.) Kaup. B 500. C 445. R 503. Glossy Ibis.
- 650. Plegadis guarauna (L.) Ridg. B —. C 445bis, 445ter. R 504. White-faced Glossy Ibis.
- 651. Eudocimus albus (L.) Wagl. B 499. C 446. R 501. White Ibis.
  - 645. N. hud-son'-i-cus. To Hudson's Bay, after Henry Hudson.
- 646. N., bor-e-a'-lis. Lat. borealis, northern; boreas, the northwind.
- 647. N. tă-ī-tēn'-sĭs. Of Otaheite, one of the Society or Friendly Islands. The original orthography, tahitiensis, is resolvable into the above, which is less barbarous in sound and look. Though named for the island called in English Otaheite, or better Otahiti, the first syllable is to be dropped as being merely the definite article the. It is the native name O-tahiti, the island; i. e., the principal island.

This is N. femoralis, Peale, of the orig. ed., Appendix.

- 648. Tān'-tǎl-ús lŏ-cŭ-lā'-tōr. Gr. Τάνταλος, Tantalus, the Phrygian king, who, admitted to the councils of the gods, betrayed their secrets, and was tormented, "tantalized," with food and water in sight but unattainable.—Lat. locus, a place; loculus, a little place, division, compartment; loculatus or loculosus, furnished with compartments, full of "pigeon-holes"; but qu. loculator and its application to this bird?
- 649. Plē'-gă-dĭs fāl-cĭn-ēl'-lŭs. Gr. πληγάs, a scythe, sickle, from πλήσσω or πλήττω, I strike. The actual form, Plegadis, may be a diminutive; if so, it is exactly Greek for the quasi-Latin falcinellus, falcicula, or falcunculus, a little scythe, small hook; falx, a reaping-hook or any thing of that falcate shape, as the bill of this bird is. See Falco, No. 498.

This stands in the orig. ed. as *Ibis falcinellus* var. ordii. But it has proved to be not satisfactorily distinguished from the European form; while as to the generic designation, see Ibis, 1878, p. 112.

650. P. gŭ-ă-raū'-nă. A barbarous word, of some South American (Brazilian) dialect. It occurs as such in Marcgrave and other early ornithologists.

This stands as *Ibis guarauna* in the orig. ed.; see No. 649. The *Ibis thalassinus* of Ridg., Am. Nat., viii, 1874, p. 110, inserted in the Appendix of the orig. ed. as No. 445 ter, proves to be the young of this species: see Coues, Bull. U. S. Geol. and Geogr. Surv. Terr., iv, No. 1, 1878, p. 57.

651. Eū-dŏc'-ĭ-mŭs āl'-būs. Gr. εὐδόκιμος, well-tried; hence, approved, famous, of high repute: from εὐ, well, and δόκιμος, assayed and found acceptable; δέχομαι, I accept. The ibis or τβις of the ancients (not this species) was a celebrated and sacred bird; it was the Egyptian bird, now called Ibis œthiopica. — Lat. albus, white.

This is Ibis alba in the orig. ed. See Elliot, Ibis, 1877, p. 482.

- 652. Eudocimus ruber (L.) Wagl. B 498, C 447, R 502. Scarlet Ibis.
- 653. Ajaja rosea (Briss.) Reich. B 501. C 448. R 505. Roseate Spoonbill.
- 654. Mycteria americana L. B —. C 448bis. R 499. (! M.)
  American Jabiru.
- 655. Ardea herodias L. B 487. C 449. R 487. Great Blue Heron.
- 656. Ardea occidentalis Aud. B 488, 489. C 450, 451. R 486. Great White Heron; Florida Heron.
- 657. Ardea cinerea L. B C R 488. (G.) European Blue Heron.
- 658. Herodias egretta (Gm.) Gr. B 486, 486\*. C 452. R 489. Great White Egret.
- 659. Garzetta candidissima (Gm.) Bp. B 485. C 453. R 490.
  Little White Egret; Snowy Heron.
- 660. Hydranassa tricolor (Müll.) Ridg. B 484. C 454. R 492.
  Louisiana Heron.
- 652. E. rub'-er. Lat. ruber, red. This is Ibis rubra in the orig. ed.
- 653. Ajaja rō'-sĕ-ĕ. Lat. roseus or rosaceus, rosy, rose-red; rosa, a rose; related to Gr. βόδον; see for instance in rhodocolpus, rose-breasted. Ajaja or ajaia or ajaja is the old Brazilian name of this bird, of signification and pronunciation alike unknown to us.

  This stands as Platalea ajaja in the orig. ed.; for the change of this long-standing name, see Ridg., Pr. Nat. Mus., iii, 1880, p. 10.
- 654. Myc-tē'-rī-ā ăm-ĕr-ĭ-cā'-nā. Gr. μυκτήρ, the nose, sncut; μυκτηρίζω, literally, "I work the nose," i. e., turn up the nose at, sneer, scorn, deride, &c.; well applied to the expression of this ugly bird.
- 655. Ar'-dĕ-ă hĕr-ō'-dĭ-ăs. Lat. ardea, a heron. Gr. ἐρώδιας, ἐρωδάς, or ἐρώδιος, a heron. There is also a proper name Herodias.
- 656. A. ōc-cĭ-dēn-tā'-lĭs. See Dendraca, No. 113.

  Note. The Ardea wurdemanni of the orig. ed. is a dichroism of this species. See Ridg., Bull. U. S. Geol. Surv. Terr., iv, No. 1, 1878, p. 227.
- 657. A. cĭn-ĕr'-ĕ-ă. Lat. cinereus, ashy. See Harporhynchus, No. 22. Not in the orig. ed. Only North American as occurring in Greenland. See Reinh., Ibis, 1861, p. 9.
- 658. Hěr-ō'-dĭ-ăs ē-grēt'-tă. Latin proper name *Herodias:* see *Ardea*, No. 655. *Egretta* is Latinized from the French *aigrette*, a top-knot, plume; whence also *egret*. These words are said to be related to heron itself, all springing from O. H. G. *hiegro*, a heron.
- 659. Gār-zēt'-tă cān-dĭ-dīs'-sĭ-mă. Garzetta is the Italian name of the corresponding European species. Lat. candidissima, very white, entirely white; superlative of candidus. See Falco, No. 501.
- 660. Hyd-ră-nās'-să trĭ'-cŏl-ŏr. Gr. ὕδωρ, water, giving in Latin hydra-, and νᾶσσα or νῆσσα, a water-fowl; from a verb meaning to swim. We have here two words very fruitful of derivatives; one giving us the compounds of hydr-, as hydrawlic, the other those relating to the sea, a ship, or swimming: nautical, aeronaut, navy, navigate, nausea; the latter is originally "sea"-sickness, and literally "ship"-sickness. Lat. tricolor, three-colored.

This stands as Ardea leucogastra var. leucoprymna in the orig. ed. See Ridg., Bull. U. S. Geol. Surv. Terr., iv, No. 1, 1878, p. 224.

- 661. Dichromanassa rufa (Bodd.) Ridg. B 482, 483. C 455. R 491. Reddish Egret.
- 662. Florida cœrulea (L.) Bd. B 490. C 456. R 490. Little Blue Heron.
- 663. Butorides virescens (L.) Cab. B 493. C 457. R 494.
  Green Heron.
- ,664. Nyctiardea grisea nævia (Bodd.) Allen. B 495. C 458. R 495. American Night Heron.
- 665. Nycterodius violaceus (L.) Reich. B 496. C 459. R 496. Yellow-crowned Night Heron.
- 666. Botaurus mugitans (Bartr.) Coues. \*B 492. C 460. R 497.

  American Bittern.
- 667. Ardetta exilis (Gm.) Gr. B 491. C 461. R 498. Least Bittern.
  - 661. Dī-chrō-mā-nās'-să rū'-fă. Gr. δίs, twice; χρω̂μα, chroma, color; originally, probably, flesh-color; and νᾶσσα, a water-fowl; alluding to the dichroism or dichromatism which prevails in this and other herons, these birds of the same species being found either pure white or variously colored. Lat. rufus, reddish.

This stands as Ardea rufa in the orig. ed. See Ridg., Bull. U. S. Geol. and Geogr. Surv. Terr., iv, No. 1, 1878, p. 246.

- 662. Flō'-rĭ-dă coē-rŭl-ĕ-ă. Lat. floridus, florid, flowery; flos, a flower; but the genus is named for the State of Florida. Lat. cæruleus, blue; see Polioptila, No. 36.
- 663. Bū-tōr-ī'-dēs vĭr-ēs'-cēns. Lat. butio or butor, a bittern; equal to bo-taur, bo-taurus, bostaurus? see Bubo, No. 462; είδοs, resemblance. There is also a proper name Butorides—Lat. virescens, present participle of viresco, I grow green, am greenish, from vireo, which see, No. 170.
- 664. Nyc-tĭ-ār'-dĕ-ă grīs'-ĕ-ă naē'-vĭ-ā. Badly formed from Gr. νύξ, gen. νυκτόs, night, and Lat. ardea, a heron; better Noctiardea, like Noctiluca, &c. Lat. griseus, see Macrorhamphus, No. 609, and Leucosticte, No. 205. Lat. nævius, see Turdus, No. 5.
- 665. Nyc-tĕr-ō'-dī-ŭs vī-ŏ-lā'-cĕ-ŭs. Gr. νύξ, night, and ἐρωδιόs, a heron, like the Latin ardea. Commonly written nyctherodius; but we see no occasion for the h, the ε not being aspirated; though the h is seen in the Lat. herodias. Lat. violaceus, violet-colored; viola, a violet, pansy.
- 636. Bō-taū'-rūs mū-gī'-tāns. The many words bittern, bitorne, bitore, butor, butio, are all onomatopæic, from the hollow guttural sound of the bird's voice, and are referable to bostaurus or bo-taurus? see Bubo, No. 462. Lat. mugitans, bellowing; mugito, I low like a cow; as the children say, "moo."
- 667. Ar-dēt'-tă ēx-ī'-līs. Ardetta is an Italian word, equivalent to ardeola, diminutive of ardea.

   Lat. exilis, contracted from exigilis, equivalent to exiguus, from exigo, this equal to ex and ago, literally, I drive out. Any thing exacted or exact, is carefully measured, considered, strictly accounted for; hence likely to be scanty, as opposed to abundant, or superfluous; therefore, poor, thin, mean, small; any of these latter adjectives well suited to this lean little bird. We have the idea in several applications in the English words exigency, an emergency; exiguous, small; the French exigeant, exacting; and in our rare though actual word exile, small. (The latter must not be confounded, however, with exile, banishment, one banished; though this might seem exactly from exigo, "I drive out," it is from another root: exsulo, exsul.)

- 668. Grus americana (L.) Temm. B 478. C 462. R 582. White Crane; Whooping Crane.
- 669. Grus canadensis (L.) Temm. B 480. C 463. R 584.
  Northern Sandhill Crane.
- .670. Grus pratensis Bartr. B 479. C —. R 583. Southern Sandhill Crane.
- 671. Aramus pictus (Bartr.) Coues. B 481. C 464. R 581. Scolopaceous Courlan; Limpkin.
- 672. Parra gymnostoma Wagl. B —. C —. R 568. (! M.) Mexican Jaçaná.
- . 673. Rallus longirostris crepitans (Gm.) Ridg. B 553. C 465. R 571. Clapper Rail; Salt Marsh Hen.
  - 668. Grus ăm-er-i-cā'-nă. Lat. grus, genitive gruis, feminine noun of the third declension, a crane. The word refers to the hollow guttural voice of the birds, and is apparently related to English grunt.
  - 669. G. căn-ă-dēn'-sīs. It was doubtless upon the northern bird, figured by Edwards, that Linnæus based this name. G. fraterculus of Cassin has been found distinct from the common sandhill crane of the United States, and identical with the northern bird. It is therefore properly a synonym of canadensis, and another name must be found for the United States bird commonly called canadensis. See next species. See Ridg., Bull. Nutt. Club, v, 1880, p. 187; Coues, ibid., p. 188.
  - 670. G. prā-tēn'-sīs. Lat. pratensis, relating to pratum, a field.

    Not in the orig. ed. See last species.
  - 671. Ar'-ă-mus pīc'-tus. The word aramus is unknown to us. Agassiz gives it as "nom. propr." A correspondent remarks: "Vieillot's Analyse is very incorrectly printed, and some letter may have been omitted or changed; hence the clue is still to seek. The origin seems hopeless, unless revealed by accident." Under these circumstances, it is consoling to reflect that the word is more decorous in form than many of known classic derivation. Lat. pictus, see Setophaga, No. 151.
  - 672. Pār'-rǎ gȳm-nō'-stŏ-mǎ. Parra is a good Latin word, being the name of some unknown bird regarded as of ill-omen; as occurring in Pliny, said to be the European Lapwing, Vanellus cristatus. Transferred by Linnæus to a mixed lot of spur-winged birds, chiefly of America. "Jaçaná" is the Brazilian name of a species of this genus; made a generic term by Brisson in 1760, and we do not see why it should not be employed instead of Parra. Gr. γυμνός, naked, and στόμα, mouth; in allusion to the caruncular skin at the base of the bill.

Not in the orig. ed.; since discovered in Texas by J. C. Merrill: see Bull. Nutt. Club, i, 1876, p. 88; Pr. U. S. Nat. Mus., i, 1878, p. 167.

673. Rāl'-lŭs lōn-gĭ-rōs'-trĭs crĕp'-ĭ-tāns. Rallus is said to be contracted from rarulus, a diminutive of rarus, rare; and to mean thin, slight; if so, the adjective has become an apt generic name for these lean narrow birds. It is more likely, however, to be onomatopœic, Latinized in late days from the French rasle, râle, a rattling cry, Engl. rail, to reproach, deride, &c., having nothing to do with the English rail (of a fence); very applicable to these clamorous birds. — Lat. longirostris, long-billed. — Lat. crepitans, present participle of crepito, I creak, crackle, clatter, crepitate; a frequentative or intensive form of crepo, of same signification.

This is R. longirostris of the orig. ed.

- 674. Rallus longirostris obsoletus (Ridg.) Coues. B —. C 466a. R 570. California Clapper Rail.
- 675. Rallus longirostris saturatus Hensh. B —, C —, R 571a.

  Louisiana Clapper Rail.
- 676. Rallus elegans Aud. B 552. C 466. R 569. King Rail; Fresh Marsh Hen.
- 677. Rallus virginianus L. B 554. C 467. R 572. Virginia Rail.
- 678. Porzana maruetta (Leach) Bp. B C R 573. (G.)
  Spotted Crake.
- 679. Porzana carolina (L.) V. B 555. C 468. R 574. Carolina Crake; Rail; Sora; Ortolan.
- 680. Porzana noveboracensis (Gm.) Cass. B 557. C 469. R 575. Yellow Crake.
- 681. Porzana jamaïcensis (Gm.) Cass. B 556. C 470. R 576. Black Crake.
  - 674. R. 1. ōb-sŏ1-ē'-tŭs. Lat. obsoletus, obsolete, grown unaccustomed, passed out of vogue; ob, opposition, and solesco, I grow accustomed; soleo, I am accustomed. The application is to the faded, as if worn out and disused, coloration.

This is R. elegans var. obsoletus, of the orig. ed., Appendix: see Bull. Nutt. Club, v, 1880, p. 139.

1000, p. 199. 175 R 1 săt-ŭ-rā/-ti

- 675. R. l. săt-ŭ-rā'-tŭs. Lat. saturatus, saturated, satiated, filled full; i. e., having eaten enough; satis, enough: whence satisfied, &c. The allusion is to the color, which is full, i. e. rich, dark, heavy.

  Not in the orig. ed.; since described. See Bull. Nutt. Club, v, 1880, p. 140.
- 676. R. &'-18-gāns. Lat. elegans or eligans, elegant; literally, choice, select; from e and lego, I pick out; quite equivalent to electus, chosen, picked, eclectic, &c.
- 677. R. vīr-gĭn-j-ā'-nŭs. To Virginia, "mother of Presidents," and wet-nurse of Secession.
- 678. Pōr-zā'-nă mă-rŭ-ēt'-tă. Porzana is an Italian word, the meaning of which we know not; it has been in book-use for several centuries, as the name of some marsh bird. Maruetta is likewise Italian: said to be applicable to anything by the sea, and hence to be equivalent to maritime. Crake is to crackle, cackle, creak, croak, quack, &c.; see Crex, No. 683, Querquedula, No. 714.

Not in the orig. ed. Only North American as occurring in Greenland. See Reinh., Ibis, 1861, p. 12.

- 679. P. că-rŏ-lī'-nă. To Carolina. This is the rail of sportsmen. It is also called sora or soree; why, we know not: the word is colloquial and local, and has scarcely crept into the books. The word "ortolan" has a curious connection with this species. It is Italian and French, equal to the Latin hortulanus, relating to a garden: the "ortolan" is Emberiza hortulana, a bunting, esteemed a great delicacy by gourmands; and our crake has been called ortolan for no better reason than that it is also edible and sapid! The same name is sometimes applied to the bobolink, Dolichonyx oryzivorus, because it is found abundantly in the same marshes in the fall, and sells in the same restaurants as the same bird as the rail, the two being brought in together by the gunners.
- 680. P. no-ve-bor-a-cen'-sis. No New York. See Vireo, No. 181.
- 681. P. jăm-ă-ĭ-cēn'-sīs. To Jamaica. The name signifies in the vernacular the island of springs, of flowing water.

- 682. Porzana jamaïcensis coturniculus Bd. B —. C 470a. R 576a. Farallone Black Crake.
- 683. Crex pratensis Bechst. B 558. C 471. R 577. (! E.)
  Corn Crake.
- 684. Gallinula galeata (Licht.) Bp. B 560. C 472. R 579. Florida Gallinule.
- 685. Ionornis martinica (L.) Reich. B 561. C 473. R 578. Purple Gallinule.
- 686. Fulica americana Gm. B 559. C 474. R 580.

  American Coot.

[See Addenda, No. 885.

- 687. Phœnicopterus ruber L. B 502. C 475. R 585. Red Flamingo.
- 688. Cygnus buccinator Rich. B 562. C 476. R 589. Trumpeter Swan.
- 689. Cygnus columbianus (Ord) Coues. B 561bis. C 477. R 588.

  American Swan.
- 682. P. j. cō-tūr-nī'-cŭ-lŭs. Lat. diminutive of Coturnix, which see, No. 579.
- 683. Crēx prā-tēn'-sīs. Gr. κρέξ, Lat. crex, a crake; all three of these words are the same, meaning the creaking, crackling cry of the bird; κρέκκω, I make such a noise.— Lat. pratensis, see Grus, No. 670. (A subgenus, "Crescicus," which passed in some American works for the black rail, was simply a misprint for creciscus, which is a Greek diminutive form of κρέξ.)
- 684. Gāl-līn'-ŭ-lă găl-ĕ-ā'-tă. Lat. gallinula, a diminutive of gallina, a hen: see Gallinago, No. 608. It is commonly but wrongly accented on the penult, and pronounced gally-new'-ler!

  But gahl-leen'-u-lah is doubtless nearer the sound a Roman would have made if he had used the word. Lat. galeata, helmeted; galea, a helmet; galeo, I crown with a helmet; very apt, in allusion to the frontal shield of a bird of this genus.
- 685. I-ōn-ōr'-nĭs mār-tĭn'-ĭ-că. Gr. τον, λωνία, a violet, and όρνις, a bird; well applied to these luxurious porphyritic or hyacinthine "sultans." English violet is from Lat. viola, and this is very easily gotten from the Greek. To the island of Martinique.
- 686. Fŭl'-ĭ-că ăm-ĕr-ĭ-cā'-nă. Lat. fulica, same as fulix, a coot, from the sooty color of the bird; fuligo, soot, whence fuliginosus, &c.
- 687. Phoē-nǐ-cōp/-tĕr-ŭs rŭb'-ĕr. Gr. φοινικόπτερος, Lat. phænicopterus, the flamingo; literally, red-winged: φοίνιξ and πτέρον: see Agelæus, No. 316. Lat. ruber, red. English flamingo seems to come directly through the Spanish flamenco, the name of this bird; both these, as the French flamant, are of course from the Latin flamma, flame, flery-red.
- 688. Cyg'-nŭs būc-cĭn-ā'-tŏr. Gr. κύκνος, Lat. cycnus or cygnus, a swan; famed for its dying song; also name of a person fabled to have been transmuted into the bird. The name is probably rooted in the idea of singing, this being one of the most persistent and ubiquitous myths. Lat. buccinator, a trumpeter, who uses his cheeks so much in blowing his instrument; buccina, or βυκάνη, a trumpet; bucca, the cheek.
- 689. C. cŏ-lūm-bĭ-ā'-nŭs. Of the Columbia River, where specimens were noted by Lewis and Clarke, afterwards named by Ord.

This stands in the orig. ed. as C. americanus. For the change, see Coues, Bull. U. S. Geol. Surv. Terr., 2d ser., No. 6, 1876, p. 444.

- 690. Cygnus musicus Bechst. B —, C —, R 586. (G.!E.)
  Whooping Swan.
- 691. Cygnus bewicki Yarr. B —. C —. R 587. (! E.)
  Bewick's Swan.
- 692. Anser albifrons (Gm.) Bechst. B —. C —. R 593. (G.) European White-fronted Goose.
- 693. Anser albifrons gambeli (Hartl.) Coues. B 565, 566. C 478. R 593α.

  American White-fronted Goose.
- 694. Chen cœrulescens (L.) Ridg. B 564. C 479. R 590.
  Blue Goose.
- ,695. Chen hyperboreus (Pall.) Boie. B 563. C 480. R 591. Snow Goose.
- 696. Chen hyperboreus albatus (Cass.) Ridg. B —. C 480a. R 591a.

  Lesser Snow Goose.
- 697. Chen rossi (Bd.) Ridg. B —. C 481. R 592.
  Ross' Snow Goose.
- 690. C. mū'-sī-cŭs. Gr. μουσικόs, Lat. musicus, relating to a muse, any one of the Muses; hence, "music" is primarily and most properly to be predicated of high ideals in general, whether in science, letters, or art. The term musicus, however, as applied to a swan, is a lucus a non lucendo, unless a relationship between the Muses and the Graces be imagined.

Not in the orig. ed., and here admitted with doubt. Greenland only, as straggler from Europe. See Reinh., Ibis, 1861, p. 13, and Freke, Zoölogist, September, 1881, p. 372. See next species.

691. C. be'-wick-i. To Thomas Bewick.

Not in the orig. ed., and here doubtfully admitted. See Pr. Nat. Mus., iii, 1880, p. 222, where Ridgway revives the record given in Fn. Bor.-Am. ii, 1831, p. 465, and states that the description of specimens killed at Igloolik, Arctic America, lat. 66°, indicates the true Bewick's Swan. But on the doubt in the case of these Arctic Swans, if any different from *C. columbianus*, see Newton, Man. Nat. Hist. Greenl., 1875, p. 113, and especially Freke, Zoölogist, September, 1881, p. 366.

692. An'sĕr āl'-bĭ-frōns. Lat. anser, a goose. How anser came about we do not know; we suppose it related more or less radically to anas, and so to νᾶσσα, a duck; see Hydranassa, No. 660. — Lat. albifrons, white forehead.

Not in the orig. ed., and here admitted with doubt, the identification of the Greenland white-fronted geese being questionable. cf. Reinh., Ibis, iii, 1861, p. 12; Newt., Man. Nat. Hist. Greenl., 1875, p. 113, and Freke, Zoölogist, September, 1881, p. —.

- 693. A. a. găm'-bĕl-lī. To William Gambel.
- 694. Chēn [pronounced cane] coē-rǔl-ēs'-cēns. Gr. χήν, a goose. See Dendraca, No. 117.
- 695. C. hy-pēr-bŏr'-ĕ-ŭs. Lat. hyperboreus, hyperborean, northern; see Lobipes, No. 603.
- 696. C. h. āl-bā'-tus. Lat. albatus, whitened, made white.
- 697. C. ros'-sī. To Bernard R. Ross, Chief Factor, H. B. Co.

- 698. Chloëphaga canagica (Sevast.) Eyt. B 573. C 482. R 598. Painted Goose.
- 699. Bernicla leucopsis (Bechst.) Boie. B 572. C 483. R 597. (! E.)
  Barnacle Goose.
- .700. Bernicla brenta (Pall.) Steph. B 570. C 484. R 595.
  Brant Goose.
- 701. Bernicla brenta nigricans (Lawr.) Coues. B 571. C —, R 596.
  Black Brant Goose.
- .702. Bernicla canadensis (L.) Boie. B 567. C 485. R 594. Canada Goose; Common Wild Goose.
- 703. Bernicla canadensis leucoparia (Brandt) Coues. B 568. C 485a. R 594b, White-cheeked Canada Goose.
- 704. Bernicla canadensis hutchinsi (Rich.) Coues. B 569. C 485b. R 594α. Hutchins' Canada Goose.
- 705. Dendrocygna fulva (Gm.) Burm. B 575. C 486. R 600. Fulvous Tree Duck.
- 706. Dendroeygna autumnalis (L.) Eyt. B 574. C 487. R 599. Autumnal Tree Duck,
- 698. Chlŏ-ē'-phă-gă că-nă'-gĭ-că. Gr. χλόα or χλόη, young grass, whence χλωρός, green; φάγω, I eat. Mr. H. W. Elliott informs us there are Eskimos of Alaska who call themselves "Kanagiamoot," i. e., "the people of the Kanag" whatever that may be; whence quasi-Lat. canagica.
- 699. Bēr'-nī-clă leū-cōp'-sĭs. Bernicla or bernicula is Latinized from the French bernicle or bernache, Engl. barnacle. We only know this word as the name of the little cirriped crustaceans out of which this goose was fabled to sprout, ripen, and fall like a fruit from its stem. A correspondent observes: "Max Müller says hibernaculum, but he gives no reason whatever (nor for hiberniculæ) founded on the word having been ever used." (cf. Lect. on the Sci. of Lang., 2d ser.) Gr. λευκός, white, and öψις, appearance.

This species is Greenlandic, but otherwise North American only as a straggler. For a résumé of occurrences, see Freke, Zoölogist, September, 1881, p. 372.

The geese of this genus stand in the orig. ed. as species of *Branta*; but that word having been found unavailable as a generic term, the name *Bernicla* is restored.

- 700. B. brēn'-tă. Latinized from brent, brant, brand, or branded goose; the forms brentus and brenthus are also found. See Campylorhynchus, No. 63. Brent or brant goose is therefore simply burnt goose, from its blackish appearance, as if charred.
- 701. B. b. nig'-ri-cans. Lat. nigricans, being blackish, like nigrescens. Not in the orig. ed.
- 702. B. că-nă-dēn'-sĭs. See Myiodioctes, No. 149.
- 703. B. c. leū-cŏ-pă-rī'-ă. Gr. λευκόs, white; παρειά, the cheek.
- 704. B. c. httch'-in-si. To ——— Hutchins, to whom we were at one time indebted for most that was known of the birds of interior British America.
- 705. Dēn-drŏ-cyg'-nă fūl'-vă. Gr. δένδρον, a tree, and κύκνος, a swan; see Cygnus, No. 688.— Lat. fulvus, fulvous, reddish.
- 706. D. aū-tūm-nā'-lĭs. Lat. autumnalis or auctumnalis, relating to the autumn, when the increase of the earth is harvested; auctumnus, the autumn; auctus, an increase, increased; auctor, a producer, author; augeo, I increase, furnish forth, augment.

- 707. Anas boscas L. B 576. C 488. R 601.
- 708. Anas obscura Gm. B 577. C 489. R 602. **Dusky Duck**.
- 709. Anas obscura fulvigula Ridg. B —. C 489a. R 603. Florida Dusky Duck.
- .710. Dafila acuta (L.) Jen. B 578. C 490. R 605. Pintail; Sprigtail.
- 711. Chaulelasmus streperus (L.) Gr. B 584. C 491. R 604. Gadwall.
- 712. Mareca penelope (L.) Selby. B 586. C 492. R 606. (! E.) European Widgeon.
- 713. Mareca americana (Gm.) Steph. B 585. C 493. R 607. American Widgeon.
- 714. Querquedula crecca (L.) Steph. B 580. C 494. R 611. (! E.) English Teal.
- 707. An'-ăs bōs'-căs. Lat. anas, a duck; doubtless related to νâσσα. See what is said under Hydranassa, No. 660. Gr. βοσκάς, Lat. boscas or boscis, a duck, probably this very species; from βόσκω, I graze. This word has almost invariably, in ornithology, been written boschas very wrongly, as Wharton was lately at pains to point out (Ibis, 1879, p. 453).
- 708. A. ōb-scū'-ră. Lat. obscurus, dark, obscure.
- 709. A. o. fūl-vī'-gŭ-lă. Lat. fulvus, fulvous, and gula, throat. This and many similar words are viciously accented on a long penult.
- 710. Dă'-fī-lă ă-cū'-tă. Dafila is a nonsense-word, invented by W. E. Leach, like  $Harelda_r$  meaning nothing. Lat. acuta, sharpened, pointed; as the tail of the bird is.
- 711. Chaū-lě-lās'-mŭs strěp'-ĕ-rŭs. Gr. χαύλιος, prominent, projecting, protuberant; and ἐλασμός, a layer, plate, lamella; referring to the denticulations of the bill.—Lat. streperus (not classic), noisy, clamorous; as we should say, obstreperous; strepitus, a noise; strepo, I make a fuss.
- 712. Mā-rē'-că pē-něl'-ŏ-pē. Mareca is said to be a Brazilian vernacular word for some kind of duck; long after, it was transferred to the widgeon. But it may also be remarked that there is the Lat. Marīca, a water-nymph. Ray has Mareca (Syn., p. 149). Penelope was the celebrated wife of Ulysses, mother of Telemachus; penelops, or in Gr. πηνέλοψ, was some kind of duck. Linnæus wrote the latter.
- 713. M. ăm-ĕr-ĭ-cā-nă. See Parula, No. 93.
- 714. Quēr-quě'-dǔ-lǔ crēc'-cǔ. Lat. querquedula, a kind of small duck; etymology obscure, and not at all to our way of thinking in the authorities consulted; apparently from καρκαίρω, κέρχω, κερκίs, κίρκη, κρέκω, κρέξ, a set of onomatopœic words formed to express a shrill or harsh creaking sound; hence related to creak, quack, crackle, &c, and quite equivalent to the very word crecca, which we have here, and which seems but an arbitrary adjective formed from κρέκω. Charleton calls one of the ducks Anas "caudacuta, The Cracke (a strepitu)." The form quacula is found in some writers; and "quack" is the usual word to express a duck's voice. See Crex, No. 683.

- 715. Querquedula carolinensis (Gm.) Steph. B 579. C 495. R 612. Green-winged Teal.
  - 716. Querquedula discors (L.) Steph. B 581. C 496. R 609.
    Blue-winged Teal.
  - 717. Querquedula cyanoptera (V.) Cass. B 582. C 497. R 610. Cinnamon Teal.
- . 718. Spatula clypeata (L.) Boie. B 583. C 498. R 608. Shoveller.
- 719. Aïx sponsa (L.) Boie. B 587. C 499. R 613.
  Summer Duck; Wood Duck.

[See Addenda, No. 886.

- 720. Fuligula marila (L.) Steph. B 588. C 500. R 614. Greater Black-head; Scaup Duck.
- 721. Fuligula affinis Eyt. B 589. C 501. R 615. Lesser Black-head; Scaup Duck.
  - 722. Fuligula collaris (Donov.) Bp. B 590. C 502. R 616. Ring-neck; Black-head.
  - 723. Fuligula ferina americana (Eyt.) Coues. B 591. C 503. R 618. American Pochard; Red-head.
  - 715. Q. că-rŏ-lĭn-ēn'-sĭs. To Carolina. The genus Nettion, in which this teal has been placed by some, is the Gr. νήττιον, a little duck; contracted from νηττάριον, a diminutive of νῆσσα or νῆττα: see Hydranassa, No. 660. Very curiously, it seems to have been used by the Greeks as a familiar term of endearment, just as we sometimes now say "little duck," or "ducky darling."
  - 716. Q. dis'-cors. Lat. discors, discordant, disagreeing, unlike; literally "two-hearted," from dis, twice, and cor, the heart; opposed to concors, concordant.
  - 717. Q. cy-ăn-ōp'-tĕ-ră. Gr. κυανός, blue, πτέρον, wing.
  - 718. Spă'-tŭ-lă clyp-ĕ-ā'-tă. Lat. spatula or spathula, Gr. σπαθίs, a spathe, spatula, spoon, ladle; with reference to the spathulous or spoon-like shape of the bird's bill. Lat. clypeatus, furnished with a shield, wearing a shield; clypeus or clipeus or clipeus or clipeum, a shield: commemorating in this case the rounded expanse of the bill.
  - 719. X-īx spōn'-să. Gr. a'\(\xi\) or ăi\(\xi\); application not obvious. Nor is the orthography settled. If the word be from the monosyllable a'\(\xi\) it should be Latinized ax; if from the dissyllable ăi\(\xi\) it becomes a\(\xi\). In the uncertainty, we do not change the accustomed form; though we suspect ax to be preferable. Lat. sponsa, a bride, a spouse, a betrothed; that is, a promised one; spondeo, I promise sacredly, I vow. Prettily applied to this lovely duck, as if the bird were arrayed for bridal.
  - 720. Fŭl-ïg'-ŭ-lā mă-rī'-lă. Lat. fuligula or fulicula, diminutive of fulica or fulix, a coot; fuligo, soot. Marila we know nothing about; qu., a proper name? qu. Gr. μαρίλη, embers, charcoal, from the scaup's pitch-black foreparts?
  - 721. F. af-fin'-is. Lat. affinis, affined, allied; ad, and finis. See Campylorhynchus, No. 64.
  - 722. F. cōl-lā'-rīs. Lat. collaris, relating to the neck, collum; this species having a ring of color, like a collar, round the neck.
  - 723. F. fĕ-rī'-nă ăm-ĕr-ĭ-cā'-nă. Lat. ferina, wild, in a state of nature, feral.

- 724. Fuligula vallisneria (Wils.) Steph. B 592. C 504. R 617. Canvas-back.
- 725. Clangula glaucium (L.) Brehm. B 593. C 505. R 620. Golden-eye.
- 726. Clangula islandica (Gm.) Bp. B 594. C 506. R 619. Barrow's Golden-eye.
- 727. Clangula albeola (L.) Steph. B 595. C 507. R 621. Buffle-head; Butter-ball; Spirit Duck.
- 728. Harelda glacialis (L.) Leach. B 597. C 508. R 623. Long-tailed Duck; Old Wife.
- 729. Camptolæmus labradorius (Gm.) Gr. B 600. C 509. R 624. Labrador Duck.
- 730. Histrionicus minutus (L.) Coues. B 596. C 510. R 622. Harlequin Duck.
- 724. F. vāl-līs-nēr'-ī-ā. Vallisneria is a genus of aquatic plants, the wild celery, V. spiralis L., named for Antoine Vallisner, a French botanist. The name was applied to the bird from its fondness for this plant as food. The name canvas-back, from the peculiar coloration of the upper parts, is an Americanism which has been in use at least since 1800. (e. g., see Barton, Med. and Phys. Journ., pt. i, vol. ii, 1805, p. 161.)
- 725. Clān'-gŭ-lă glaū'-cĭ-ŭm. Lat. clangula, diminutive of clangor, a clang, noise; the corresponding Gr. κλαγγή means particularly the outery of wild animals; κλάζω, future κλάγξω, I cry out. It was applied to this bird several centuries ago. Gr. γλαύκιον or γλαυκίον, a kind of wild duck, perhaps this very species. Under the varying forms of glaucion, glaucium, glaucius, and glaucia, it has been definitely applied to this duck for more than three centuries.
- 726. C. īs-lānd/-ĭ-că. To Iceland. See Falco, No. 500.
- 727. C. āl-bě'-ŏ-lă. Diminutive (irregular) form of albus, white: albula would be better form. "Buffle-head" is a corruption of buffalo-head, from the puffiness of the head: "butter-ball" from the fatness of the bird at times: "spirit duck," from the quickness of diving.
- 728. Hār-ēl'-dă glă-cĭ-ā'-līs. Harelda is a nonsense-word, invented by Leach. Lat. glacialis, glacial, iey, relating to ice; glacies, ice. (Unde derivatur? cf. Gr. γλαῦκος.)
- 729. Cāmp-tŏ-laēm'-ŭs lāb-ră-dō'-rĭ-ŭs. Gr. καμπτόs, flexible, as leather is, for instance; κάμπω, I bend; and λαιμόs, the throat; but the whole word refers to the soft leathery expansion of the bill, as if Camptorhynchus, for which latter word, preoccupied in zoölogy, it was proposed as a substitute. To Labrador; which name is said to have been given to the country by the Spaniards, it being considered cultivable, as Greenland was not; Span. labrado, cultivated land; labrador, laborer; labrar, to work.
- 730. Hīs-trī-ō'-nī-cŭs mī-nū'-tŭs. Lat. histrionicus, histrionic, relating to histrio, a stage-player; because the bird is tricked out in various colors, as if it were dressed to play some part on the stage. The word is related in the most interesting manner to historia, history, and histology, the science of tissues of the body; the idea being the weaving together of things, to make, as history, a connected account, as in histology, a tissue of organs. We still say, for example, a tissue of falsehood, &c. These words are all related to lotos, a loom, or the web woven on it.

- 731. Somateria stelleri (Pall.) Jard. B 598. C 511. R 625. Steller's Duck.
- 732. Somateria fischeri (Brandt) Coues. B 599. C 512. R 626. Spectacled Eider.
- 733. Somateria mollissima (L.) Boie. B—. C—. R 627. Eider Duck.
- 734. Somateria mollissima dresseri (Sharpe) Coues. B 606. C 513. R 627a.

  American Eider Duck.
- 735. Somateria v-nigra Gray. B 607. C 514. R 628. Black-throated Eider.
- 736. Somateria spectabilis (L.) Boie. B 608. C 515. R 629. King Eider.
- 737. Œdemia americana Sw. B 604. C 516. R 630.

  American Black Scoter.
- 738. Œdemia fusca (L.) Flem. B 601. C 517. R 631. Velvet Scoter; White-winged Scoter.
- 739. Œdemia perspicillata (L.) Flem. B 602. C 518. R 633. Surf Duck.
  - 731. Sō-măt-ĕ'-rī-ā stěl'-lĕr-ī. Gr. σῶμα, genitive σώματος, the body, and ἔριον, wool, down; with reference to the famous "eider-down" produced by species of this genus. To G. W. Steller, the surgeon and naturalist of Behring's second voyage, 1741-42.
  - 732. S. fisch'-ĕr-i. To Gotth. Fischer von Waldheim, a Russian naturalist.
  - 733. S. mol-lis'-si-ma. Lat. mollissima, superlative degree of mollis, soft; this a contraction for movilis, mobile, moveable, from moveo, I move. The reference is of course to the downy plumage.

See next species. Since the American bird has been distinguished from the European, the latter has been said to be also found in North America, on the west side of Cumberland Gulf. See Ridg., Pr. Nat. Mus., iii, 1880, p. 222. This requires us to restore the name S. mollissima, but it is No. 734 that equals No. 513 of the orig. ed.

- 734. S. m. dres-ser-i. To Henry E. Dresser, of London, author of the "Birds of Europe," &c.

  This is the S. mollissima of writers on American birds and of the orig. ed. of the
  Check List. See Sharpe, Ann. Mag. Nat. Hist., 1871, p. 51. See last species.
- 735. S. v-nīg'-ră. This is a queer way of saying that the bird has a black v-shaped mark on the throat—"digammated," indeed!
- 736. S. spēc-tă'-bĭ-lĭs. Lat. spectabilis, that may be seen, hence, worth seeing, a spectacle; specto, spicio, specio, I look at; whence a thousand derivatives.
- 737. Oē-dē'-mĭ-ă ăm-ĕr-ī-cā'-nă. Gr. οἴδημα, Lat. αdema, a swelling, tumefaction; οἰδάω, I swell; referring to the humpiness or gibbosity of the bill.
- 738. O. fūs'-că. Lat. fuscus, fuscous, dark; not well applied to this black bird.
- 739. O. pēr-spic-īl-lā/-tă. Irregularly formed from perspicio; equivalent to perspicibilis, contracted to perspicilis, and then given a participial termination, as if from a verb perspicillo; meaning perspicuous, that may be clearly seen, hence conspicuous, spectacular; see Somateria, No. 736.

- 740. Œdemia perspicillata trowbridgii (Bd.) Coues. B 603. C 518a. R —. Long-billed Surf Duck.
- 741. Erismatura rubida (Wils.) Bp. B 609. C 519. R 634. Ruddy Duck.
- 742. Nomonyx dominica (L.) Ridg. B 610. C 520. R 635. St. Domingo Duck.
- 743. Mergus merganser L. B 611. C 521. R 636. Merganser; Goosander.
- 744. Mergus serrator L. B 612. C 522. R 637. Red-breasted Merganser.
- 745. Mergus cucullatus L. B 613. C 523. R 638. Hooded Merganser.
- 746. Sula bassana (L.) Briss. B 617. C 524. R 650. Gannet; Solan Goose.
- 747. Sula leucogastra (Bodd.) Salv. B 618. C 525. R 652.
  Booby Gannet.
- 748. Pelecanus trachyrhynchus Lath. B 615. C 526. R 64'.
  American White Pelican.
  - 740. O. p. trow-brid'-gi-i. To W. P. Trowbridge, who collected in California.
- 741. Er-īs-mă-tū'-ră rūb'-ĭ-dă. Gr. ἔρεισμα, a stay, prop, pier, and οδρα, tail, as the stiffened member of the bird might seem to be. Lat. rubidus, ruddy, reddish.
- 742. Nŏm-ō'-nyx dŏm-īn'-ĭ-că. Gr. νόμος, law, order, regular way, and ὅννξ, nail. The nail at the end of the bill in all the species of so-called Erismatura, except rubida, is formed in a particular way. See Dendraca, No. 129.
- 743. Mēr'-gūs mēr-gān'-sĕr. Lat. mergus, a diver; mergo, I dive, mergere, mersi, mersum; whence submerged, immersed, &c. Merganser is simply mergus + anser, i. e., divinggoose.
- 744. M. sēr-rā'-tōr. Lat. serrator, a sawyer; serratus, sawn, i.e., saw-shaped, serrate, serried, as the prominent teeth of the bill look like those of a saw; serra, a saw; supposed to be equal to secra, from seco, I cut.
- 745. M. cŭ-cūl-lā'-tŭs. Lat. cucullatus, hooded, wearing the cucullum, a kind of hood, a capuchon, perhaps from its circular shape (κύκλος). Very appropriate in this case.
- 746. Sŭ'-lā bās-sā'-nā. Sula, by Agassiz given as a proper name, was Latinized lately from the French name, Le Sula. Quasi-Lat. bassanus is an adjective derived from the name of one of the great haunts of the bird, the Bass Rock, Firth of Forth, Scotland.
- 747. S. leũ-cŏ-gās'-tră. Gr. λευκόs, white, and γαστήρ, the belly. This stands as S. fiber in the orig. ed. See Salv., Tr. Z. S. ix, pt. ix, 1875, p. 496.
- 748. Pěl-ĕ-cā'-nùs trāch-ȳ-rhȳnch'-ūs. Gr. πελεκάν, or πελεκῶνοs, or Lat. pelecanus, a pelican. The etymology is obscure; but the pelican was fabled to strike and wound its own breast, that the young might be nourished with blood; and there are various Greek and Latin words signifying some cutting and striking instrument, as an axe, which are nearly identical in form with the above. Gr. τραχύs, rough, uneven, and ῥύγχοs, the beak; with reference to the deciduous excrescence or "centre-board" on the upper mandible.

- 749. Pelecanus fuscus L. B 616. C 527. R 641. Brown Pelican.
- 750. Phalacrocorax carbo (L.) Leach. B 620. C 528. R 642.
- 751. Phalacrocorax dilophus (Sw.) Nutt. B 623. C 530. R 643.

  Double-crested Cormorant.
- 752. Phalacrocorax dilophus cincinnatus (Brandt) Ridg. B 622. C 529.
  White-tufted Cormorant. [R 649b.
- 753. Phalacrocorax dilophus floridanus (Bartr.) Coues. B 624. C 530a. R 643a. Florida Cormorant.
- 754. Phalacrocorax mexicanus (Brandt.) S. & S. B 625. C 531. R 644.

  Mexican Cormorant.
- 755. Phalacrocorax penicillatus (Brandt) Heerm. B 626. C 532. R 645.

  Tufted Cormorant.
- 756. Phalacrocorax perspicillatus Pall. B 621. C 533. R 648.

  Pallas's Cormorant.
- 757. Phalacrocorax bicristatus Pall. B —. C 534. R 647. Red-faced Cormorant.
- 758. Phalacrocorax violaceus (Gm.) Ridg. B 627. C 535. R 646. Violet-green Cormorant.
  - 749. P. fūs'-cŭs. Lat. fuscus, fuscous, dark.
  - 750. Phăl-ā-crŏ'-cŏr-āx cār'-bō. Gr. φαλακροκόραξ, Lat. phalacrocorax, a cormorant; from φαλακρόs, bald, and κόραξ, a raven. Compare Phalacopus, No. 604. The cormorant was often ealled "sea-crow," and "cormorant" is nothing but corvus marinus; Fr. cormoran; Ital. corvo marino; Span. cuervo marino or cuervo calvo (bald-headed crow). Lat. carbo, a coal, charcoal; whence carbon; from the black color.

The cormorants are all given as *Graculus* in the orig. ed. But this was according to a way which G. R. Gray had of determining the types of genera, which has been found not available. *Graculus* signifies that the bird is so like a crow in color; cf. English "sea-crow," above.

- 751. P. di'-loph-us. Gr. δίs, twice, and λόφος, crest.
- 752. P. d. cîn-cīn-nā'-tŭs. Lat. cincinnatus, having curly hair; Lat. cincinnus, Gr. κίκιννος, a curly lock.
- 753. P. d. flō-rǐ-dā/-nŭs. To Florida. Bartram named the bird before Audubon did.
- 754. P. mēx-ĭ-cā/-nŭs. To Mexico. See Sialia, No. 28.
- 755. P. pē-nī-cīl'-lā'-tūs. Lat. penicillum, a pencil, or painter's brush; equivalent to peniculus, a little brush; this from penis, a tail, or the male organ: compare pendeo, I hang; as something pendent or appendaged. The reference is to the tufts of lengthened feathers on the bird.
- 756. P. pēr-spic-īl-lā'-tŭs. See Œdemia, No. 739.
- 757. P. bī-crīs-tā'-tŭs. Lat. bis, twice, and cristatus, crested. Exactly equal to the Gr. δίλοφος.
- 758. P. vi-ŏ-lā'-cĕ-ŭs. Lat. violaceus, violet-colored; viola, a violet. See Ionornis, No. 685.

- 759. Phalacrocorax violaceus resplendens (Aud.) Ridg. B.—. C.—. R 646a.

  Baird's Cormorant.
- 760. Plotus anhinga L. B 628. C 536. R 649. Anhinga; Darter; Snake-bird.
- 761. Tachypetes aquilus (L.) V. B 619. C 537. R 639. Frigate Bird; Man-of-war Bird.
- 762. Phaëthon æthereus L. B —. C —. R 655. (?!)
  Red-billed Tropic-bird.
- 763. Phaethon flavirostris Brandt. B 629. C 538. R 654. Yellow-billed Tropic-bird.
- 764. Stercorarius skua (Brünn.) Coues. B 652. C 539. R 696.
- .765. Stercorarius pomatorhinus (Temm.) Lawr. B 653. C 540. R 697. Pomatorhine Jäger.
  - 759. P. v. rēs-plēn'-dēns. Lat. resplendens, resplendent, splendid, or lustrous; resplende or splende, I shine, gleam. Splendor is derived by some etymologists from  $\sigma\pi\lambda\eta(\nu)\delta\delta s$ , live coals.

Not in the orig. ed. Since recognized by Ridgway, Pr. Nat. Mus., iii, 1880, p. 222. Farallone Islands.

- 760. Plō'-tus ān-hīn'-gă. Gr. πλωτόs, being a good swimmer; from πλώω or πλέω, I swim, navigate; Lat. plotus; and very early applied, in ornithology, to divers swimming birds. Anhinga is a barbarous word, from the Portuguese anhina, and equivalent to the Lat. anguina, snaky; anguis, a snake; very well applied to this curious bird, which in its subaqueous excursions strangely resembles a swimming serpent. See Coues, Bull. Nutt. Orn. Club, iii, 1878, p. 101. We should like to substitute the Latin form of the word, but that would probably be going too far.
- 761. Tāch-ȳ'-pēt-ēs ā'-quīl-ūs. Gr. ταχυπέτης, Lat. tachypetes, flying rapidly; ταχθς, swift, and πέτομαι, I fly. Lat. aquilus, swarthy, dark-colored. The word is vaguely supposed by most persons to have something to do with aquila, an eagle, in consideration of the raptorial prowess of this piratical high-flyer; but it would in that case be either aquila, substantive, an eagle, or aquilinus, adjective, aquiline. Aquila and aquilus are doubtless the same word, etymologically; but the present specific name has nothing further to do with the genus Aquila, which see, No. 532.
- 762. P. aē-thě'-rě-ŭs. Gr. αἰθέριος, Lat. αthereus, etherial, relating to the αἰθήρ, αther, ether, or serene upper air, as opposed to ἀήρ, αër, the lower aërial region; the birds of this genus being noted for soaring aloft. Th. αἴθω, ἄω.

Not in the orig. ed. If there be no mistake in identification, this species has straggled to Newfoundland. See Freke, Comp. List B. of Eur. and N. A., p. 44 (repaged from Proc. Roy. Soc. Dubl., 1879).

- 763. Phă'-ĕ-thōn fiā-vĭ-rōs'-trĭs. Gr. Φαέθων, Lat. Phaëthon, a proper name, an epithet of the sun; Phaëthon having once undertaken to drive the chariot of the sun, his father Helios; well applied to these highly aerial birds of the Tropics. Sometimes very wrongly written Phæthon, and even Phæton. Lat. flavirostris, yellow-billed.
- 764. Stēr-cŏr-ā'-rĭ-ŭs skŭ'-ă. Lat. stercorarius, having to do with ordure, a scavenger; stercus, excrement; from the filthy habits of the bird. Skua is the name applied to the bird by the Færoëse.
- 765. S. pō-mă-tŏ-rhīn'-ŭs. Gr. πωμα, genitive πωματος, a flap, lid, cover; and βls, genitive βινός,

- 766. Stercorarius parasiticus (Brünn.) Gray. B 654. C 541. R 698. Parasitic Jäger.
- 767. Stercorarius buffoni (Boie) Coues. B 655. C 542. R 699.
  Arctic Jäger; Long-tailed Jäger.
- 768. Larus glaucus Brünn. B 656. C 543. R 660. Glaucous Gull.
- 769. Larus leucopterus Faber. B 658. C 544. R 661. White-winged Gull.
- 770. Larus glaucescens Licht. B 657, 659. C 545. R 662. Glaucous-winged Gull.
- 771. Larus marinus L. B 660. C 546. R 663. Great Black-backed Gull.
- 772. Larus argentatus Brünn. B —. C 547. R 666. Herring Gull.
- 778. Larus argentatus smithsonianus Coues. B 661. C 547a. R 666a. Smithsonian Herring Gull.

the nose; from the scale-like covering of the nostrils. Temminck, habitually careless in such matters, originally wrote pomarinus, and we have almost always said "pomarine" jäger, with some vague notion of the sea in the case of this marine bird; but Newton's explanation of the word, as above, is undoubtedly correct. Säger or jæger is the German for hunter, these birds being habitual hunters and plunderers of the gulls and terns. The name was originally applied to a class of wild huntsmen who lived on the banks of the Rhine, and supported themselves entirely by plunder and robbery.

- 766. S. păr-ă-sī'-tǐ-cŭs. Gr. παρασιτικόs, Lat. parasiticus, parasitic; Gr. παράσιτοs, Lat. parasitus, a parasite, from παρά, by the side of, and σῖτοs, grain, food; literally, one who sits at the table of another; as we should say now, in vulgar parlance, a "free-luncher," "bummer," "dead-beat"; hence, in general, any kind of a hanger-on.
- 767. S. buf-fon'-i. To Jean Louis Le Clerc, Compte de Buffon, the famous French panegyrist of nature, particular friend of Linnæus, who wrote a great history of birds with the help of the Abbé de Montbeillard, and caused Daubenton to prepare the celebrated 1008 Planches Enluminées.
- 768. Lăr'-ŭs glaū'-cŭs. Gr. λάρος, Lat. larus, a gull. Lat. glaucus, glaucous, bluish, γλαυκός. See Glaucidium, No. 484. Gull is supposed to be named for its gluttony, from gulo, a glutton (gula, the gullet); Welsh, gwylan; Fr., goeland.
- 769. L. leū-cōp'-tĕr-ŭs. Gr.  $\lambda \epsilon \nu \kappa \delta s$ , white, and  $\pi \tau \epsilon \rho \delta \nu$ , wing.
- 770. L. glaū-cēs'-cēns. Lat. (decidedly post-classic) glaucescens, the present participle of a suppositious inceptive verb glaucesco, I grow bluish; meaning here somewhat bluish.
- 771. L. mă-rī'-nŭs. Lat. marinus, marine; mare, the sea.
- 772. L. ār-gēn-tā'-tŭs. Lat. argentatus, silvered, silvery; the participle of an obsolete verb argento; argentum, silver, money, from ἄργυρος, silver, ἀργός, white, the color of the metal. One writer has criticised the use of argentatus to denote a silvery color, arguing that argentatus would mean silvered over, silver-plated, or frosted, and proposed to substitute some other derivative of argentum. But this is hypercriticism; the word is more apt or fit for the bird than most specific names are.
- 773. L. a. smith-son-i-ā'-nus. To the Smithsonian Institution; this named for James Smithson, illegitimate son of Hugh Percy, Duke of Northumberland.

- 774. Larus occidentalis Aud. B 662. C 547 b. R 664. Western Herring Gull.
- 775. Larus cachinnans Pall. B —. C —. R 667. Pallas's Gull.
- 776. Larus affinis Reinh. B —. C —. R 665. (G.)
  Reinhardt's Gull.
- 777. Larus californicus Lawr. B 663. C 548a. R 668.
  Californian Gull.
- 778. Larus delawarensis Ord. B 664. C 548. R 669. Ring-billed Gull.
- 779. Larus canus L. B —. C —. R 671. (! E.)
  Mew Gull.
- 780. Larus brachyrhynchus Rich. B 665, 673. C 549. R 670.
  American Mew Gull.
- 781. Larus heermanni Cass. B 666. C 551. R 672. White-headed Gull.
- -782. Rissa tridactyla (L.) Bp. B 672. C 552. R 658.
  Kittiwake Gull.
  - 774. L. ōc-cīd-ēn-tā'-līs. See Dendræca, No. 113.
  - 775. L. că-chīn'-nāns. Lat. cachinnans, laughing immoderately; cachinno, I roar with laughter; Gr. καχάζω οι καγχάζω, of same meaning. Well expressing the outery of the gull. Not in the orig. ed.; since determined to occur in Alaska. This is L. borealis of Baird, Trans. Chicago Acad., i, 1869, p. 305.
  - 776. L. āf-fī'-nĭs. Lat. affinis, allied; ad and finis. See Campylorhynchus, No. 64.
    Not in the orig. ed.; since determined to be a good species; North American only as accidentally Greenlandic. See Reinh., Vid. Medd. Nat. För. Kjφb., 1853, p. 78.
  - 777. L. căl-ĭ-fōr/-nĭ-cŭs. To California.
  - 778. L. děl-ă-wăr-ēn'-sĭs. To the State of Delaware; named for Lord De La Ware.
  - 779. L. cā'-nŭs. Lat. canus, ashy, hoary-gray. Mediately derived from καίω, to burn, consume, the root here seen giving rise to many words, as cinereus, kinetic, &c.
    Not in the orig. ed. Since ascertained to inhabit Labrador. See Saunders, P. Z. S., 1878, p. 178, and Brewer, Bull. Nutt. Club, iii, 1878, p. 50.
- 780. L. brāch-y-rhynch'-us. Gr. βραχύs, short, and βύγχοs, beak. Note. — We give all these Lari, excepting one, as good species, in deference to recent investigation; but much doubt that the method of treating them in the orig. ed. is not more natural after all.
- 781. L. heēr'-mān-nī. To Dr. Adolphus L. Heermann, of Philadelphia, who collected extensively in the south-west.
  This stands in the orig. ed., very erroneously, as L. belcheri.
- 782. Rīs'-să trǐ-dāc'-tỹ-lă. Rissa or Ritsa is the Icelandic vernacular name. Lat. tris, thrice, and dactylus, digit, whether finger or toe: Gr.. δάκτυλος. This bird has the hind toe rudimentary, leaving only three perfect digits. Kittiwake is an old Scotch name of this species; perhaps from its cry.

- 783. Rissa tridactyla kotzebuii (Bp.) Coues. B —. C 552a. R 658a. (?) Kotzebue's Kittiwake.
- 784. Rissa brevirostris Brandt. B 674, 675. C 553. R 659. Short-billed Kittiwake.
- 785. Pagophila eburnea (Gm.) Kaup. B 676, 677. C 550. R 657. Ivory Gull.
- 786. Chroïcocephalus atricilla (L.) Lawr. B 667. C 554. R 673. Laughing Gull.
- 787. Chroïcocephalus franklini (Rich.) Bruch. B 668, 669. C 555. R 674. Franklin's Rosy Gull.
- 788. Chroïcocephalus philadelphia (Ord) Lawr. B 670. B 556. R 675. Bonaparte's Rosy Gull.
- 789. Rhodostethia rosea (Macg.) Bruch. B 678. C 557. R 676. Wedge-tailed Gull.
- 790. Xema sabinii (Sab.) Leach. B 680. C 558. R 677. Fork-tailed Gull.
- 791. Xema furcata (Prov. & Des Murs) Bruch. B 679. C 559. R 678. Swallow-tailed Gull.
- 792. Sterna anglica Mont. B 681. C 560. R 679. Gull-billed Tern; Marsh Tern.
- 783. R. t. kŏt-zĕ-buī/-ī. To Otto de Kotzebue, the Russian navigator.
- 784. R. brě-vĭ-rōs'-trĭs. Lat. brevis, short, and rostris, pertaining to the bill, rostral; from rostrum.
- 785. Pă-gŏ'-phĭ-lă ĕ-būr'-nĕ-a. Gr. πάγος, ice, and φίλος, loved. Lat. εburnea, of ivory, like ivory (in whiteness or hardness); εbur, ivory; directly from the Sanscrit word for elephant.
- 786. Chrō-ĭ-cŏ-cĕph/-ă-lŭs ā-trǐ-cīl/-lă. Gr. χρωικόs, colored, and κεφαλή, head. This word has given great trouble from Eyton's, the founder's, saying it was from κροικοs, there being no such word. Various attempts to derive it from χροιά or χροια, or from χρώs, χροόs, color, and to rectify the supposed erroneous orthography, have resulted in kroikocephalus, chracocephalus, chroicocphalus. Wharton has shown Eyton's original orthography to be correct, lacking only the diæresis over the i, there being actually such an adjective as χρωικόs, not given in the common dictionaries. (See Zoologist, March, 1878, p. —.) Lat. atricilla, black-tailed; only applicable to the young bird. See Motacilla, No. 86.
- 787. C. frank'-lin-i. To Sir John Franklin.
- 788. C. phĭl-ă-dēl'-phĭ-ă. To the City of Brotherly-Love. See Gcothlypis, No. 142.
- 789. Rhŏ-dō-stē'-thĭ-ă rŏs'-ĕ-ă. Gr.  $\dot{\rho}\dot{\phi}\delta o\nu$ , the rose, and  $\sigma\tau\hat{\eta}\theta os$ , the breast; rose-breasted. Lat. roseus, rosy.
- 790. Xĕ'-mă să-bĭn'-ĭ-ī. Xema is a nonsense word, invented by Leach: it is sometimes written zema. To Edward Sabine, by his brother.
- 791. X. fūr-cā'-tă. Lat. furcatus, forked, furcate, bifurcate, forficate; furca, a fork.
- 792. Stēr'-nă āng'-lī-că. Sterna is not classic, having nothing to do with sturnus, a starling, or with sternum, the breast-bone, or sterno, to strew. Agassiz gives the latter etymon. It is

- 793. Sterna caspia Pall. B 682. C 561. R 680. Caspian Tern.
- 794. Sterna maxima Bodd. B 683. C 562. R 681. Cayenne Tern; Royal Tern.
- 795. Sterna elegans Gamb. B 684. C 563. R 682. Elegant Tern.
- 796. Sterna cantiaca Gm. B 685. C 564. R 683. Sandwich Tern.
- 797. Sterna hirundo L. B 689. C 565. R 686.
  Common Tern or Sea Swallow.
- 798. Sterna forsteri Nutt. B 686, 691. C 566. R 685. Forster's Tern.
  - 799. Sterna macrura Naum. C 690, 693. C 567, 568. R 687 Arctic Tern.
  - 800. Sterna dougalli Mont. B 692. C 569. R 688.
    Roseate Tern.
  - 801. Sterna superciliaris antillarum (Less.) Coues B 694. C 570. R 690. Least Tern.

a Latinization, perhaps not older than about 1523, of the English tern, or stern, or sterne, or stirn, there being all these, and other old forms of the word; Danish tærne, &c. We have a vague impression that the word is onomatopæic, from the cry of the bird. One of the names of the bird is the Swiss Schnirring. Most languages, however, have a different set of words, equivalent to our sea-swallow; as Fr. Hirondelle-de-mer; Germ. Ste-Schwalbe, &c. — Lat. anglica, English; Montagu having named the bird after a country where it is comparatively seldom seen.

- 793. S. cās'-pĭ-ă. To the Caspian Sea.
- 794. S. māx'-ī-mă. Lat. maximus, superlative degree of magnus, large.

  This is S. regia of the orig. ed. We are now willing to accept Boddaert's name.
- 795. S. ĕ'-Iĕ-gāns. See Rallus, No. 673.
  This is S. galericulata of the orig. ed. We are glad to return to the orig. name of this species, which H. S. has shown to be not galericulata Licht., as S. & S. had it.
- 796. S. cān-tǐ-ā'-că. An adjective formed from Cantium, a place in Britain, mentioned by Julius Cæsar; now Kent, England.
- 797. S. hĭr-ūn'-dō. See Hirundo, No. 159.
- 798. S. fōr'-stĕr-ī. To John Reinhold Forster, who wrote, among many other things, a valuable account of Hudson's Bay birds, published in 1772.
- 799. S.  $m\bar{a}c-r\bar{u}'-r\bar{a}$ . Gr.  $\mu\alpha\kappa\rho\delta s$ , long, and  $\delta\bar{b}\rho\alpha$ , tail. The word is often written macroura, and defensibly so, the full form being macrooura. But it is permissible to shorten oou into long  $\bar{u}$ , as we habitually do in leucurus for leucoourus.
- 800. S. doū'-găl-lī. To Dr. McDougall, of Scotland.

  This stands as S. paradisea Brünn., of the orig. ed. But Brünnich's bird being unquestionably the Arctic Tern, No. 799, we do not see why the latter should not be called S. paradisea.
- 801 S. sŭp-ēr-cĭl-ĭ-ā'-rĭs. Lat. superciliaris or superciliosus, supercilious; i.e., relating to the eye-brow, supercilium; super and cilium, a hair; because one raises the eyebrows in expres-

- 802. Sterna trudeaui Aud. B 687. C 571. R 684. (! S. A.) Trudeau's Tern.
- 803. Sterna aleutica Bd. B —. C 572. R 689.
  Aleutian Tern.
- 804. Sterna fuliginosa Gm. B 688. C 573. R 691. Sooty Tern.
- 805. Sterna anæsthetica Scop. B —. C 574. R 692. Bridled Tern.
- 806. Hydrochelidon lariformis (L., 1758) Coues. B 695. C 575. R 693. Black Tern.
- 807. Hydrochelidon leucoptera (Meisn.) Boie. B —. C 575bis. R 694. (! E.) White-winged Black Tern.
- 808. Anoüs stolidus (L.) Gray. B 696. C 576. R 695.
- 809. Rhynchops nigra L. B 697. C 577. R 656.
  Black Skimmer.

sion of certain emotions, as, surprise. But when surprised at anything, we question it, or doubt it, and this implies a feeling of superiority in ourselves; hence haughtiness, loftiness, even disdain and scorn, for the person or object which makes us supercilious. Super is the Gr.  $\delta\pi\epsilon\rho$ . Cilium is the eyelid, before transferred to the eyelashes; it is the Greek  $\kappa\delta\lambda\alpha$ , the eyelids. Cilia, in the plural, has latterly been much used in the sciences for any sort of little hairs or fringes, or flagella; as, ciliated epithelium, &c. — Lat. antillarum, of the Antilles; in the genitive plural.

- 802. S. trū-deaū'-ī. To Dr. James Trudeau, of Louisiana.
  Included as North American on the authority of Audubon.
- 803. S. ă-leū'-tĭ-că. To the Aleutian Islands.
- 804. S. fū-lī-gĭn-ō'-să. See Canace, No. 559.
- 805. S. ăn-aēs-thē'-tī-că. Gr. ἀναισθητικόs, insensible, unfeeling, not perceiving; hence, as applied to this bird, stupid, foolish; ἀ or ἀν, privative, and αἰσθητικόs, sensible, &c.; αἴσθησιs, sensation, perception, feeling; αἰσθάνομαι, I perceive. We have the English æsthetic direct from the Greek, though this has experienced a refinement of meaning the original did not possess; also in medicine, anæsthesia, the state of insensibility produced by such drugs as æther or chloroform, called from their property, anæsthetic. The word has been brutally written anosthæta; anæstheta is one amendment already introduced, and the above is a further improvement.
- 806. H̄ȳ-drō-chĕ-lī'-dōn lăr-ĭ-fōr'-mĭs. Gr. ὕδωρ, water, and χελιδών, a swallow, i.e., seaswallow. Lat. lariformis, gull-like, shaped like a gull: larus and forma.
- 807. H. leū-cōp'-tĕ-rā. Gr. λευκόs, white, and πτέρον, wing. North America in one known instance (Wisconsin); see Brewer, Am. Nat., 1874, p. 188.
- 808. Ă'-nŏ-ŭs stŏl'-ĭ-dŭs. Gr. ἄνοος or ἄνους, literally mindless, unmindful of; α privative and νοῦς, the mind, intellect, understanding. It is applied to the bird as exactly equivalent to stolidus, or anæsthetica, as stolid, apathetic, insensible, in view of its indifference to the presence of man.—Lat. stolidus, stolid; related to stultus, foolish, silly.
- 809. Rhÿnch'-ōps nīg'-rā. Gr. ρύγχος, the beak, and ωψ, the face; well applied to a bird whose extraordinary beak is such a prominent feature. Lat. niger, feminine nigra, black.

- 810. Diomedea brachyura Temm. B 631. C 578. R 701. Short-tailed Albatross.
- 811. Diomedea nigripes Aud. B —, C 579. R 700. Black-footed Albatross.
- 812. Phœbetria fuliginosa (Gm.) Coues. B 633. C 580. R 703. Sooty Albatross.
- 813. Ossifraga gigantea (Gm.) Reich. B 634. C 581. R 704. (!)
  Giant Fulmar.
- 814. Fulmarus glacialis (L.) Steph. B 635. C 582. R 705. Fulmar.
- 815. Fulmarus glacialis pacificus (Aud.) Coues. B 636. C 582a. R 705a. (?)

  Pacific Fulmar.
- 816. Fulmarus glacialis rodgersi (Cass.) Coues. B —. C 582b. R 705b. (?)
  Rodgers's Fulmar.
- 817. Priocella tenuirostris (Aud.) Ridg. B 637. C 583. R 706. (!)
  Slender-billed Fulmar.
- 810. Dǐ-ŏ-mē-dē/-ă brāch-ȳ-ū/-ră. Lat. Diomedeus, adjective relating to Diomedes or Διομήδης, Jove-counselled, a Grecian hero famous at the siege of Troy: application probably fanciful. Pliny's Diomedeæ ares were birds living on the Island Diomedea in the Adriatic. Gr. βραχύς, short, and οἶρα, tail.
- 811. D. nīg'-rī-pēs. Lat. niger, black, and pes, foot.
- 812. Phoē-bē'-trī-ă fū-lī-gĭn-ō'-să. Gr. φοιβήτρια, a prophetess, soothsayer, like φοιβάστρια, Phæbastria, another genus of this family invented by Reichenbach; φοιβάζω is to prophesy; literally, to "play Apollo" with oracular utterances; Φοίβος, Phæbus, a synonym of Apollo. These words are with great propriety and correct sentiment applied to albatrosses, the import of whose weird presaging will be felt by one who reads Coleridge's "Antient Mariner," or himself goes down the deep in ships.
- 813. Os-sī'-fră-gă gĭ-gān'-tĕ-ă. Lat. ossifragus, bone-breaking, from os, genitive ossis, a bone, and frango, I break; in the perfect, fregi, participle fractus: three forms of the word repeated in English in frangible, fragile, fracture: the Latin digammated from Gr. ρήγνυμι; the stem here seen giving an immense crop of words. Lat. giganteus, gigantic, giant; the original "giants," gigantes, Γίγαντες, were a race of Titans, who attempted to scale high heaven; they were the sons of Tartarus and Earth; but, being probably illegitimate, took the name of their mother; "gigantic" meaning literally "earth-born," γηγενής; γη, and γίγνομαι.

Only North American as astray on the high sea.

- 814. Fūl'-mă-rŭs glă-c\[Ta'-l\]is. Fulmarus is arbitrary Latinization of fulmar, which is said to be akin to fulmart, foulmart, or foumart, a polecat; probably from foul (dirty), and the root of the word murder (Wharton's MS.). Glacialis, see Harelda, No. 728.
- 815. F. g. pā-ci'-fi-cus. See Anorthura, No. 77.
- 816. F. g. rŏd'-gĕr-sī. To Commodore John Rodgers, U. S. Navy.
- 817. Prī-ŏ-cēl'-lă těn-ŭ-ī-rōs'-trīs. Priocella we do not recognize, unless, perhaps, it is a frightful concatenation of Prion and Procellaria, two well-known genera of this family. French ornithologists were frequently guilty of such atrocities; see Embernagra, No. 311, for example. Agassiz gives it as Prion and Procella. Prion is the Gr. πρίων, a saw, from the prominent teeth of the bill; for Procellaria, see below.—Lat. tenurostris, slender-

- 818. Daptium capense (L.) Steph. B 639. C 584. R 719. (!)
  Pintado Petrel; Cape Pigeon.
- 819. Œstrelata hæsitata (Kuhl) Coues. B 638. C 585. R 717. (!)
  Black-capped Petrel. [See Addenda, No. 887.
- 820. Œstrelata bulweri (Jard. & Selb.) Coues. B —. C —. R 718. (G. ! E.)
  Bulwer's Petrel.
- 821. Halocyptena microsoma Coues. B —. C 586. R 720. Least Petrel.
- 822. Procellaria pelagica L. B 645. C 587. R 721. Stormy Petrel.
- 823. Cymochorea leucorrhoa (V.) Coues. B 642. C 588. R 723. Leach's Petrel.

billed; tenuis, slender, slight; more literally thin, as if spread out thin; from tenuo, I make thin, dilute, rarefy; from Gr.  $\tau\epsilon\ell\nu\omega$ , I stretch out, spread out, extend.

The bird is questionably North American, unless as astray on the high sea.

- 818. Dāp'-tǐ-tim cặp-ēn'-sẽ. Gr. δάπτιον or δύπτιον, a diminutive of δύπτης or δύπης, a diver. This set of words vary in the vowels in different dictionaries, and may not all be found; compounds of them are seen in ornithology in eudyptes, eudytes, &c. They are all from one root. The above is almost universally written daption, but in transliteration from Greek to Latin becomes properly daptium. Capense, of the Cape of Good Hope, which was the cape in those days; Caput Bonee-Spei, as it was called; caput, head, a headland. "Pintado" is painted; i. e., of variegated colors; pingo, I paint.
  Only North American as astray on the high sea.
- 819. Oēs-trē'-lă-tă haēs-ĭ-tā'-tă. Gr. oἰστρήλατοs, literally, goaded on by a gad-fly, (i. e., a goad-fly), οἶστροs, æstrus, as cattle are; hence, goaded on in any way, as these wideranging ocean birds seem to be by some mysterious impulse which drives them over the waves. The latter part of the word, -lata, the "goaded on" part of the whole idea, is from the Gr. ἐλαύνω, I urge on, drive. Lat. hæsitata, literally, stuck fast; hæsito, I stick fast, intensified from hæreo, I hang to, cleave to, adhere; in a tropical sense, I hesitate; the latter is the application in this case, the describer of the bird being uncertain about it, and therefore hesitating to name it. When at length the above generic and specific terms were combined, the bird was put in the bad way of a stuck-fast gad-about!

  Only North American as astray on the high sea.
- 820. O. bŭl'-wĕr-ī. To Bulwer.

Only North American as a straggler to Greenland. See Newton, Man. Nat. Hist. Greenl., 1875, p. 108; Freke, Zoölogist, September, 1881, p. 378.

- 821. Hăl-ō-cyp-tē'-nă mīc-rŏ-sō'-mă. Gr. ἄλs, genitive άλόs, the salt sea, ἀκύs, swift, πτηνόs, winged. Gr. μίκροs, small, σῶμα, body; "the sharp-winged little sea-body."
- 822. Prŏ-cēl-lā'-rǐ-ă pěl-ă'-gĭ-că. Lat. procellaria or procellosa, stormy, tempestuous, relating to storm; procella, a storm. Gr. πελαγικόs, pelagic, relating to the sea; thoroughly Greek, but transliterable into Latin. Petrel is commonly fancied to be a diminutive of Peter, Petrus, who attempted to walk on the sea of Galilee, as these little birds seem to be continually doing, in the way they patter over the ocean waves; but there are many forms of petrel, as petteril, peterel, &c., and the word may be related to the verb to patter, just used.
- 823. Cỹ-mŏ-chŏr-ē'-ă leū-cōr'-rhŏ-ă. Gr. κῦμα, genitive κύματοs, the surging billows, and χορεία or χορῆα, a choir, a dancing; literally, the wave-dancers. One of my critics has favored me with an excellent reason why, according to his faithful dictionary, the

- 824. Cymochorea melæna (Bp.) Coues. B 643. C 589. R 724. Black Petrel.
- 825. Cymochorea homochroa Coues. B —. C 590. R 725.
  Ashy Petrel.
- 826. Oceanodroma furcata (Gm.) Bp. B 640. C 591. R 726. Fork-tailed Petrel.
- 827. Oceanodroma hornbyi (Gr.) Bp. B 641. C 592. R 727. Hornby's Petrel.
- 828. Oceanites oceanicus (Kuhl) Coues. B 644. C 593. R 722. Wilson's Petrel.
- 829. Fregetta grallaria (V.) Bp. B 646. C 594. R 728. (!)
  Lawrence's Petrel.
- 830. Priofinus melanurus (Bonn.) Ridg. B 651. C 595. R 707. (!)

  Black-tailed Shearwater.

word ought to have been *cymatochoreutes*. We would refer him to his dictionary again for certain words beginning with *sync*- and *euph*-. The stem of the first part of the word is seen in *accumulate*, to roll up; of the second in *chord*, *choir*, *choral*, *choresis*, or *chorea* (St. Vitus's dance), &c. — Gr. λευκόs, white, and δρόρs, the rump.

- 824. C. měl-aē'-nă. Gr. μέλας, feminine μέλαινα, black. The orthography introduced by Bonaparte, melania, requires to be emended as above.
- 825. C. hŏ-mō'-chrŏ-ă. Gr. δμός, equal, like, and χρόα, color; in allusion to the unicolor plumage.
- 826. O-cĕ-ăn-ŏ'-drŏ-mă fūr-cā'-tă. Gr. 'Ωκεανός, Oceanus, the divinity of, and the ocean itself; supposed to be ἀκύς, swift, and νέω, I flow. See Ammodramus, No. 238, and Hydranassa, No. 660.— Lat. furcatus, forked; furca, a fork.
- 827. Ö. hörn'-bÿ-ī. To Admiral Hornby, R. N.
- 828. Ō-cĕ-ăn-ī'-tēs ō-cĕ-ăn'-ĭ-cŭs. Gr. ἀκεανίτης, a son of the sea; sprung from Oceanus. See Oceanodroma, No. 826. Gr. ἀκεανικός, oceanic.
- 829. Fre-get'-tă grāl-lā'-rī-ă. Fregetta, fregeta, fregeta, as variously spelled, is from the Ital. fregeta, Span. fragata, Fr. frégete, Eng. frigate; according to Diez, the Lat. fabricata; originally applied in French ornithology to the bird called man-of-war, Tachypetes aquilus; applied by English ornithologists about 1790 to some species of the present family, and very lately taken by Bonaparte for a generic term.— Grallæ, among the Romans, was a pair of stilts, the word being contracted from gradula, this from gradus, a step; and the Grallatores were people who acted on the stage on stilts. The word was early taken in ornithology for wading birds, called grallæ or grallatores, from their length of leg; from these words we have derived the English adjectives grallarial and grallatorial; and grallaria is an obvious easy Latin derivative, though probably never used by the Romans.

Only North American as astray on the high sea.

830. Prǐ-ŏ'-fin-ŭs mĕl-ăn-ū'-rŭs. Priofinus, unless we are mistaken, is a dreadful concoction of prion and puffinus, by the same victims of misapplied ingenuity who gave us Priocella; see this, No. 817, and Puffinus, next below. — Gr. μέλαs, genitive μέλανος, black, and οδρα, tail.

Only North American as astray on the high sea.

- 831. Puffinus kuhli (Boie) Bp. B —. C 596. R 708. Cinereous Shearwater.
- [See Addenda, No. 888.
- 832. Puffinus major Faber. B 647. C 597. R 709.
  Greater Shearwater.
- 833. Puffinus creatopus Coop. B —. C 598. R 710. Flesh-footed Shearwater.
- 834. Puffinus anglorum Temm. B 649. C 599. R 711.

  Manks Shearwater.
- . 835. Puffinus obscurus (Gm.) V. B 650. C 600. R 712. Dusky Shearwater.
  - 836. Puffinus opisthomelas Coues. B —. C 601. R 713.

    Black-vented Shearwater.
  - 837. Puffinus fuliginosus A. Strickl. B 648. C 602. R 714. Sooty Shearwater.
  - 831. Puf'-fin-us kuhl'-i. Two very different kinds of birds early received the name of puffein or puffin: one of these, the Fratercula arctica, has retained it in English, in place of the old English coulterneb ("ploughshare-nose"), which soon gave way; the French now call it macareux. The other, namely, the shearwater, soon lost the name of puffin; but meanwhile puffin had been taken into the books, and, at the pen of those who wrote their treatises in Latin, became puffinus or puphinus; and this was subsequently fixed as a generic term for the Shearwater Petrels. We do not know the exact meaning of the word, but suppose it has something to do with puff, as suggested by the stout, "puffy" shape of the bodies of the Auks, as if puffed up. The species is dedicated to Dr. Heinrich Kuhl, whose early death left much promise unfulfilled.
  - 832. P. mā'-jor. Lat. major, greater, comparative degree of magnus, great.
  - 833. **P.** crĕ-ăt'-ŏ-pūs. Gr. κρέαs, genitive κρέατοs, flesh, and ποῦs, foot; in allusion to the color of the feet. We see the same stem in the anatomical term pan-creas, "all-flesh."
  - 834. P. ān-glō'-rǔm. "Puffinus anglorum" is a curiosity. It simply says in Latin "the puffin of the English," just as one might cite Puffinus jonstoni, the puffin of Jonston's treatise. Willughby, edited in Latin in 1676, called it "Puffinus Anglorum," meaning only that it was the bird "called puffin in English"; and Temminck, in 1820, not unhappily made the phrase generic and specific as the technical name of the bird.—"Manks" or "Manx" is the name of the people and of their language, of the Isle of Man; so "manx shearwater" is as if we were to say "the puffin of the Isle of Man."—"Shearwater" is defined by early ornithologists as "avis aquæ superficiem radens,"—the bird that grazes, skims, shaves, shears over the surface of the water; rado, I shave, scrape; the stem is seen in erase, razor, &c. See above, Puffinus, No. 831.
  - 835 P. ōb-scū'-rūs. Lat. obscurus, dark-colored. Note. — There is doubt that the small dark shearwater of our South Atlantic coast is the P. obscurus of Gmelin, and Finsch has lately proposed to call it P. auduboni. But until we have more light on this obscure group, we prefer not to disestablish several well-settled names in this genus. See Ridg., Pr. Nat. Mus., ii, 1880, p. 12.
  - 836. P. ŏ-pīs-thŏ'-mĕ-lăs. Gr.  $\delta\pi\iota\sigma\theta\epsilon$ , backward, and  $\mu\epsilon\lambda\alpha s$ , black; a Greek way of saying black behind.
    - Note. This is supposed by some to be Puffinus gavia (Forst.).
  - 837. P. fū-lī-gīn-ō'-sŭs. See Canace, No. 559.

- 838. Puffinus amaurosoma Coues. B —. C 603. R 715. Spectral Shearwater.
- 839. Puffinus tenuirostris Temm. B —. C 604. R 716.
  Slender-billed Shearwater.
- 840. Colymbus torquatus Brünn. B 698. C. 605. R 736. Great Northern Diver or Loon.
- 841. Colymbus torquatus adamsi (Gr.) Coues. B —. C 605a. R 737. Yellow-billed Loon.
- 842. Colymbus arcticus L. B 699. C 606. R 738. Black-throated Diver.
- 843. Colymbus arcticus pacificus (Lawr.) Coues. B 700. C 606a. R 739. Pacific Black-throated Diver.
- \*844. Colymbus septentrionalis L. B 701. C 607. R 740. Red-throated Diver.
- 845. Æchmophorus occidentalis (Lawr.) Coues. B 704. C 608. R 729. Western Grebe.
- 846. Æchmophorus occidentalis clarki (Lawr.) Coues. B 705. C 608a. R 730. Clark's Grebe.
- 847. Podicipes griseigena holbælli (Reinh.) Coues. B 702. C 610. R 731.

  American Red-necked Grebe.
  - 838. P. ă-maū-rŏ·sō'-mă. Gr. ἀμαυρός, dark, dim, dusky, and σῶμα, body. Note. — This is probably Proc. grisea Gm., as held by Finsch and Salvin.
  - 839. P. těn-ŭ-ĭ-rōs'-trĭs. See Priocella, No. 817.
  - 840. Cŏ-lym'-būs tōr-quā'-tūs. The Latin colymbus is simply a transliteration from the Greek, and has nothing to do, notwithstanding the great similarity, with the purely Latin columba, a dove; the latter being not Greek at all, nor the former Latin, except as directly transferred from the Greek. The two words are consequently not related, unless it be in a radical manner; Corssen, however, considers them to be the same. Gr. κόλυμβος οr κολυμβίς, a diver or swimmer; κολυμβάω, I dive, swim. The κόλυμβις of Aristotle was a species of grebe (Podicipes).—Lat. torquatus, see Asyndesmus, No. 456.—
    "Loon" is an old Scotch word. See No. 874.
  - 841. C. t. ă'-dăms-ī. To Dr. C. B. Adams, of the British Navy.
- 842. C. ārc'-tĭ-cŭs. See Sialia, No. 29.
- 843. C. a. pā-cĭ'-fī-cŭs. See Anorthura, No. 77.
- 844. C. sēp-tēn-trī-ō-nā'-līs. Lat. septentrionalis, northern; septentriones, the north, northern regions; septem-trio, the constellation of the Wain. See Parus, No. 45.
- 845. Aēch-mŏ'-phŏr-ŭs ōc-cĭd-ēn-tā'-lĭs. Gr. αἰχμή, a spear, and φορόs, bearing; in allusion to the long, slender, sharp bill. For occidentalis, see Dendraca, No. 113. Grebe is a French word, the meaning of which we do not know.
- 846. A. o. clārk'-ī. To J. H. Clark.
- 847. Pŏd-i'-cī-pēs grīs-ĕi'-gĕn-ă hŏl'-boĕl-lī. The extraordinary word "podiceps" has excited much curiosity, and stimulated some ingenious surmises. As it stands, podiceps seems to be the Greek ποῦs, genitive ποδόs, foot, and the Latin termination -ceps, denoting head; and "foot-head" it has doubtless been taken to be by many, who, if thinking of it at all, have felt vaguely that some allusion was intended to the bird's somersaulting

- 848. Podicipes cornutus (Gm.) Lath. B 706. C 611. R 732. Horned Grebe.
- 849. Podicipes auritus (L.) Lath. B 708. C —. R 733. (G.) European Eared Grebe.
- 850. Podicipes auritus californicus (Heerm.) Coues. B 707. C 612. R 733a.

  American Eared Grebe.
- 851. Podicipes dominicus (L.) Lath. B 708a. C 613. R 734. St. Domingo Grebe.
- 852. Podilymbus podicipes (L.) Lawr. B 709. C 614. R 735. Pied-billed Grebe; Dab-chick.
- 853. Fratercula corniculata (Naum.) Gray. B 713. C 617. R 744.
  Horned Puffin.

in the water,—turning "heels over head," as we should say. In deriving the name of the family of grebes, some curious words have been ventured; as Podicipinæ, as if the genitive were podicipis, or Podicipitinæ, as if the genitive were podicipitis. There is no doubt that podiceps, and everything derived from it, is absurd. We have not traced the word back of 1758, when it probably originated in a misprint. Going back further in the annals of ornithology, we soon come upon the word in its proper form, viz., podicipes, occurring repeatedly in Willughby and various writers of about that period. The word is the Latin podex, genitive podicis, the rump, buttocks, and pes, foot; being simply a translation into Latin of a very vulgar English name. Having crystallized in the shape of podiceps, by Latham's employ of the word as a generic term, and then been used for a century, it will not be easy to eradicate; but the attempt should be made to substitute the proper podicipes. The genitive of this is podicipedis, and the family name should be Podicipedide.—Lat. griseus, gray: gena, cheek.

Note. — There is no technical reason or excuse for using the word at all. For Colymbus, Brisson, 1760, is the proper name for the genus of grebes, having meant Grebe, not Loon, from the time of Aristotle to that of Linnæus, when the latter used it for loons and grebes indiscriminately. The loons were called Mergus by Brisson; and Eudytes, Illiger, 1811, seems to be the tenable generic name for them.

- 848. P. cōr-nū'-tŭs. Lat. cornutus, horned; cornu, a horn; in reference to the tufts of feathers on the head.
- 849. P. aūr-ī/-tŭs. Lat. auritus, eared; auris, an ear; Gr. οδs, genitive ἀτόs, ear; in allusion to the auricular tufts of feathers.
  Not in the orig. ed. Only North American as occurring in Greenland.
- 850. P. căl-ĭ-fōr/-nĭ-cŭs. To California.
- 851. P. dom-in'-i-cus. To the Island of St. Domingo. See Dendraca, No. 129.
- 852. Pŏd-ĭ-lym'-bŭs pŏd-ĭ'-cĭ-pēs. The word podilymbus, sometimes aggravated into podylimbus, is a peculiarly villanous miscegenation of podi[ceps and co]lymbus; see the latter word, No. 840, and Podicipes, No. 847.
- 853. Frā-tēr'-cŭ-lă cōr-nǐ-cŭl-ā'-tă. "Fratercula" is a singular word, the application of which to this bird is not obvious, and the form of which seems absurd: a feminine noun meaning "little brother." Fraterculus is a proper classical word, a diminutive of frater, brother. But there is no larger bird sufficiently near this species for the latter to be called the "little brother." Fratercula in ornithological writing is much older than 1760, when Brisson made a genus of it, and we are inclined to think that it is humorously used; all the more so by being made feminine, in the same spirit that prompted the comic writer Plautus to invent the verb fraterculo, as he did sororio, to signalize the swellings of the breasts of boys, like twin-brothers, at puberty. If there be anything in

- 854. Fratercula arctica (L.) Steph. B 715. C 618. R 743. Common Puffin; Sea Parrot.
- 855. Fratercula arctica glacialis (Leach) Coues. B 714. C 618a. R 743a. Large-billed Puffin.
- 856. Fratercula cirrata (Pall.) Steph. B 712, 716. C 619. R 745.
  Tufted Puffin.
- 857. Ceratorhina monocerata (Pall.) Cass. B 717, 718. C 620. R 746.
  Horn-billed Auk.
- 858. Simorhynchus psittaculus (Pall.) Schl. B 725. C 621. R 747.

  Parroquet Auk.
- 859. Simorhynchus cristatellus (Pall.) Merr. B 719, 720. C 622. R 748.
- 830. Simorhynchus pygmæus (Gm.) Ridg. B 721. C 623. R 749. Whiskered Auk.
- 861. Simorhynchus pusillus (Pall.) Coues. B 722, 723. C 624. R 750. Knob-billed Auk.
- 862. Ptychorhamphus aleuticus (Pall.) Brdt. B 724. C 625. R 751.

  Aleutian Auk.

this, the application of the word to the birds is to be sought in their stout puffy shape, that which appears to have caused the English word puffin. See Puffinus, No. 831.—Lat. corniculata, horned, a diminutive of cornutus: referring to the acute epidermal process on the upper eyelid, which is deciduous, being shed like the horns of deer.

- 854. F. ārc'-tĭ-că. See Sialia, No. 29.
- 855. F. a. glă-cĭ-ā'-lĭs. See Harelda, No. 728.
- 856. F. cīr-rā'-tă. Lat. cirratus or cirrhatus, having curled locks, or ringlets, curly-haired; cirrus or cirrhus, a curl of hair; well applied to this oddly feather-tufted bird.
- 857. Cĕr-āt-ŏ-rhī'-nă mŏ-nŏ-cĕr-ā'-tă. Gr. κέρας, genitive κέρατος, a horn, and ρίς, genitive ρ̂ινός, the nose; alluding to the prominent deciduous horn which grows up from the base of the bill, over the nostrils.—Gr. μόνος, only, alone, single, transliterated as Lat. mono-, in composition, and κέρας, horn, Latinized as cĕras, ceratis, whence an adjective form, ceratus, horned; that is, unicorn, one-horned.
- 858. Si-mŏ-rhyn'-chŭs psīt-tă'-cŭ-lŭs. Gr. σιμόs, flat-nosed, snub-nosed, like the negro; δύγχοs, beak; well applied to these birds, whose bills are singularly shaped. The same idea is expressed in the Latin simia, an ape, whence our English simian, become a common word since Darwinism has been so much discussed.—Lat. psittaculus, a little parrot, diminutive of psittacus. See Sitta, No. 57.
- 859. S. crīs-tă-tēl'-lus. Lat. diminutive of cristatus, crested.
- 860. S. pyg-maē'-ŭs. See Sitta, No. 61.

  This is S. camtschaticus of the orig. ed.; but Alca pygmwa Gm. is based on the young of the same species, called S. cassini by Coues.
- 861. S. pŭs-īl'-lŭs. See Sitta, No. 60.
- 862. Pty-chŏ-rhām'-phùs ă-leū'-tǐ-cùs. Gr. πτόξ, genitive πτυχός, a fold, and βάμφος, the beak; well alluding to the wrinkled covering of the bill; which, by analogy with what is known of other species, may be taken as an indication that the soft part concerned will be found to grow some kind of excrescence, not yet discovered.—Lat. aleuticus, of the Aleutian Islands,—the country of the people called Aleuts.

- 863. Alle nigricans Link. B 738. C 626. R 752. Sea Dove; Dovekie.
- 864. Synthliborhamphus antiquus (Gm.) Coues. B 736. C 627. R 753. Black-throated Auk.
- 865. Synthliboramphus umizusume (Temm.) Coues. B 737. C 628. R 754. Temminck's Auk.
- 866. Brachyrhamphus marmoratus (Gm.) Brdt. B 732, 733. C 629. R 755.

  Marbled Murrelet.
- 867. Brachyrhamphus kittlitzi Brdt. B 735. C 630. R 756. Kittlitz's Murrelet.
- 868. Brachyrhamphus hypoleucus Xant. B —. C —. R 757. White-bellied Murrelet.
- 869. Brachyrhamphus craverii (Salvad.) Coues. B —. C —. R 758. (?)
  Craveri's Murrelet.
- 870. Brachyrhamphus brachypterus Brdt. B 734. C —. R 759. (?)
  Short-winged Murrelet.
- 863. Al'-lĕ nĭg'-rĭ-cāns. Alle is a local designation of this species and of Uria grylle. Its meaning we do not know. It was long the specific name of the bird, taken for the generic by Link in 1806. Lat. nigricans, present participle of a supposed verb nigrico, equivalent to nigresco, I grow black, am blackish.

This stands in the orig. ed. as *Mergulus alle*: for the reason of the change, see Coues, Bull. Nutt. Club, iv, 1879, p. 244.

- 864. Sỹn-thlǐ-bŏ-rhām'-phùs ān-tī'-qǔ-uš. Gr. συνθλίβω, I compress; σύν, with, and θλίβω, I press; βάμφος, beak.—Lat. antiquus, antique, ancient; with reference to the gray of the head, like an old man's. Antiquus is simply for anticuus, this a form of anticus, from ante, before; one having retained the idea of being before in space, that is, in front of, the other having acquired the idea of priority in time, like antea; the opposition in either case is with post, postea, behind, after.
- 865. S. ū-mǐ-zŭ'-sŭ-mē. This appalling word we know nothing about except that it is transliterated from the Japanese, Temminek having described the species from that country; "son nom japonais est wumizusume," he says. We drop the w.
- 866. Brāch-y-rhām'-phùs mār-mŏ-rā'-tùs. Gr. βραχύs, short, and βάμφοs, beak. Lat. marmoratus, marbled; marmor, marble; in allusion to the veined and clouded color. Marmor is the Gr. μάρμαροs, from μαρμαίρω, I shine, glitter, sparkle, as did the beautiful white stone which Praxiteles carved. Murrelet is a word coined by Coues in 1868 as a diminutive of murre, like rivulet from river.
- 867. B. kitt'-litz-i. To F. H. von Kittlitz, traveller and naturalist.
- 868. B. hÿp-ŏ-leū'-cŭs. Gr. ὑπό, under, and κευκόs, white, meaning neither whitish nor under the white, but white underneath.
  Not in the orig. ed.; since recognized as probably valid.
- 869. B. cră-vē'-rĭ-ī. To Sig. Federico Craveri.

  Not in the orig. ed.; since recognized as perhaps distinct.
- 870. B. bră-chyp'-tĕ-rŭs. Gr.  $\beta\rho\alpha\chi\dot{v}s$ , short, and  $\pi\tau\dot{\epsilon}\rho\sigma\nu$ , wing. Not in the orig. ed. Since recognized by Ridgway. Pacific Coast.

- 871. Uria grylle (L.) Brünn. B 726, C 631. R 760.
  Black Guillemot; Sea Pigeon.
- 872. Uria columba (Pall.) Cass. B 727. C 632. R 761.
  Pigeon Guillemot.
- 873. Uria carbo (Pall.) Brdt. B 728. C 633. R 762. Sooty Guillemot.
- 874. Lomvia troile (L.) Brdt. B 729, 730. C 634. R 763. Common Guillemot; Murre.
- 875. Lomvia troile californica (Bry.) Coues. B —. C —. R 763a. (?)
  California Guillemot.
- 876. Lomvia arra (Pall.) Coues. B 731. C 635. R 764, 764a.
  Thick-billed Guillemot.
- 877. Utamania torda (L.) Leach. B 711. C 616. R 742.
  Razor-billed Auk.
- 878. Alca impennis L. B 710. C 615. R 741. Great Auk.
- 871. U'-rı-a gryl'-lē. Gesner and others state that οὐρία is the Greek name of a guillemot, or some other water-bird. Uria occurs all through ornithology from Gesner, and was made a genus by Brisson in 1760. The meaning we do not know; perhaps akin to urinari, Skr. nāri, water. Gylle is said to be from Gr. γρυλλίζω, I grunt; the bird has been called sibilans by some; but grisla and grylle are N. European names.
- 872. U. cŏl-ūm'-bă. Lat. columba, a pigeon, applied in the same way that we call the bird "sea-pigeon" in English.
- 873. U. cār'-bō. Lat. carbo, a coal, charcoal; here used in allusion to the uniformly sooty color, as if the bird were charred.
- 874. Lōm'-vǐ-ā trō-ī'-lē. Lomwia and lomvia are two of many forms in which is found spelled the vernacular name of the bird, in Scotch, Færoëse, and related languages; as Dan. lûn, Dutch loen, Eng. loon or loom. It was taken by Linnæus for the specific, and much later by Brandt for the generic name.— Troile, on the contrary, may be of classic origin, Troilus being the son of Priam; also used as synonymous with Trojan; application in this case arbitrary, if any. Newton says "possibly a compliment to Troil, the Icelander." Brünnich wrote it Troille in 1764 (Orn. Bor., p. 27).
- 875. L. t. căl-ī-fōr'-nī-că. Pertaining to the State of California.

  Not in the orig. ed. Since recognized by Ridgway.
- 876. L. ār'-rā. Lat. arra or arrha was purchase-money, or a pledge in earnest of a contract, and might have been applied by Pallas to a bird in such demand by the natives as to serve as a sort of unit or standard of exchange in barter. "Salerne says the great blue parrot of Brazil is called Arras or Aras; this seems here transferred to the sea-parrot." (Wharton's MSS.)
- 877. Ŭ-tă-măn'-ĭ-ă tōr'-dă. Both these words are mere Latinizations of vernacular names.

  \*Utamania or utumania\* was in the bird-books long before Leach made a genus of it, and so was tord or tordmule. We do not know what these words mean, further than that they signify this species. Ray says (Syn., 1713, p. 119): "Ad litora Cretæ invenitur; indigenis 'Utamania' dicta."
- 878. Al'-că îm-pēn'-nĭs. Alca is not classic, being merely a Latinization of the vernacular name, found in several different forms, as alk, alck, alka, auk, auk. The third of these

is found in the old treatises written in Latin, and the change to alca is of course immaterial. The meaning of the word is in question. The form awk (which we observe some late English scholars use) might suggest a relationship with awkward, in view of these ungainly fowl; but awkward means simply left-handed. Quite probably alk is related, and not distantly, to elk, the bird and the beast being the largest, or most notable, or most prevailing animals of their respective kinds in the consideration of the people. But elk is in Latin alce (quite like alca), and this is uniform with the Greek ἀλκή, meaning strength, prowess; one of the names of Hercules, for example, being derived therefrom. The probability that alk, elk, alce, and άλκή are radically if not still more closely related, is heightened by the other vernacular names of this bird, gare-foul, goir-fugel, &c., these qualifying prefixes being similar to those seen in gerfalcon, and recognized by Steenstrup in inventing his genus Gyralca, the idea of size, strength, or other predominance being evident. If this be so, the alk, the Gare-fowl, is the fowl, par excellence, as elk, alce, is the great beast, as Ger-falco is the falcon; with the implication of some honor or special esteem. We are thus led directly to Hierofalco, which see, No. 498. — Lat. impennis, featherless, i. e., wingless, with reference to the diminutive wings, unfit for flight; in, negative, and penna, a feather.

Though the Great Auk is extinct in North America, and has doubtless disappeared from the face of the earth, we still keep the place in memoriam of this "most honourable and antient fowle."

## ADDENDA.

The foregoing list of 878 names agrees with the analysis of the original and of the present edition of the Check List: 778 - 10 + 110 = 878.\* But in the course of the year during which this edition has been printing, the following ten additions to the bird-fauna of North America have been announced:—

- 879. Parus meridionalis Scl. B 292. C —. R 43. Mexican Titmouse.
- 880. Myiarchus crinitus cooperi (Bd.) Coues. B 132. C —, R 311.

  Mexican Great-crested Flycatcher.
- 881. Antrostomus vociferus arizonae Brewst. B —. C —. R —. Arizona Whippoorwill.
- 882. Buteo brachyurus V. B —. C —. R —. (!S. A.)
  Short-tailed Buzzard.
  - 879. Pā'-rǔs mĕ-rī-dǐ-ō-nā'-lĭs. Lat. meridionalis, southern, southerly; for meridialis, from meridies, midday, noon.

    Not in the orig. ed. Since ascertained to occur in Arizona. Brewst., Bull. Nutt. Club, vi, No. 4, October, 1881, p. 252.
- 880. Mỹĩ-ār'-chṃs crī-nī'-tms coō'-pĕr-ī. To William Cooper.

  Not in the orig. ed. Since ascertained to occur in Arizona. Brewst., Bull. Nutt.

  Club, vi, No. 4, October, 1881, p. 252.
- 881. An-trō'-stŏ-mŭs vō-cī'-fĕr-ŭs ă-rī-zō'-naē. To the Territory of Arizona.

  Not in the orig. ed. Since described from Arizona. Brewst., Bull. Nutt. Club, vi,
  No. 2, April, 1881, p. 69.
- 882. Bū'-tĕ-ō bră-chy-ū'-rūs. Gr. βραχύs, short, and οὖρα, tail. Not in the orig. ed. Since ascertained to occur in Florida. See Ridg., Bull. Nutt. Club, vi, No. 4, October, 1881, p. 210.
- \* The stereotyped plates of the introductory pages, indicating 878 names, with 110 additions, have been punched to give the total of 888 with 120 additions.

- 883. Buteo fuliginosus Scl. B —. C —. R —. (!S. A.) (?)
  Sooty Buzzard.
- 884. Eurynorhynchus pygmæus (L.) Pearson. B —. C —. R —. (!A.) Spoon-billed Sandpiper.
- 885. Fulica atra L. B —. C —. R —. (G.) European Coot.
- 886. Fuligula rufina (Pall.) Steph. B —. C —. R —. (! E.)
  Rufous-crested Duck.
- 887. Œstrelata gularis (Peale) Brewst. B —. C —. R —. (!) Gular Petrel.

are informed that the alleged occurrence is questionable.

- 888. Puffinus borealis Cory. B —. C —. R —. (?)
  Northern Shearwater.
- 883. B. fū-lī-gĭn-ō'-sŭs. Lat. fuliginosus, sooty, of a dark sooty color; fuligo, soot.

  Not in the orig. ed. Since ascertained to occur in Florida (if really distinct from B. brachyurus). See Ridg., Bull. Nutt. Club, vi, No. 4, October, 1881, p. 212.
- 884. Eū-ry-nŏ-rhyn'-chus pyg-maē'-us. Gr. εὐρύνω, I dilate, widen, spread out; from εὐρύς, broad; and ρύγχος, beak. It is found spelled in many different ways; often eurin- or eurhin-, as if supposed to be εὖ and ρίς or ρίν, the nose. See Sitta, No. 61.
  Not in the orig. ed. of the Check List. Since given by Ridgway as occurring at Point Barrow, Arctic coast of Alaska, in Bull. U. S. Nat. Mus., No. 21, 1881, p. 85. We
- 885. Fül'-ĭ-că ā'-tră. Lat. ater, atra, atrum, black. Not in the orig. ed. of the Check List. Since reported to have been obtained in Greenland in 1876. See Ridg., Bull. U. S. Nat. Mus., No. 21, 1881, p. 85, and Freke, Zoölogist, September, 1881, p. 374.
- 886. Fŭl-ĭg'-ŭ-lă rū-fi'-nă. Late Lat. rufinus, reddened, reddish, formed from rufus, of same meaning.

Not in the orig. ed. of the Check List. Specimen said to have been procured in Fulton Market, New York, February, 1872, and to be now in the Nat. Mus. at Washington. Ridg., Bull. U. S. Nat. Mus., No. 21, 1881, p. 85; Allen, Bull. Nutt. Club, vi, 1881, p. 173.

- 887. Oēs-trē'-lā-tā gŭl-ā'-rīs. Lat. gularis, pertaining to gula, the throat. Not in the orig. ed. Since ascertained to occur in New York. See Brewst., Bull. Nutt. Club, vi, No. 2, April, 1881, p. 94. Only North American as astray on the high sea.
- 888. Pŭf'-fin-ŭs bŏr-ĕ-ā'-līs. Lat. borealis, northern.

  Not in the orig. ed. Since described as new from Massachusetts. See Cory, Bull.

  Nutt. Club, vi, No. 2, April, 1881, p. 84.

## LIST OF WORDS DEFINED.

- Note (1). This Index contains matter additional to or corrective of that in the body of the text.
- Note (2). The figures refer, not to the pagination of the book, but to the numeration of the names:

  e. q., Turdus, No. 1, not page 1.
- Note (3).— When a word occurs in the List more than once,—as in cases of most generic and many specific or subspecific names,—the reference is usually to the place where it is first or best defined.
- Note (4). Words differing only in termination, as american-us,-a,-um, are usually not duplicated.
- Note (5). Generic names are distinguished by a capital initial letter.

abbreviatus, 522 aberti, 309 acadicus, 384, 483 Accipiter, 494 accipitrinus, 473 Actodromas, 614 aculeata, 58 acuminata, 619 acutipennis, 402 adamsi, 841 Æchmophorus, 845 Ægialites, 584 Ægiothus, 207 ægocephala, 630 æneus, 315 æstiva, 111 æstivalis, 251 æthereus, 762 affinis, 64 Agelæus, 316 agilis, 139 aglæus, 337 agrestis, 271

aikeni, 262

Ajaja, 653

Alauda, 85

alascensis, 78

Aïx, 719

alaudinus, 229 alba, 86 albatus, 696 albeola, 727 albicilla, 533 albicollis, 275, 395 albifrons, 542, 692 albigula, 307 albilora, 130 albocaudatus, 513 albolarvatus, 442 Alca, 878 alcvon, 423 aleutica, 803, 862 alexandri, 410 aliciæ, 12 Alle, 863 alleni (C. A.) 412 alleni (J. A.) 302 alpestris, 82 alpina, 623 altiloquus, 172 Aluco, 461 amabilis, 545 amaurosoma, 838 Amazilia, 419 ambiguus, 422 americana, 93

Ammodramus, 238 amœna, 294 Ampelis, 166 Amphispiza, 258 anæsthetica, 805 Anas, 707 Ancylochilus, 625 anglica, 792 anglorum, 834 angustifrons, 455 anhinga, 760 ani, 425. Said to be the Brazilian name of the bird annæ, 414 annectens, 264, 351 Anorthura, 76 Anoüs, 808 Anser, 692 anthinus, 228 anthracina, 528 Anthus, 88 antillarum, 801 antiquus, 861 Antrostomus, 396 Aphelocoma, 354 Aphriza, 594 Aquila, 532 aquilus, 761

Aramus, 671 Archibuteo, 525 arctica, 29 arctoa, 206 Ardea, 655 Ardetta, 667 arenaria, 627 argentatus, 772 arizonæ, 253 Arquatella, 620 arra, 876. Both conjectures in the text wrong. Simply onomatopœic; Russian arrie or arra arvensis, 85 Asio, 472 asio, 465 Astragalinus, 213 Astur, 496 Asturina, 527 Asyndesmus, 456 ater, 313 atrata, 201, 538 atricapillus, 44, 185 atricilla, 786 atrigularis, 274 atrocristatus, 42 Atthis, 416. It is also a proper name auduboni, 9, 120 aura, 537 aurantia, 121 auratus, 457 auricapillus, 135

bachmani, 103
bahamensis, 153
bairdi, 224
barbatulus, 172
Bartramia, 640
Basilinna, 407
bassana, 746
belli, 183
bendirii, 21
berlandieri, 70
Bernicla, 699
bewicki, 71
bicolor, 40, 286
bicristatus, 757
bilineata, 258

aurifrons, 451

Auriparus, 56

australis, 202 autumnalis, 706

auritus, 849

blackburnæ, 121. This probably should stand D. aurantia, Coues bonapartii, 617 Bonasa, 565 borealis, 32 boscas, 707 Botaurus, 666 boucardi, 256. Peucæa ruficeps boucardi is not in the orig. ed. Since discovered in Arizona by H. W. Henshaw, Ornith. Wheeler's Surv., 1875, p. 117. Brachyrhamphus, 866 brachyrhynchus, 780 brachypterus, 870 brachyura, 810 brenta, 700 brevirostris, 784 breweri, 273 brewsteri, 211. The dubious Linota brewsteri may be a hybrid between Ægiothus linaria and Chrysomitris pinus. See Brewst., Bull. Nutt. Club, vi, no. 4, Oct. 1881, p. 225 brunneicapillus, 63 Bubo, 462 buccinator, 688 Budytes, 87 buffoni, 767 bullocki, 327 bulweri, 820 Buteo, 512 Butorides, 663

cabanisi, 424 cachinnans, 775 cærulea. See cœrulea Calamospiza, 286 calendula, 33 Calidris, 627 calliope, 417 Callipepla, 577 Calothorax, 418 calurus, 517 Calypte, 415 Campephilus, 431 Camptolæmus, 729 Campylorhynchus, 63 Canace, 555 canadensis, 59, 149 canagica, 698 candicans, 501

candissima, 659 caniceps, 265 cantiaca, 796 cantianus, 591 canus, 779 canutus, 626 capense, 818 capitalis, 362 carbo, 873 Cardellina, 150 Cardinalis, 299 carolina, 450, 679 carolinensis, 16 carpalis, 257 Carpodacus, 194 caspia, 793 cassini, 178 castanea, 123 Catharista, 538 Cathartes, 537 Catherpes, 66 caudacutus, 240 caurinus, 342 cedrorum, 166 celata, 107 Centrocercus, 560 Centrophanes, 220 Centurus, 450 Ceratorhina, 857 Certhia, 62 Certhiola, 153 cerviniventris, 420 Cervle, 423 Chætura, 405 Chamæa, 39 Chamæpelia, 547 Charadrius, 581 Chaulelasmus, 711 Chen, 694 cheriway, 535 Chloëphaga, 698 chlorurus, 310 Chondestes, 281. P. 57, penultimate line, for chondrestes? read chondredestes? Chordediles, 399 Chroïcocephalus, 786 chrysaëtus, 532 chrysoïdes, 458 chrysolæma, 84 Chrysomitris, 212 chrysoparia, 115 chrysoptera, 102 cincinnatiensis, 101. thophaga cincinnatiensis may

be a hybrid between H. pinus and Oporornis formosa. See Ridg., Bull. Nutt. Club, v, 1880, p. 237 cincinnatus, 752 Cinclus, 30 cinctus, 52. Parus cinctus is not in the orig. ed. Since ascertained to occur in Alaska. See Bull. Nutt. Club, ii, no. 1, Jan. 1877, p. 37 cinerescens, 375 cinereus, 22, 250 circumcinetus, 588 Circus, 489 ciris, 292 cirrata, 856 Cistothorus, 81 citrea, 95 Clangula, 725 clypeata, 718 Coccygus, 428 cœrulea, 36 cœrulescens, 117 Colaptes, 457 collaris, 722 colubris, 409 Columba, 539 columba, 872 columbarius, 505 columbianus, 344 Colymbus, 840 confinis, 3, 233 conspersus, 67 Contonus, 380 Conurus, 460 cooperi, (J. G.) 156 cooperi, (Wm.) 495 corax, 338 corniculata, 853 cornutus, 848 coronata, 119, 279 Corvus, 338 costæ, 415 Cotile, 163 Coturniculus, 234 coturniculus, 682 coturnix, 579 couchi, 372 couesi, 621 craverii, 869 creatopus, 833 crecca, 714 crepitans, 673

Crex, 683

crinitus, 373 crissalis, 25, 308 cristata, 349 cristatellus, 859 Crotophaga, 425 cryptoleucus, 339 cucullatus, 328, 745 cunicularia, 487 cupido, 563 Cupidonia, 563 curonicus, 590 curvirostra, 199 curvirostris, 19 cyanea, 295 Cyanecula, 31 cyaneus, 489 cyanocephalus, 332 Cyanocitta, 349 cyanoptera, 717 Cygnus, 688 Cymochorea, 823 Cyrtonyx, 578

dactylisonans, 579 Dafila, 710 Daptium, 818 delawarensis, 778 Dendrocygna, 705 Dendræca, 111 derbianus, 364 Dichromanassa, 661 difficilis, 389. Empidonax flaviventris difficilis was not recognized in the orig. ed. dilophus, 751 Diomedea, 810 discolor, 127 discors, 716 Dolichonyx, 312 domesticus, 74 dominica, 129 dominicensis, 369 dorsalis, 266 dougalli, 800 dresseri, 734

eburnea, 785 Ectopistes, 543 egretta, 658 Elanoïdes, 493 Elanus, 492 elegans, 521, 676 Embernagra, 311 Empidonax, 384 Engyptila, 542 enucleator, 190 ěrēmoē/că, 256 bis. Peucæa r.e., Brown, Bull. Nutt. Club, vii, Jan. 1882, p. 26, Texas. (?)— Gr. έρημος, a desert; οἰκέω, Ι inhabit Eremophila, 82 Ereunetes, 612 Erismatura, 741 erythrina, 540 erythrocephalus, 453 erythrocercus, 374 erythrophthalmus, 301 Eudocimus, 651 Eugenes, 408 Eurynorhynchus, 884 excubitorides, 188 exilipes, 210 exilis, 667

falcinellus, 649 Falco, 497 fallax, 245 familiaris, 62 fasciata, 39, 244 ferina, 723 ferrugineus, 331, 485 fischeri, 732 flammeolus, 471 flammeus, 461 flaviceps, 56 flavifrons, 176 flavipes, 634 flavirostris, 211 flaviventris, 388 flaviviridis, 171 flavus, 87 Florida, 662 floridanus, 341 fœda, 628 forficatus, 367 formicivorus, 454 formosa, 140 forsteri, 798 franklini, 556 Fratercula, 853 Fregetta, 829 frontalis, 353, 196 frugivorus, 340 fulgens, 408 Fulica, 686 fulicarius, 604 fuliginosa, 559 Fuligula, 720 Fulmarus, 814

hæsitata, 819

fulvifrons, 392 fulvigula, 709 fulvus, 582 fumifrons, 360 funerea, 480 furcata, 791, 826 fuscescens, 7 fuscicœrulescens, 511 fuscocaudata, 419 gairdneri, 441 galbula, 326 galeata, 684 Gallinago, 607 Gallinula, 684 gallipavo, 553 gambeli, 278 Garzetta, 659 Geococcyx, 427 Geothlypis, 141 Geotrygon, 550 gigantea, 813 gilvus, 174. Should probably stand as Vireo ochroleucus, Coues glacialis, 728 glottis, 635 glaucescens, 770 Glaucidium, 484 glaucium, 725 glaucus, 492, 768 gnoma, 484 graciæ, 128 grallaria, 829 gramineus, 232 grammicus, 281 griseigena, 847 griseinucha, 205 griseus, 609 gruberi, 529 Grus, 668. See Müller, Nat. Syst. Suppl., 1776, p. 110; Cass., Pr. Phila. Acad., 1864, p. 245 grylle, 871 guarauna, 650 gubernator, 317

hæmastica, 629 Hæmatopus, 595

Gymnocitta, 345

gymnostoma, 672

Guiraca, 291 gularis, 887

guttatus, 231

Haliaëtus, 533 haliaëtus, 530 Halocyptena, 821 hammondi, 390 Harelda, 728 harlani, 515 Harporhynchus, 17 harpyia, 531 harrisi, 439 heermanni, 248 Helmintherus, 96 Helminthophaga, 98. According to Ridgway, Bull. Nutt. Club, vii. no. 1, Jan. 1882, p. 53, the name Helminthophaga is preoccupied, and it has consequently been changed by him to Helminthophila heloisæ, 416 helvetica, 580 henryi, 400 henslowi, 236 hepatica, 157 Herodias, 658 herodias, 655 Hesperocichla, 5 Hesperophona, 189 Heteroscelus, 642 hiaticula, 589 hiemalis, 76 Himantopus, 601 himantopus, 611 Hirundo, 159 hirundo, 797 Histrionicus, 730 holbælli, 208 homochroa, 825 hornbyi, 827 hornemanni, 209 horreorum, 159 hudsonicus, 49, 645 hutchinsi, 704 huttoni, 182 Hydranassa, 660 Hydrochelidon, 806 hvemalis, see hiemalis Hylocichla, 6 Hylotomus, 432 hyperboreus, 603 hypochrysea, 133 hypogæa, 487 hypoleucus, 868

Iache, 421

Icteria, 144 icterocephalus, 319 Icterus, 323 Ictinia, 491 ignea, 300 iliaeus, 4, 282. Turdus iliaeus is not in the orig. ed. Greenland. Reinh., Ibis, 1861, p. 6 illinoensis, 252 imberbe, 393 impennis, 878 inca, 549 incanus, 642 inornatus, 41 intermedia, 277 interpres, 598 Ionornis, 685 Iridoprocne, 160 islandicus, 500

jamaïcensis, 681 Junco, 261

kennicotti, 466 kirtlandi, 131 kittlitzi, 867 kotzebuii, 783 krideri, 519 kuhli, 831

labradorius, 729 Lagopus, 568 lagopus, 525 lapponicus, 220 lariformis, 806 Larus, 768 latirostris, 421 lawrencii, 99. lawrencii may between H.

lawrencii, 99. Helminthophaga lawrencii may be a hybrid between H. pinus and H. chrysoptera. See Brewst. Bull. Nutt. Club, vi, no. 4, Oct. 1881, p. 218.

lecontii, 24 lecontii, 237

leucobronchialis, 100. Helminthopaga leucobronchialis may be a hybrid between H. pinus and H. chrysoptera. See Brewst., Bull. Nutt. Club, vi, no. 4, Oct. 1881, p. 218 leucocephalus, 534

leucocephalus, 534 leucogaster, 72, 747 leucolæma, 83 leucoparia, 703

leucophrys, 276 leucopsis, 699 leucoptera, 198, 546 leucorrhoa, 823 Leucosticte, 201 leucurus, 570 Limosa, 628 linaria, 207 lincolni, 242 lineatus, 520 Linota, 211 litoralis, 204 Lobipes, 603 loculator, 648 Lomvia, 874 longicauda, 145 longirostris, 18 Lophophanes, 40 Lophortyx, 575 Loxia, 198 lucasanus, 436 luciæ, 104 lucifer, 418 ludovicianus, 68, 187 lunifrons, 162 luteiventris, 365 lutescens, 108 luxuriosa, 358

maccalli, 468 maccowni, 223 macgillivrayi, 143 Machetes, 639 macrolopha, 352 Macrorhamphus, 609 macrurus, 333, 799 macularius, 638 maculata, 616 maculosa, 125 magna, 320 major, 334, 832 Mareca, 712 marila, 720 marinus, 771 maritimus, 238 marmoratus, 866 martinica, 550, 685 maruetta, 678 massena, 578 maxima, 794 maxwellæ, 467 media, 607 megalonyx, 305 megarhyncha, 285

melæna, 824

melancholicus, 372 Melanerpes, 453 melanocephalus, 599, 290 melanoleucus, 633 melanotis. 55. Psaltriparus 1997 melanotishas lately been definitely ascertained to occur in Arizona melanura, 37. According to Brewster, Polioptila melanura should read P.californica. See Bull. Nutt. Club, vi, p. 103 melanurus, 830 Meleagris, 553 melodus, 587 Melopelia, 546 Melospiza, 242 merganser, 744 Mergus, 743 meridionalis, 879 mesoleucus, 306 mexicana, 28 miamiensis, 69. Thryothorus ludovicianus miamiensis is not in the orig. ed. Since described from Florida. Am. Nat., ix, 1875, p. 469 Micrathene, 486 . Micropalama, 611 microsoma, 821 migratorius, 1 Milvulus, 366 Mimus, 15 minimus, 53, 387 minor, 605, 401 minutilla, 614 minutus, 730 mitratus, 146 Mitrephorus, 392. This generic term being preoccupied in Coleoptera, Mitrephanes is substituted (Gr. μίτρη, mitre, φαίνω, I appear). See Coues, Bull. Nutt. Club, vii, no. 1, Jan. 1882, p. 55 Mniotilta, 91 mollissima, 733 Molothrus, 313 monocerata, 857 montanus, 14 monticola, 268 moreleti, 296 morio, 346 Motacilla, 86

motacilla, 138

mugitans, 666 musicus, 690 mustelinus, 6 Myeteria, 654 Myiadestes, 169 Myiarchus, 373 Myiodioetes, 146 Myiodynastes, 365

nævius, 5, 136 nanus, 10 nebulosa, 476 neglectus, 51. Parus rufescens neglectus is not in the orig. ed. Since described by Ridgway, Pr. Nat. Mus., i, 1879, p. 485. California nelsoni, 241 Neocorys, 90 Nephœcetes, 404 Nettion, 715 nevadensis, 260 niger, 404 nigrescens, 116, 237 nigricans, 378, 701, 863 nigrilora, 94 nigripes, 811 nitens, 167 nivalis, 219 nivosus, 591 Nomonyx, 742 notabilis, 137 notatus, 218 noveboracensis, 181 nuchalis, 447 Numenius, 643 nuttalli, 348 Nyctala, 482 Nyctea, 479 Nycterodius, 665 Nyctiardea, 664 Nyctidromus, 395

obscurus, 314 obsoletus, 65, 499, 674 occidentalis, 46, 113, 478 oceanicus, 828 Oceanicus, 828 Oceanodroma, 826 ochropus, 636 Œdemia, 737 Œnanthe, 26 Œstrelata, 819 olivaceus, 170 Onychotes, 529 opisthomelas, 836 Oporornis, 139 oregonus, 263 ornatus, 222 Ornithium, 393 Orortyx, 574 Oroscoptes, 14 Ortalis, 552 Ortyx, 571 oryzivorus, 312 Ossifraga, 813 ostrilegus, 595

pacificus, 77, 464 Pagophila, 785 palliatus, 596 pallida, 272 pallidicineta, 564 palmarum, 132 palmeri, 20 paludicola, 80 palustris, 79, 243 Pandion, 530 Panyptila, 403 parasiticus, 766 parisiorum, 329 parkmani, 75 Parra, 672 Parula, 93 Parus, 44 Passer, 192 Passerculus, 224 Passerella, 282 Passerina, 292 passerinus, 234, 547 pealii, 504 Pediœcetes, 561 pelagica, 822 pelasgica, 405 Pelecanus, 748 Pelidna, 623 penelope, 712. Qu. penelops penicillatus, 755 pennsylvanica, 124 peregrina, 109, 503 Perisoreus, 359 perpallidus, 235 perspicillata, 739, 756 pertinax, 381 Petrochelidon, 162 Peucæa, 251 Peucedramus, 110

phæopus, 644

Phaëthon, 762

Phaïnopepla, 167

Phalacrocorax, 750 Phalænoptilus, 398 Phalaropus, 604 phasianellus, 561 philadelphia, 142 philadelphicus, 173 Philohela, 605 Phœbetria, 812 phœniceus, 316 Phœnicopterus, 687 Phonipara, 297 Phylloscopus, 32 Pica, 347 Picicorvus, 344 Picoïdes, 443 pieta, 151, 221, 574 Picus, 433 pileatus, 432 pileolatus, 148 Pinicola, 190 pinus, 98 Pipilo, 301 Pitangus, 364 plagata, 527 platycercus, 413 Plectrophanes, 219 Plegadis, 649 Plotus, 760 plumbea, 38 pluvialis, 583 Podasocys, 592 Podicipes, 848 podicipes, 852 Podilymbus, 852 Polioptila, 36 Polyborus, 535 polyglottus, 15 pomatorhinus, 765 Poœcetes, 232 popetue, 399 Porzana, 678 pratensis, 88, 670 pratincola, 461 princeps, 225 principalis, 431 Priocella, 817 Priofinus, 830 Procellaria, 822 Progne, 165 propinquus, 2. Turdus migratorius propinquus is not in

the orig. ed.

scribed by Ridgway, Bull.

Nutt. Club, ii, no. 1, Jan.

1877, p. 9. Western U. S.

Since de-

Protonotaria, 95 psaltria, 215 Psaltriparus, 53 Pseudogryphus, 536 Psilorhinus, 346 psittaculus, 858 ptilocnemis, 622 Ptychorhamphus, 862 pubescens, 440 Puffinus, 831 pugnax, 639 purpureus, 194 pusilla, 60 pygmæa, 61 Pyranga, 154 Pyrocephalus, 394 Pyrrhula, 191. The difficulty with P. cassini may be owing to wrong sexing of the single known specimen Pyrrhuloxia, 298

Querquedula, 713 querula, 280 Quiscalus, 333

Rallus, 673 Recurvirostra, 600 redivivus, 23 Regulus, 33 resplendens, 759 rhodocolpus, 197 Rhodostethia, 789 Rhyacophilus, 636 Rhynchophanes, 223 Rhynchops, 809 richardsoni, 383 riparia, 163 Rissa, 782 rodgersi, 816 rosea, 653, 789 rossi, 697 rostratus, 230 Rostrhamus, 490 ruber, 448, 652 rubida, 741 rubineus, 394 rubra, 154 rubrifrons, 150 rufescens, 50 ruficapilla, 106 ruficeps, 255 rufina, 247, 886 rufovirgata, 311 rufus, 17

rupestris, 569 rustica, 347 rusticula, 606 ruticilla, 152

sabinii (J.), 567 sabinii (E.), 790 sacer, 498 Salpinetes, 65 samuelis, 249 sancti-johannis, 525 sandvicensis, 226 satrapa, 34 saturatus, 675 savana, 227 saxatilis, 403 Saxicola, 26 sayi, 377 Sayiornis, 377 scalaris, 434 scandiaca, 479 Scardafella, 549 schistacea, 284 Scolecophagus, 331 scolopaceus, 610 Scolopax, 606 Scops, 465 Scotiaptex, 474 Selasphorus, 411 semipalmatus, 586 seniculus, 430 septentrionalis, 45 serrator, 744 serripennis, 164 Setophaga, 151 Sialia, 27 sialis, 27 Simorhynchus, 858 sinuata, 298 Sitta, 57 Siurus, 135 skua, 764 smithsonianus, 773 sociabilis, 490 solitarius, 177, 637 Somateria, 731 sparverioïdes, 510 sparverius, 509 spectabilis, 736 Spectyto, 487 Spermophila, 296 Sphyropicus, 446 spilurus, 73 Spiza, 287

Spizella, 268

sponsa, 719 spraguii, 90 spurius, 324 squamata, 577 Squatarola, 580 Starnænas, 551. Ital. starna, a partridge Steganopus, 602 Stegidopteryx, 164 stellaris, 81 stelleri, 350, 731 Stellula, 417 Stercorarius, 764 Sterna, 792 stolidus, 808 streperus, 711 Strepsilas, 598 striata, 122 striatulus, 497 stricklandi, 437 Strix, 474 Sturnella, 320 Sturnus, 363 subarquatus, 625 subcœrulea, 491 subis, 165 subviridis, 384 suckleyi, 506 suecica, 31 Sula, 746 sulcirostris, 426 superciliaris, 801 Surnia, 480 swainsoni, 13, 97 Symphemia, 632 Synthliboramphus, 865, pessime,

Tachycineta, 161 Tachypetes, 761 taïtensis, 647 Tantalus, 648 Telmatodytes, 79 tengmalmi, 482 tenuirostris, 817, 839 tephrocotis, 203 Tetrao, 563 texensis, 402 thalassina, 161 Thrasyaëtus, 531 Thryothorus, 68 thyroïdes, 449 tigrina, 126 torquatus, 456

Synthliborhamphus, 864

lege

Totanus, 633 townsendi, 114, 169 trachyrhynchus, 748 trailli, 385 trichas, 141 trichopsis, 470 tricolor, 318 tridactyla, 782 Tringa, 626 Tringoïdes, 638 tristis, 213 Trochilus, 409 Troglodytes, 74 Trogon, 422 troile, 874 trowbridgii, 740 trudeaui, 802 Tryngites, 641 Turdus, 1. Tyrannus, 368 tyrannus, 366

ultramarina, 357 ulula, 481 umbella, 565 umbelloïdes, 566 umizusume, 865 unalascæ, 8 unalascensis, 283 unicinetus, 512 Uria, 871 urophasianus, 560 uropygialis, 452 Urubitinga, 528. U. anthracina is not in the orig. ed. Since discovered in Arizona by H. W. Henshaw, Am. Sportsm., v, Feb. 1878, p. 328 ustulatus, 11

Utamania, 877

vallisneria, 724 Vanellus, 593 varia, 91 varius, 446 vauxi, 406 vermivorus, 96 versicolor, 293 verticalis, 370 vespertina, 189 vicinior, 180 villosus, 438 violaceus, 665, 758 virens, 112, 144 Vireo, 170

virescens, 663 virgata, 594 virginiæ, 105 virginiana, 299 v-nigra, 735 vociferans, 371 vociferus, 397, 584 vulgaris, 363 whitneyi, 486 wilsonius, 585 wilsonianus, 472 wollweberi, 43 woodhousii, 355

Xanthocephalus, 319 Xanthura, 358 xantusi, 407 Xema, 790 Xenopicus, 442

Zamelodia, 289 Zenaida, 545 Zenaidura, 544 Zonotrichia, 275

### APPENDIX.

# CATALOGUE OF THE AUTHOR'S ORNITHOLOGICAL PUBLICATIONS, 1861–1881.

\* Anonymous.

† REVIEWS.

#### 1861.

- \*Ornithology. Washington Described, pp. 24-27. (12mo. Washington, Philp & Solomons, 1861.)
   A slight sketch of the Birds of the District of Columbia.
- 2. A Monograph of the Tringeæ of North America. Proc. Acad. Nat. Sci. Phila., xiii, July, 1861, pp. 170-205.
- 3. Notes on the Ornithology of Labrador. Proc. Acad. Nat. Sci. Phila., xiii, August, 1861, pp. 215–257.
- 4. A Monograph of the Genus Ægiothus, with descriptions of new Species. *Proc. Acad. Nat. Sci. Phila.*, xiii, November, 1861, pp. 373–390.

#### 1862.

- 5. List of Birds ascertained to inhabit the District of Columbia, with the times of Arrival and Departure of such as are non-residents, and Brief Notices of Habits, etc. By Elliott Coues and D. Webster Prentiss. Sixteenth Ann. Rep. Smiths. Inst., for 1861, 1862, pp. 399-421.
- Synopsis of the North American Forms of the Colymbide and Podicipide. —
   Proc. Acad. Nat. Sci. Phila., xiv, April, 1862, pp. 226-233.
   Abstract of a monograph published in full in Birds of the Northwest, 1874.
- Revision of the Gulls [Larinæ] of North America; based upon specimens in the Museum of the Smithsonian Institution. — Proc. Acad. Nat. Sci. Phila., xiv, June, 1862, pp. 291–312.

Abstract of a monograph published in full in the Birds of the Northwest, 1874, pp. 589-717.

- Supplementary Note to a "Synopsis of the North American Forms of the Colymbide and Podicepidee."— Proc. Acad. Nat. Sci. Phila., xiv, September, 1862, p. 404.
- 9. A Review of the Terns [Sterninæ] of North America. Proc. Acad. Nat. Sci. Phila., xiv, December, 1862, pp. 535-559.

- Additional Remarks on the North American Ægiothi. Proc. Acad. Nat. Sci. Phila., xv, February, 1863, pp. 40, 41.
- On the Lestris richardsoni of Swainson; with a Critical Review of the Subfamily Lestridine. Proc. Acad. Nat. Sci. Phila., xv, May, 1863, pp. 121-138.
- 12. [On the specific validity of Larus smithsonianus.] *The Ibis*, v, July, 1863, p. 367.

#### 1864.

- The Crania of Colymbus torquatus and C. adamsii compared.—Proc. Acad. Nat. Sci. Phila., xvi, February, 1864, pp. 21, 22.
- 14. A Critical Review of the Family Procellaridæ; Part I., embracing the Procellarieæ, or Stormy Petrels. — Proc. Acad. Nat. Sci. Phila., xvi, March, 1864, pp. 72-91.
- A Critical Review of the Family Procellaridæ: Part II.; Embracing the Puffineæ. — Proc. Acad. Nat. Sci. Phila., xvi, April, 1864, pp. 116-144.
- Notes on certain Central-American Laridæ, collected by Mr. Osbert Salvin and Mr. F. Godman. — The Ibis, vi, July, 1864, pp. 387–393.

#### 1865.

- 17. Ornithology of a Prairie-Journey, and Notes on the Birds of Arizona. *The Ibis*, 2d ser., i, April, 1865, pp. 157-165.
- [Notes on Birds observed at Fort Whipple, Arizona.] The Ibis, 2d ser.,
   i, October, 1865, pp. 535-538.

#### 1866.

- Field Notes on Lophortyx Gambeli. The Ibis, 2d ser., ii, January, 1866, pp. 46-55.
- 20. List of the Birds of Fort Whipple, Arizona: with which are incorporated all other species ascertained to inhabit the Territory; with brief critical and field Notes, descriptions of new species, etc. Proc. Acad. Nat. Sci. Phila., xviii, March, 1866, pp. 39-100.

Fifty copies reissued, repaged, under the title: [Reprinted from the Proceedings of the Philadelphia Academy of Natural Sciences, January 1866.] | — | Prodrome of a Work |

- on the | Ornithology | of | Arizona Territory. | By Elliott Coues, M.A., M.D. | (Ass't Surgeon U. S. Army.) | | Philadelphia: | Merrihew & Son, Printers. | 1866. 8vo. pp., 1-64.

  Note. Part I. of the work here forecast pub. 1877 under the title: Birds of the Colorado Valley, etc.
- A Critical Review of the Family Procellariidæ: Part III; embracing the Fulmareæ. — Proc. Acad. Nat. Sci. Phila., xviii, March, 1866, pp. 25-33.
- Critical Review of the Family Procellariide: Part IV; Embracing the Æstre-lateæ and Prioneæ. Proc. Acad. Nat. Sci. Phila., xviii, May, 1866, pp. 134-172.
- 23. Critical Review of the Family Procellariidæ; Part V; embracing the Diomedeinæ and the Halodrominæ. With a General Supplement. Proc. Acad. Nat. Sci. Phila., xviii, May, 1866, pp. 172-197.
- 24. From Arizona to the Pacific. The Ibis, 2d ser., ii, July, 1866, pp.259-275.
  Notes on the birds observed during the journey.
- 25. The Osteology of the Colymbus torquatus; with Notes on its Myology.—

  Mem. Bost. Soc. Nat. Hist., i, pt. ii, November, 1866, pp. 131-172, figg. 2, pl. 5.

Separately reissued, under the title: On the | Osteology and Myology | of | Colymbus torquatus. | By Elliott Coues, A.M., M.D., | Brevet Captain and Assistant Surgeon United States Army. | [From the Memoirs read before the Boston Society of Natural History, vol. i, part ii.] | Cambridge: | printed at the Riverside Press. | November, 1866. 4to. pp. 131-172, figg. 2, pl. 5.

#### 1867.

26.\*†The Birds of New England. — The Round Table, No. 140, Sept. 28, 1867, pp. 213, 214.

E. A. Samuels' work.

#### 1868.

27. A Monograph of the Aleidæ. — Proc. Acad. Nat. Soc. Phila., xx, January, 1868, pp. 2-81, figg. 1-16.

Also separate, retitled: A Monograph | of | the Alcidæ. | By | Elliott Coues, A.M., M.D. | (Ass't. Surgeon U. S. Army.) | — | Philadelphia: | Merrihew & Son, printers, | No. 243 Arch Street, | 1868. Svo, pp. and figg. as above. See also under 1870.

- List of Birds collected in Southern Arizona by Dr. E. Palmer; with remarks.
   Proc. Acad. Nat. Sci. Phila., xx, January, 1868, pp. 81–85.
- 29. Synopsis of the Birds of South Carolina. Proc. Bost. Soc. Nat. Hist., xii, October 7, 1868, pp. 104-127.
- 30. Catalogue of the Birds of North America contained in the Museum of the Essex Institute; with which is incorporated A List of the Birds of New England. With Brief Critical and Field Notes. Proc. (Comm.) Essex Inst., v, 1868, pp. 249-314.

Fifty copies reissued, with new index, repaged, retitled: A List | of the | Birds of New England. | By Elliott Coues, | Asst. Surgeon, U. S. A. | — | [Reprinted from the

- Proceedings of the Essex Institute, Vol. v.] | | Salem, Mass. | Essex Institute Press. | 1868. 8vo. pp. 1–71.
- 31. Instances of Albinism among our Birds. Amer. Nat., ii, No. 3, May, 1868, pp. 161, 162.

About a dozen cases, chiefly of North American species.

32. Bird's-Eye Views. — Amer. Nat., ii, No. 10, December, 1868, pp. 505-513;
ii, No. 11, January, 1869, pp. 571-583, figg.
On the structure of the eye in Birds.

#### 1869.

- 33. On a Chick with supernumerary Legs. Proc. Bost. Soc. Nat. Hist., May 19, 1869, pp. 78-82.
- 34. \*Of Doves and Thorns. *The Liberal Christian*, July 24, 1869.

  Breeding of Zenaidura carolinensis.
- 35. \*Of a "Fast" Bird [Geococcyx californianus].—The Liberal Christian, Aug. 14, 1869.
- 36. \*A Skeleton in the House. The Liberal Christian, Sept. 11, 1869.
  Molothrus ater × Polioptila cœrulea.
- 37. Sea-side Homes. *Amer. Nat.*, iii, No. 7, September, 1869, pp. 337–349. On the breeding of Sterna antillarum and Ægialites wilsonius.
- On Variation in the Genus Ægiothus. Proc. Acad. Nat. Sci. Phila., xxi, October, 1869, pp. 180–189.
   Supplementary to the article in op. cit., 1861, p. 373.
- 39. \*Structure of Feathers. The Liberal Christian, Oct. 9, 1869.
- On the Classification of Water Birds. Proc. Acad. Nat. Sci. Phila., xxi, December, 1869, pp. 193–218.

Reissued, repaged. On the | Classification | of | Water Birds. | By Elliott Coues, A. M., M. D., Ph. D., | [etc., 4 lines.] | — | [Reprinted from the Proceedings of the Philadelphia Academy of Natural Sciences | for December, 1869.] | — | Philadelphia: | Merrihew & Son, printers. | No. 243 Arch Street. | 1870. 8vo. pp. 28.

- Extracts from a Memoir intituled 'A Monograph of the Alcide.'—Zoologist,
   2d ser., v, 1870, pp. 2004–2016, 2081–2090, 2124–2132, 2155–2163, 2205–2214, 2245–2253, 2289–2296, 2327–2334, 2369–2378, 2396–2403.
   Reprinted, modified, from Proc. Phila. Acad., 1868, pp. 2-81.
- 42. The Clapper Rail [Rallus crepitans]. Amer. Nat., iii, No. 11, January, 1870, pp. 600-607.
- 43. The Great Auk [Alca impennis].—*Amer. Nat.*, iv, No. 1, March, 1870, p. 57.

- The Cow Bird [Molothrus ater]. Amer. Nat., iv, No. 1, March, 1870,
   p. 58.
- 45. Foot-notes from a Page of Sand. Amer. Nat., iv, No. 5, July, 1870, pp. 297–303.
  Tracks made by Sandpipers, &c.
- **46.** The Natural History of Quiscalus major. *The Ibis*, 2d ser., vi, July, 1870, pp. 367-378.
- 47. †Ornithological Results of the Exploration of the North-west. Amer. Nat., iv, No. 6, August, 1870, pp. 367-371.

Review of Dall and Bannister's and Baird's papers on the Birds of Alaska, in *Trans. Chicago Acad.*, 1869.

#### 1871.

- **48.** Notes on the Natural History of Fort Macon, N. C., and Vicinity. (No. 1.) [Vertebrates.] *Proc. Acad. Nat. Sci. Phila.*, xxiii, May 2, 1871, pp. 12–49.
  - Birds, pp. 18-47. There are 5 Nos. of this, 1871-1879, the 1st and 4th relating to Birds.
- 49. The Yellow-headed Blackbird [Xanthocephalus icterocephalus].—Amer. Nat.,
  v, No. 4, June, 1871, pp. 195-200, fig.
  Biography of the species, with references to other birds observed in Kansas.
- 50. †Recent Ornithological Publications. Amer. Nat., v, No. 4, June, 1871, pp. 234–238.

Bannister on Anserinæ, Pr. Phila. Acad., 1870, p. 130; Ridgway on Falconidæ, ibid., p. 138.

- †Progress of American Ornithology. Amer. Nat. v, No. 6, August, 1871, pp. 364-373.
   Review of J. A. Allen's memoir on Florida Birds, Bull. Mus. Comp. Zool., 1871.
- Mechanism of Flexion and Extension in Birds' Wings. Amer. Nat., v,
   Nos. 8 and 9, September, 1871, pp. 513, 514.
   Abstract of the paper in the Proc. Amer. Assoc. Adv. Sci. for 1871.
- Bullock's Oriole [Icterus bullocki]. Amer. Nat. v, No. 11, November, 1871, pp. 678-682, fig. 120.
- **54.** Singular Albino [Dolichonyx oryzivorus]. Amer. Nat., v, No. 11, November, 1871, p. 733.
- 55. The Long-crested Jay [Cyanocitta macrolopha]. Amer. Nat., v, No. 12, December, 1871, pp. 770-775, fig.
- 56. †Gray's Hand List of Birds. Amer. Nat., v, No. 12, December, 1871, pp. 775-779.

#### 1872.

 Mechanism of Flexion and Extension in Birds' Wings. — Proc. Amer. Assoc. Adv. Sci., xx, for 1871, 1872, pp. 278–284, figg.

- 58. Observations on Picicorvus Columbianus. *The Ibis*, 3d ser., ii, January, 1872, pp. 52-59.
- Ornithological Query [Turdus migratorius]. Amer. Nat., vi, No. 1, January, 1872, p. 47.
- 60. †Two Late American Papers on Ornithology. Amer. Nat., vi, No. 3, March, 1872, pp. 165, 166.
  - Ogden on Chettusia, Pr. Phila. Acad., 1871, p. 191; Lawrence on New Troglodytidæ and Tyrannidæ, ibid., p. 233.
- 61.\*†An Ornithological Blunder [Bonasa jobsii]. Amer. Nat., vi, No. 3, March, 1872, pp. 172, 173.
- Contribution to the History of the Blue Crow [Gymnokitta cyanocephala] of America. — The Ibis, 3d ser., ii, April, 1872, pp. 152–158.
- 63. †Newton's Ornithological Register. Amer. Nat., vi, No. 6, June, 1872, pp. 360, 361.
- 64. The Nest, Eggs, and Breeding Habits of Harporhynchus crissalis. Amer. Nat., vi, No. 6, June, 1872, pp. 370, 371.
- 65. A New Bird [Glaucidium ferrugineum] to the United States. Amer. Nat., vi, No. 6, June, 1872, p. 370.
- Studies of the Tyrannidæ. Part I. Revision of the Species of Myiarchus.
   Proc. Acad. Nat. Sci. Phila., xxiv, June 25-July 16, 1872, pp. 56-81.
- 67. †The Boston Society's Ornithological Catalogue. Amer. Nat., vi, No. 8, August, 1872, pp. 472, 473.
  Review of Λ. Hyatt's paper on Spheniscidæ.
- 68. Nest and Eggs of Helminthophaga luciæ. Amer. Nat., vi, No. 8, August, 1872, p. 493.
- 69. Occurrence of Couch's Flycatcher [Tyrannus melancholicus couchi] in the United States. Amer. Nat., vi, No. 8, August, 1872, p. 493.
- 70. †Giebel's Thesaurus. *Amer. Nat.*, vi, No. 9, September, 1872, pp. 549-551.
- 71. Material for a Monograph of the Spheniscidæ. Proc. Acad. Nat. Sci. Phila., xxiv, September, 1872, pp. 170-212, pll. 4, 5.
- 72. †Recent Discoveries in Ornithotomy. Amer. Nat., vi, No. 10, October, 1872, pp. 631–635.
  - Morse on the Carpus and Tarsus of Birds, Ann. Lyc. Nat. Hist., N. Y., 1872; and Reinhardt, Cranial Bone of Musophagidæ, Vidd. Meddel. Nat. Förh.  $Kj\phi b$ ., 1871.
- 73. Key | to | North American Birds | containing a concise account of every species of | Living and Fossil Bird | at present known from the continent north of the Mexican | and United States Boundary. | Illustrated by 6 steel plates, and upwards of 250 woodcuts. | By | Elliott Coues, | Assistant Surgeon United States Army. | | Salem: Naturalists' Agency. | New York: Dodd and Mead. | Boston: | Estes and Lauriat. | 1872. 1 vol. imp. 8vo.

- 4 prel. ll., pp. 1-361, 1 l., pll. 1-6, figg. 1-238. (Pub. Oct. 1872. Ed. of 2,200 cop. Copyright, Putnam and Coues.)
- 74. [Contributions to] Sharpe and Dresser's Hist. of the Birds of Europe, Parts xi, xii, xv, 1872.

- 75. [Contributions to] Sharpe and Dresser's Hist. of the Birds of Europe, Parts xvi, xx, xxi, 1873.
- 76. †Dubois' Conspectus. Amer. Nat., vii, No. 1, January, 1873, pp. 40–42.
  Review of C. F. Dubois' Conspectus Avium Europæarum.
- 77. †New England Ornithology. Amer. Nat., vii, No. 1, January, 1873, pp. 42, 43.
  Review of C. J. Maynard's paper in Pr. Bost. Soc., xiv, 1872, p. 356.
- 78. [Circular relating to the "Birds of the Northwest."] Headquarters Department of Dakota, Feb. 14, 1873.
- †Handbook of British Birds. Amer. Nat., vii, No. 3, March, 1873, pp. 163-165.
   Review of J. E. Harting's work of that name.
- †Ornithology of the West. Amer. Nat., vii, No. 4, April, 1873, pp. 220-223.
   Review of J. A. Allen's paper in Bull. Mus. Comp. Zool., iii, 1872, p. 113.
- 81. †African Ornithology. Amer. Nat., vii, No. 4, April, 1873, pp. 226, 227. Review of J. H. Gurney, Sr.'s, Andersson's Birds of Damara Land.
- 82. Some United States Birds, New to Science, and other Things Ornithological.

   Amer. Nat., vii, No. 6, June, 1873, pp. 321–331, figg. 65–70.
- New Avian Subclass [Odontornithes]. Amer. Nat., vii, No. 6, June, 1873, p. 364.
- Color-variation in Birds Dependent upon Climatic Influences. Amer. Nat.,
   vii, No. 7, July, 1873, pp. 415–418.
   Criticism of R. Ridgway's papers in Am. Journ. Sci., iv, 1872, p. 454; v, 1873, p. 39.
- 85. †Late Local Lists. Amer. Nat., vii, No. 7, July, 1873, pp. 418–421.
  Reviews of W. H. Dall, Pr. Cala. Acad., 1873; C. H. Holden and C. E. Aiken, Pr. Bost. Soc., xv, 1872, p. 193; W. D. Scott, ibid., p. 219.
- 86. "Birds walking under Water." Forest and Stream, Oct. 16, 1873.
- 87. Notes on Two little-known Birds of the United States [Passerculus bairdi, Neocorys spraguii]. Amer. Nat., vii, No. 11, November, 1873, pp. 695-697.
- 88. Use of small shot. Amer. Sportsm., Nov. 22, 1873, p. 117.
- Specimens of Bird Architecture [Icterus]. Amer. Sportsm., Nov. 29, 1873, p. 129.

- 90. Range of the Eared Grebe [Podiceps auritus californicus]. Amer. Nat., vii, No. 12, December, 1873, p. 745.
- 91. Notice of a Rare Bird [Coturniculus lecontii]. Amer. Nat., vii, No. 12, December, 1873, pp. 748, 749.
- 92. Report | on the | Prybilov Group, or Seal Islands, of Alaska. | By | Henry W. Elliott, | Assistant Agent Treasury Department. | | Washington: | Government Printing Office. | 1873. 1 vol. oblong 4to. (Appendix. Ornithology of the Prybilov Islands. By Dr. Elliott Coues, U. S. A.)
  This is the orig. ed., very scarce (150 copies.). See 1875.
- 93. A | Check List | of | North American Birds. | | By | Elliott Coues. | | Salem. | Naturalists' Agency. | 1873. 8vo. 2 prel. ll., pp. 1–137, 2 ll.

  This is the orig. ed. Separately published December, 1873. Also published with "Field Ornithology," 1874.

- 94. Field Ornithology. | Comprising a | Manual of instruction | for | procuring, preparing, and preserving Birds, | and a | Check List of North American Birds. | By | Dr. Elliott Coues, U. S. A. | [Monogram.] | Salem: | Naturalists' Agency. | Boston: Estes & Lauriat. | New York: Dodd & Mead. | 1874. 1 vol. 8vo. pp. i-iv, 1-116, 1-137, 2 ll.
  - Published January, 1874. The "Check List" originally published separately, 1873.
- 95. [On the Classification of Birds, with Characters of the Higher Groups, and Analytical Tables of North American Families.] Baird, Brewer, and Ridgway's Hist. N. A. Birds, i, 1874, pp. xiv-xxviii.
- 96. Glossary of Technical Terms used in Descriptive Ornithology. Including a number of prominent Anatomical and Physiological Terms. Baird, Brewer, and Ridgway's Hist. N. A. Birds, iii, 1874, pp. 535-560.
- 97. Specimens of Bird Architecture [Cinclus mexicanus]. Amer. Sportsm., iii, Jan. 17, 1874, p. 245.
- 98.\*†The Birds of North America. The Nation, No. 447, Jan. 22, 1874, p. 65.
  Review of Theodore Jasper's work.
- 99. Specimens of Bird Architecture [Chætura pelasgica]. Amer. Sportsm., iii, Feb. 14, 1874, p. 313.
- 100. Hybrid Ducks [Anas boscas × Hyonetta moschata]. Forest and Stream, Feb. 19, 1874.
- 101. Hybrid Ducks [Anas boscas × Dafila acuta]. Forest and Stream, March 5, 1874.
- 102. Pet Owls [Bubo virginianus]. Amer. Sportsm., iii, March 7, 1874, p. 354.
- 103. †The New Work on Birds. Amer. Sportsm., iii, March 28, 1874, p. 412.
  Review of Baird, Brewer, and Ridgway's work.

- 104. †Avifauna of Colorado and Wyoming. Amer. Nat., viii, No. 4, April, 1874, p. 240.
- 105. The Snow-bird [Junco hiemalis] as a Sparrow. Field and Stream, April 4, 1874.
  With reference to the construction of a game law.
- Powder-down. Forest and Stream, ii, April 9, 1874, p. 134.
   Luminosity of these feathers in Ardeidæ.
- Specimens of Bird Architecture [hole-breeders]. Amer. Sportsm., iv, April 11, 1874, p. 19.
- 108.\*†North American Ornithology. *The Nation*, No. 460, April 23, 1874, p. 270. 'Review of Baird, Brewer, and Ridgway's work.
- 109.\*†History of North American Birds. Forest and Stream, April 30, 1874,
  p. 179.
  Review of Baird, Brewer, and Ridgway's work.
- 110. Habits and Characteristics of Swainson's Buzzard [Buteo swainsoni].—

  Amer. Nat., viii, No. 5, May, 1874, pp. 282-287.
- 111. †Birds of Illinois. Field and Stream, May 2, 1874.
  Review of R. Ridgway's paper, Ann. Lyc. N. Y., x, 1874, p. 364.
- 112. Small Shot. Reply to "Arrow." Amer. Sportsm., May 30, 1874.
- 113. The Californian Vulture [Cathartes californianus]. Amer. Sportsm., iv, June 13, 1874, p. 160, fig.
- 114. Wild Turkeys [Meleagris gallipavo americana] Grouse. Field and Stream, June 13, 1874.
- 115. Dusky Grouse; Blue Grouse; Pine Grouse [Canace obscura]. Field and Stream, June 27, 1874, p. 154; July 11, 1874, p. 170.
- 116. English Sparrows [Passer domesticus]. Amer. Nat. viii, No. 7, July, 1874, p. 436.
  cf. Bull. U. S. Geol. and Geogr. Surv Terr., v, No. 2, 1879, p. 178.
- 117. Shooting Wood Ibises [Tantalus loculator]. Amer. Sportsm., iv, July 11, 1874, p. 225.
- 118. Plumed Quail [Lophortyx gambeli]. Field and Stream, July 25, 1874, p. 187; Aug. 8, 1874, p. 203; Aug. 22, 1874, p. 224.
- 119. New Species of North American Bird [Tringa ptilocnemis]. Amer. Nat. viii, No. 8, August, 1874, pp. 500, 501.
- 120. The Cranes [Gruidæ] of America. Forest and Stream, iii, Aug. 20, 1874, p. 20.
- 121. The Blue Quail [Callipepla squamata]. Field and Stream, Aug. 29, 1874.

- 122. †Recent Publications in Ornithology. Amer. Nat., viii, No. 9, September, 1874, pp. 541–546.
  - Reviews of several papers, chiefly on N. Am. Birds.
- 123. New Variety of the Blue Grosbeak [Guiraca cœrulea eurhyncha]. Amer. Nat., viii, No. 9, September, 1874, p. 563.
- 124. How to Shoot. Field and Stream, Sept. 26, 1874.
- 125. On the Nesting of Certain Hawks, etc. Amer. Nat., viii, No. 10, October, 1874, pp. 596–603.

Falco communis, Buteo swainsoni, Archibuteo ferrugineus, and other birds of Montana.

- 126.\*†A History of North American Birds. Field and Stream, Oct. 31, 1874.
  Review of Baird, Brewer, and Ridgway's work.
- 127. The Rails Family Rallidæ. Amer. Sportsm., v, Oct. 31, 1874, p. 65.
- 128. The Sparrow [Passer domesticus] War. Amer. Sportsm., v, Nov. 21, 1874, p. 113.
  cf. Bull. U. S. Geol. and Geogr. Surv. Terr., v, No. 2, 1879, p. 178.
- 129. Shells Paper or Brass? Forest and Stream, Dec. 24, 1874.
- 130. Department of the Interior. | United States Geological Survey of the Territories. | F. V. Hayden, U. S. Geologist-in-Charge. | | Miscellaneous Publications No. 3. | | Birds of the Northwest: | A Hand-book | of | The Ornithology | of the | Region drained by the Missouri River | and its Tributaries. | | By Elliott Coues, | Captain and Assistant Surgeon U. S. Army. | | Washington: | Government Printing Office. | 1874. 1 vol. 8vo. pp. i-xii, 1-791.

Pub. Dec. 1874. Ed. of 2,200 copies. 214 copies rebound, reissued, retitled: Birds of the North-West: | a Handbook | of | American Ornithology, | containing accounts of all the birds inhabiting the | Great Missouri Valley, | and many others, together representing a large majority of the | Birds of North America, | with copious biographical details from personal | observation, and an extensive synonymy. | . . . Boston: | Estes & Lauriat, | Salem Naturalists' Agency, | 1877.

131. Monograph of the North American Laridæ.—Birds of the Northwest, December, 1874, pp. 589-717.

- 132. A Report | upon the | Condition of Affairs | in the | Territory of Alaska. |
   | By Henry W. Elliott, | Special Agent Treasury Department. | |
  Washington: | Government Printing Office. | 1875. 1 vol. 8vo. pp. 277.
  (Chap. IX. Ornithology of the Prybilov Islands. By Dr. Elliott Coues,
  U. S. A. pp. 166-212.)
  - Reprinted from the orig. ed., 1873.
- 133. The Fauna | of the | Prybilov Islands | abridged from the | "Report on the Prybilov Group or Seal Islands of Alaska," | by Henry W. Elliott; with an Appendix on the | Ornithology by Dr. Elliott Coues (Washington, 1873).

- | By J. E. Harting, F. L. S. F. Z. S. | London | reprinted from the Natural History columns of | "The Field" for private circulation | 1875 8vo. pp. 38, pl. 1.
- 134. The Sparrows [Passer domesticus]. Amer. Sportsm., v, Jan. 23, 1875, p. 264.
- 135. On the Breeding of certain Birds [of Montana Territory]. Amer. Nat., ix, No. 2, February, 1875, pp. 75-78.
- 136. [On the Nest and Eggs of Gymnocitta cyanocephala.] The Ibis, 3d ser., v, April, 1875, pp. 270, 271.
- 137. Albino Black-Bird [Xanthocephalus icterocephalus]. Rod and Gun, vi, April 10, 1875, p. 24.
- 138. Duck Shooting à Cheval. Amer. Sportsm., April 24, 1875.
- 139. Sparrows [Passer domesticus] more evidence. Rod and Gun, vi, July 17, 1875, p. 249.
- 140. Ornithology of the Transit-of-Venus "Centennial." Forest and Stream, Aug. 19, 1875.
- 141. Fasti Ornithologiæ Redivivi. No. I. Bartram's 'Travels.' *Proc. Acad. Nat. Sci. Phila.*, May-September, 1875, pp. 338-358.
- 142.\*†[Notice of Brewer's Catalogue of the Birds of New England.] N. Y. Independent, Oct. 7, 1875.
- 143. †A late paper on Birds. Amer. Nat., ix, No. 10, November, 1875, pp. 570, 571.
  W. Brewster's, in Ann. Lyc. Nat. Hist., N. Y. xi, 1875, p. 129.
- Contributions | to the | Natural History | of | Kerguelen Island, | made in connection with the American Transit-of-Venus | Expedition, 1874–75. |
  By | J. H. Kidder, M.D., | Passed Assistant Surgeon U. S. Navy. | |
  I. | Ornithology. | Edited by Dr. Elliott Coues, U. S. A. | | Bull. U. S. Nat. Mus., No. 2, 1875, pp. i-ix, 1-51. Washington: Government Printing Office. 1875.

Published November, 1875.

#### 1876.

- 145. [Peucedramus, g. n., Coues MS.] Rep. Expl. W. 100 Merid., Vol. v, 1875, p. 202. (Pub. 1876.)
- 146. On the Breeding-Habits, Nest, and Eggs, of the White-tailed Ptarmigan (Lagopus leucurus). Bull. U. S. Geol. and Geogr. Surv. Terr., 2d ser., No. 5, Jan. 8, 1876, pp. 263–266.

Also published separately. 8vo. Washington, 1876.

147. Bewick's Wren, Thryothorus Bewicki. — Amer. Nat., x, No. 1, January, 1876, p. 48.

- 148. Range of the Bay Ibis [Plegadis falcinellus]. Amer. Nat., x, No. 1, January, 1876, p. 48.
- 149. Coues to "Boone" [on Brant]. Rod and Gun, vii, Jan. 15, 1876, p. 248.
- 150. An Account of the various Publications relating to the Travels of Lewis and Clarke, with a Commentary on the Zoological Results of their Expedition.

   Bull. U. S. Geol. and Geogr. Surv. Terr., 2d ser., No. 6, Feb. 8, 1876, pp. 417–444.

Also separate. 8vo. Washington, 1876.

- 151. Contributions to the Natural History of Kerguelen Island, [etc.]. Oology, etc. By J. H. Kidder and Elliott Coues. Bull. U. S. Nat. Mus., No. 3, February, 1876, pp. 7–20.
- 152. A Study of Chionis minor with reference to its Structure and Systematic Position. By J. H. Kidder, U. S. N., and Elliott Coues, U. S. A.—Bull. U. S. Nat. Mus., No. 3, February, 1876, pp. 85–116.
- 153. Reply to Mr. J. A. Allen's "Availability of certain Bartramian Names in Ornithology." Amer. Nat., x, No. 2, February, 1876, pp. 98–102.
- 154. Breeding Range of the Snow-Bird [Junco hiemalis].—Amer. Nat., x, No. 2, February, 1876, pp. 114, 115.
- 155. Unusual Nesting Sites of the Night Hawk [Chordediles popetue] and Towhee Bunting [Pipilo erythrophthalmus]. Amer. Nat., x, No. 4, April, 1876, p. 239.
- 156. Dr. Coues upon Quail, etc. Rod and Gun, viii, April 1, 1876, p. 9.
- 157. Dr. Coues on Brant, etc. Rod and Gun, viii, April 1, 1876, p. 8.
- 158. †Mr. Gentry's Book about Birds. Rod and Gun, viii, April 29, 1876,
  p. 71.
  Review of Life Histories of the Birds of Eastern Pennsylvania, Vol. I.
- 159. The Labrador Duck [Camptolemus labradorius]. Amer. Nat., x, No. 5, May, 1876, p. 303.
- 160.\*†Life-Histories of the Birds of Eastern Pennsylvania. The Nation, May 4, 1876.
  - Review of Vol. I. of T. G. Gentry's work.
- 161. The European Woodcock [Scolopax rusticula] shot in Virginia. Amer. Nat., x, No. 6, June, 1876, p. 372.
- Notable Change of Habit of the Bank Swallow [i. e., Stelgidopteryx serripennis]. Amer. Nat., x, No. 6, June, 1876, pp. 372, 373.
- 163. Letters on Ornithology. No. 1.—The Oregon Robin [Turdus nævius].—
  Chicago Field, June 24, 1876, fig.

This illustrated series of 30 Letters, running from above date to July, 1879, at various intervals, is in part new, partly from the "Birds of the Northwest."

- 164. Tarsal Envelope in Campylorhynchus and allied Genera. Bull. Nutt. Orn. Club, i, No. 2, July, 1876, pp. 50, 51.
- **165.** Brant once more. *Rod and Gun*, July 8, 1876.
- 166. Letters on Ornithology. No. 2.—The American Tree-Creeper [Certhia familiaris].— *Chicago Field*, Aug. 12, 1876, fig.
- 167. Letters on Ornithology. No. 3.—The Blue-gray Gnatcatcher [Polioptila cœrulea].—Chicago Field, Aug. 26, 1876, fig.
- 168. On the Number of Primaries in Oscines. Bull. Nutt. Orn. Club, i, No. 3, September, 1876, pp. 60-63.
- 169. Letters on Ornithology. No. 4. The Horned or Shore Lark [Eremophila alpestris]. *Chicago Field*, Oct. 7, 1876, fig.
- 170. Peculiar Nesting-Site of the Bank-Swallow [i.e., Stelgidopteryx serripennis].
   Bull. Nutt. Orn. Club, i, No, 4, November, 1876, p. 96.
- 171. Dr. Coues on "Partridge," "Quail," Etc. Rod and Gun, Nov. 11, 1876.
- 172. Letters on Ornithology. No. 5. Marsh Wrens [Telmatodytes palustris, Cistothorus stellaris]. Chicago Field, Nov. 18, 1876, figg.
- 173. \*Anecdote of a Crow's [Corvus maritimus] Intelligence. N. Y. Independent, Nov. 23, 1876.
- 174. The Destruction of Birds by Telegraph Wire. Amer. Nat., x, No. 12, December, 1876, pp. 734-736.
   Copied abridged by the press at large.
- 175. Letters on Ornithology. No. 6.—The Shrike, or Butcher Bird [Lanius borealis].—Chicago Field, Dec. 2, 1876, fig.
- 176.\*†Recent Text-books of Zoology. N. Y. Independent, Dec. 2, 1876.

  Reviews of E. S. Morse's and S. Tenney's works.
- 177. Letters on Ornithology. No. 7.—The Catbird (Mimus carolinensis].—
  Chicago Field, Dec. 9, 1876, fig.
- 178. Letters on Ornithology. No. 8. Nuthatches [Sittidæ]. Chicago Field, Dec. 16, 1876, fig.
- 179.\*†Life-Histories of Animals, including Man. The Nation, No. 369, 1876.
  Review of A. S. Packard's work.
- 180. Letters on Ornithology. No. 9.— The Red-tailed Buzzard [Buteo borealis] and other Hawks.— *Chicago Field*, Dec. 23, 1876, fig.
- 181.\*†The Land-Birds and Game-Birds of New England. The Nation, Dec. 28, 1876.Review of H. D. Minot's work.
- 182. Letters on Ornithology. No. 10.—Titmice, Tomtits, or Chickadees [Paridæ].—Chicago Field, Dec. 30, 1876, figg.

- 183. Note on Podiceps dominicus. Bull. Nutt. Orn. Club, ii, No. 1, January, 1877, p. 26.
- 184. Eastward Range of the Ferruginous Buzzard [Archibuteo ferrugineus]. Bull. Nutt. Orn. Club, ii, No. 1, January, 1877, p. 26.
- 185. Letters on Ornithology. No. 11. Swallows [Hirundinidæ]. Chicago Field, Jan. 6, 1877, figg.
- 186. Letters on Ornithology. No. 12. Woodpeckers [Picidæ]. Chicago Field, Jan. 13, 1877, 6 figg.
- 187.\*†The Land Birds and Game Birds of New England. Forest and Stream, Jan. 25, 1877.
  Review of H. D. Minot's work.
- 188.\*†Land Birds and Game Birds of New England. Rod and Gun, Jan. 27, 1877.

Review of H. D. Minot's work.

- 189. Letters on Ornithology. No. 13.—The Harrier [Circus cyaneus hudsonius].—Chicago Field, Feb. 3, 1877.
- 190. To the Swallow. Rod and Gun, Feb. 3, 1877.
- 191. Letters on Ornithology. No. 13 [bis = 14]. Grasshoppers [in connection with habits of the Sharp-tailed Grouse, Pediocetes phasianellus columbianus]. Chicago Field, March 17, 1877, fig.
- 192. Corrections of Nomenclature in the Genus Siurus. Bull. Nutt. Orn. Club, ii, No. 2, April, 1877, pp. 29–33.
- 193. †Minot's "Birds of New England." Bull. Nutt. Orn. Club, ii, No. 2, April, 1877, pp. 49, 50.
  Review of H. D. Minot's work.
- 194. Western Range of Conurus carolinensis. Bull. Nutt. Orn. Club, ii, No. 2, April, 1877, p. 50.
- Note on the Cinnamon Teal (Querquedula cyanoptera). Bull. Nutt. Orn. Club, ii, No. 2, April, 1877, p. 51.
- 196. Remarks on the Birds of the District of Columbia. By Drs. E. Coues and D. W. Prentiss. Field and Forest, ii, No. 11, May, 1877, pp. 191-193.

  Also in a separate pamphlet entitled: Catalogue of the Birds of the District of Columbia, prepared by Pierre Louis Jouy, with Remarks on the Birds of the District, by Drs. Coues and Prentiss. 8vo. Washington, 1877, pp. 11.
- 197. The Song that the Bluebird Sings. Harper's Magazine, May, 1877, p. 891. Reprinted in many places.
- 198. Notes on the Ornithology of the Region about the Source of the Red River of Texas, from Observations made during the Explorations conducted by Lieut. E. H. Ruffner, Corps of Engineers, U. S. A. By C. A. H.

McCauley, Lieut. Third United States Artillery. Annoted by Dr. Elliott Coues, U. S. A. — Bull. U. S. Geol. and Geogr. Surv. Terr., ii, No. 3, May, 1877, pp. 655-695.

Also separate, new cover-title, same pagination.

- 199. Birds [etc.]. The (Baltimore) Mirror, June 1, July 1, Aug. 1, Sept. 1, Oct. 1, Nov. 1, Dec. 1, 1877.
  - From the "Birds of the Northwest."
- 200. Leptoptila [lege Engyptila] albifrons, a Pigeon new to the United States Fauna. Bull. Nutt. Orn. Club, ii, No. 3, July, 1877, pp. 82-83.
- 201. Melopelia leucoptera in Colorado. Bull. Nutt. Orn. Club, ii, No. 3, July, 1877, p. 83.
- 202. Dr. Coues replies to Dr. Brewer [in the Sparrow (Passer domesticus) controversy]. Washington Gazette, July 8, 1877.
- 203. Letters on Ornithology. No. 15. Curious Crows [Picicorvus columbianus, Gymnocitta cyanocephala]. *Chicago Field*, July 14, 1877, figg.
- 204. Letters on Ornithology. No. 16. The English Sparrow [Passer domesticus]. Chicago Field, July 21, 1877.
  Many reprints elsewhere.
- 205.\*†Our Birds of Prey; or, the Eagles, Hawks, and Owls of Canada. The Nation, 1877, p. 341.
  Review of H. G. Vennor's work.
- Letters on Ornithology. No. 16 [bis]. The American Warblers. Family Sylvicolidæ. Chicago Field, Dec. 15, 1877, fig.

#### 1878.

- 207. Department of the Interior | United States Geological Survey of the Territories | F. V. Hayden, U. S. Geologist-in-Charge | | Miscellaneous Publications No. 11 | | Birds of the Colorado Valley | A Repository of | Scientific and Popular Information | concerning | North American Ornithology | By | Elliott Coues | | ἸΠλθ, ἢλθε χελιδών καλὰς ὧρας ἄγουσα | | Part First | Passeres to Laniidæ | Bibliographical Appendix | Seventy Illustrations | | Washington | Government Printing Office | 1878 8vo. pp. i-xvi, 1-807, figg. 70.
- 208. List of Faunal Publications relating to North American Ornithology. Birds Colorado Valley, Part I, 1878, Appendix, pp. 567-784 or [1]— [218].

This constitutes the First Instalment of Ornithological Bibliography; for 2d, 3d, and 4th, see 1879 and 1880.

- 209. Notes on the Natural History of Fort Macon, N. C., and Vicinity. (No. 4.) By Drs. Elliott Coues and H. C. Yarrow. — Proc. Acad. Nat. Sci. Phila., 1878, pp. 21–28.
  Supplementary to No. 1. Vertebrates, 1871. Pirile for 22, 24. (No. 2, Fisher is by
  - Supplementary to No. 1, Vertebrates, 1871. Birds, pp. 22-24. (No. 3, Fishes, is by Dr. Yarrow.)
- Note on Passerculus bairdi and P. princeps. Bull. Nutt. Orn. Club, iii.
   No. 1, January, 1878, pp. 1–3, pl. col'd.
- 211. The Northern Phalarope [Lobipes hyperboreus] in North Carolina. Bull. Nutt. Orn. Club, iii, No. 1, January, 1878, pp. 40, 41.
- 212. The Willow Grouse [Lagopus albus] in New York. Bull. Nutt. Orn. Club, iii, No. 1, January, 1878, p. 41.
- 213. Pipilo erythrophthalmus with spotted Scapulars. Bull. Nutt. Orn. Club, iii, No. 1, January, 1878, pp. 41–42.
- Melanism in Turdus migratorius. Bull. Nutt. Orn. Club, iii, No. 1, January, 1878, pp. 47, 48.
- 215. The Sparrow [Passer domesticus] Pest. The Country, Jan. 19, 1878.
- 216. Letters on Ornithology. No. 17.—The Aquatic Wood-Wagtail, or New York Water Thrush. (Siurus nævius.)—Chicago Field, Feb. 2, 1878.
- 217. Notes on the Ornithology of the Lower Rio Grande of Texas, from Observations made during the season of 1877. By George B. Sennett. Edited, with Annotations, by Dr. Elliott Coues, U. S. A. Bull. U. S. Geol. and Geogr. Surv. Terr., iv, No. 1, Feb. 5, 1878, pp. 1-66.
  Also separate, new cover-title, same pagination.
- 218. Peculiar Feathers of Young Ruddy Duck [Erismatura rubida]. Amer. Nat. xii, No. 2, February, 1878, pp. 123, 124, fig.
- 219. Justice to the English Sparrows [Passer domesticus]. *The Country*, Feb. 16, 1878.
- 220. †A Book on Bird-Architecture. The Country, March 16, 1878. Ernest Ingersoll's proposed treatise.
- 221. On the Moult of the Bill and Palpebral Ornaments in Fratercula arctica. Bull. Nutt. Orn. Club, iii, No. 2, April, 1878, pp. 87-91.
- 222. Habits of the Kingfisher [Ceryle aleyon]. Bull. Nutt. Orn. Club, iii, No. 2, April, 1878, p. 92.
- 223. Nesting of Vireo olivaceus. Bull. Nutt. Orn. Club, iii, No. 2, April, 1878, p. 95.
- 224. Nest and Eggs of Selasphorus platycercus. Bull. Nutt. Orn. Club, iii, No. 2, April, 1878, p. 95.
- 225. Meaning of the word "Anhinga." *Bull. Nut. Orn. Club*, iii, No. 2, April, 1878, p. 101.

- Letters on Ornithology. No. 18. The Yellow-breasted Chat. Chicago
   Field, June 29, 1878.
- 227. The Eave, Cliff, or Crescent Swallow (Petrochelidon lunifrons). Bull. Nutt. Orn. Club, iii, No. 3, July, 1878, pp. 105-112.
- 228. †Mr. H. Saunders on the Sterninæ. Bull. Nutt. Orn. Club, iii, No. 3, July, 1878, pp. 140-144.
  Review of the paper in the Proc. Zool. Soc. Lond, 1876, p. 638.
- 229. Swallow-tailed Kite [Elanoïdes forficatus] in Dakota in Winter. Bull. Nutt. Orn. Club, iii, No. 3, July, 1878, p. 147.
- 230. New Birds [five species] for the United States Fauna. The Country, July, 13, 1878, p. 184.
- 231. Field-Notes on Birds observed in Dakota and Montana along the Forty-ninth Parallel during the Seasons of 1873 and 1874. Bull. U. S. Geol. and Geogr. Surv. Terr., iv, No. 3, July 29, 1878, pp. 545-662.
  Also separate, new cover-title, same pagination, 8vo, Washington, 1878.
- 232. The Ineligibility of the European House Sparrow [Passer domesticus] in America. Amer. Nat., xii, No. 8, August, 1878, pp. 499–505.
  Reprinted, Chicago Field, Aug. 31, 1878; reprinted, The Country, Aug. 3, 1878; and elsewhere.
- 233. A Hint to Egg-Collectors. Bull. Nutt. Orn. Club, iii, No. 4, October, 1878, p. 191, cut.
- 234. Nest and Eggs of Helminthophaga pinus. Bull. Nutt. Orn. Club, iii, No. 4, October, 1878, p. 194.
- 235. †Wilson's and Bonaparte's American Ornithology. The Nation, Nov. 7, 1878.Review of the Porter and Coates' edition of 1878.

- 236. †Jones and Shulze's Illustrations of the Nest and Eggs of the Birds of Ohio.
   Bull. Nutt. Orn. Club, iv, No. 1, January, 1879, p. 52.
- 237. [Note on Dendrœca chrysoparia]. Bull. Nutt. Orn. Club, iv, No. 1, January, 1879, p. 60.
- 238. Nests and Eggs of the Clay-colored Bunting [Spizella pallida]. The Oölogist, iv, No. 7, February, 1879, p. 50.
- 239. Coues on the Nest and Eggs of the Water Thrush [Siurus nævius]. The Oölogist, iv, No. 8, March, 1879, p. 57.
- 240. The Sparrow [Passer domesticus] Nuisance. The Washington World, March 17, 1879.

- 241. History of the Evening Grosbeak [Hesperophona vespertina]. Bull. Nutt. Orn. Club, iv, No. 2, April, 1879, pp. 65-75.
- 242. †Langdon's Revised List of Cincinnati Birds. Bull. Nutt. Orn. Club, iv, No. 2, April, 1879, pp. 112, 113.
- 243. Note on Dendræca townsendi. Bull. Nutt. Orn. Club, iv, No. 2, April, 1879, p. 117.
- **244.** Note on Bucephala islandica. *Bull. Nutt. Orn. Club*, iv, No. 2, April, 1879, pp. 126, 127.
- 245. Letters on Ornithology. No. 19.—The Curlews [Numenius] of North America.— The Chicago Field, April, 26, 1879.
- 246. Letters on Ornithology. No. 20. The American Bittern [Botaurus mugitans]. Chicago Field, May 10, 1879.
- 247. Letters on Ornithology. No. 21. History of the Red-breasted, or Cinnamon Teal [Querquedula cyanoptera]. Chicago Field, May 17, 1879.
- 248. Letters on Ornithology. No. 22.—The Snow Goose, or White Brant [Chen hyperboreus].—Chicago Field, May 24, 1879.
- 249. Private Letters of Wilson, Ord and Bonaparte.—Penn Monthly, June, 1879, pp. 443-455.
- 250. Letters on Ornithology. No. 23. The American Coot [Fulica americana]. Chicago Field, June 9, 1879.
- 251. Letters on Ornithology. No. 24. The Wood Ibis [Tantalus loculator]. Chicago Field, June 14, 1879.
- 252. Letters on Ornithology. No. 25.—The Solitary Tattler; Wood Tattler [Rhyacophilus solitarius].— Chicago Field, June 21, 1879.
- 253. Letters on Ornithology. No. 26.—Semipalmated Tattler, Willet, Stone Snipe [Symphemia semipalmata].—Chicago Field, June 28, 1879.
- 254. To prevent Grease from injuring the Plumage of Birds. Amer. Nat., xiii, No. 7, July, 1879, p. 456.
- 255. [On the Use of Trinomials in Zoological Nomenclature]. Bull. Nutt. Orn. Club, iv, No. 3, July, 1879, p. 171.
- 256. Le Conte's Thrasher (Harporhynchus lecontii). The Oölogist, iv, No. 12, July, 1879, pp. 99–100.
- 257. Letters on Ornithology. No. 27. Bartramian Sandpiper or Tattler; Upland Plover [Bartramia longicauda]. Chicago Field, July 5, 1879.
- 258. Letters on Ornithology. No. 28. The Buff-breasted Sandpiper [Tryngites rufescens]. Chicago Field, July 12, 1879.

- 259. Letters on Ornithology. No. 29. Great Marbled Godwit [Limosa fœda].
   Chicago Field, July 19, 1879.
- 260. Letters on Ornithology. No. 30. The Great White Egret [Herodias egretta]. *Chicago Field*, July 26, 1879.
- 261. †Ingersoll's Nests and Eggs of American Birds. Amer. Nat., xiii, No. 8, Aug. 1879, pp. 515, 516.
- 262. †Texan Ornithology. Amer. Nat., xiii, No. 8, August, 1879, pp. 516-519.
- 263. Nesting of the Great Blue Heron [Ardea herodias] in the West. Chicago Field, Aug. 2, 1879.
- 264. On the Present Status of Passer domesticus in America, with Special Reference to the Western States and Territories. Bull. U. S. Geol. and Geogr. Surv. Terr., v, No. 2, Sept. 6, 1879, pp. 175-193.
  Including the bibliography of the subject. Also sep. pamphlet.
- 265. Second Instalment of American Ornithological Bibliography. Bull. U. S. Geol. and Geogr. Surv. Terr., v, No. 2, Sept. 6, 1879, pp. 239-330.
  The First Instalment forms the Appendix of "Birds of the Colorado Valley," Part I, 1878. Also sep. pamphlet, new cover-title, same pagination.
- 266. Note on the Black-capped Greenlet, Vireo atricapillus of Woodhouse. *Bull. Nutt. Orn. Club*, iv, No. 4, October, 1879, pp. 193, 194, pl. I (colored).
- 267. Southward Range of Centrophanes lapponica. Bull. Nutt. Orn. Club, iv, No. 4, October, 1879, p. 238.
- Obituary. [Miss Genevieve E. Jones.] Bull. Nutt. Orn. Club, iv, No. 4, October, 1879, p. 228.
- 269. A Correction [respecting Buteo borealis].—Bull. Nutt. Orn. Club, iv, No. 4, October, 1879, p. 242.
- Note on Alle nigricans, Link. Bull. Nutt. Orn. Club, iv, No. 4, October, 1879, p. 244.
- 271. Destructiveness of English Sparrows [Passer domesticus]. Amer. Nat., xiii, No. 11, November, 1879, p. 706.
- 272. Further Notes on the Ornithology of the Lower Rio Grande of Texas, from Observations made during the Spring of 1878. By George B. Sennett. Edited, with Annotations, by Dr. Elliott Coues, U. S. A. Bull. U. S. Geol. and Geogr. Surv. Terr., v, No. 3, Nov. 30, 1879, pp. 371-440.

  Also separate, new cover-title, same pagination.

273. The Origin of the Turkey [Meleagris gallipavo]. — Forest and Stream, xiii, No. 22, Jan. 1, 1880, p. 947.

- 274. Sketch of North American Ornithology in 1879. Amer. Nat., xiv, No. 1, January, 1880, pp. 20–25.
- On the Nesting in Missouri of Empidonax acadicus and Empidonax trailli.
   Bull. Nutt. Orn. Club, v, No. 1, January, 1880, pp. 20-25.
- 276. †Ingersoll's Nests and Eggs of American Birds. Bull. Nutt. Orn. Club, v, No. 1, January, 1880, pp. 38, 39.
- 277. †The Misses Jones and Shulze's Nests and Eggs of Ohio Birds. Bull. Nutt. Orn. Club, v, No. 1, January, 1880, pp. 39, 40.
- 278. Description of the Female Dendræca kirtlandi. Bull. Nutt. Orn. Club, v, No. 1, January, 1880, pp. 49, 50.
- 279. Note on Limosa hæmastica. *Bull. Nutt. Orn. Club*, v, No. 1, January, 1880, pp. 59, 60.
- 280. Capture of Phaëthon flavirostris in Western New York. Bull. Nutt. Orn. Club, v, No. 1, January, 1880, p. 63.
- 281. Depredations of the European Sparrow [Passer domesticus]. Amer. Nat., xiv, No. 2, February, 1880, p. 130.
- 282. Advent of Passer domesticus in North Carolina. Amer. Nat., xiv, No. 3, March, 1880, p. 213.
- 283. Notes and Queries concerning the Nomenclature of North American Birds.
   Bull. Nutt. Orn. Club, v, No. 2, April, 1880, pp. 95-102.
- 284. Further Light on the Moult of the Bill in certain Mormonidæ. Bull. Nutt. Orn. Club, v, No. 2, April, 1880, pp. 127-128.
- 285. Fourth Instalment of Ornithological Bibliography: being a List of Faunal Publications relating to British Birds. Proc. U. S. Nat. Mus., ii, May 31, 1880, pp. 359-482.
  This Instalment antedates the Third, below. Also separate, with new cover-title.
- 286. [Letters on Passer domesticus in America and Australia.] Forest and Stream, April 15, 1880, p. 204.
- 287. Shufeldt's Memoir on the Osteology of Speotyto cunicularia hypogæa. Bull. Nutt. Orn. Club, v, No. 3, July, 1880, pp. 129, 130, pll. I-III.
- 288. †Gentry's Nests and Eggs of the Birds of Pennsylvania. Bull. Nutt. Orn. Club, v, No. 3, July, 1880, p. 179.
- 289. †Ober's Camps in the Caribees. Bull. Nutt. Orn. Club, v, No. 3, July, 1880, p. 179.
- 290. Nest and Eggs of Catherpes mexicanus conspersus. Bull. Nutt. Orn. Club, v, No. 3, July, 1880, pp. 181, 182.
- 291. Number of Eggs of Ardea herodias. Bull. Nutt. Orn. Club, v, No. 3, July, 1880, p. 187.

- 292. Note on Grus fraterculus of Cassin. Bull. Nutt. Orn. Club, v, No. 3, July, 1880, p. 188.
- 293. Third Instalment of American Ornithological Bibliography. Bull. U. S. Geol. and Geogr. Surv. Terr., vi, No. 4, Sept. 30, 1880, pp. 521-1066.

  Not published till after the 4th, above. Not separate, occupying the whole No. of the Bull.
- 294. "Behind the Veil." Bull. Nutt. Orn. Club, v, No. 4, October, 1880, pp. 193–204.

Gossip over letters and other relics of Wilson and Audubon.

295. †Marsh's Palæornithology. — *Bull. Nutt. Orn. Club*, v, No. 4, October, 1880, pp. 234–236.

Review of O. C. Marsh's "Odontornithes."

296. Rural Bird Life | being | Essays on Ornithology | with instructions for preserving objects | relating to that science | by | Charles Dixon | with forty-five illustrations; and a preface | By Dr. Elliott Coues, U. S. A. . . . Boston | Estes and Lauriat | 299 to 305 Washington Street. | [1880.] 1 vol. sm. 8vo. Title and pp. i-xvi, 1-374, 45 illust.

American Editor's preface, pp. iii-viii.

- 297. A curious Colaptes [auratus × mexicanus]. Bull. Nutt. Orn. Club, vi, No. 3, July, 1881, p. 183.
- 298. A Correction [Trogonidæ]. Bull. Nutt. Orn. Club, vi, No. 3, July, 1881, p. 188.
- 299. Probable Occurrence of Sarcorhamphus papa in Arizona. Bull. Nutt. Orn. Club, vi, No. 4, October, 1881, p. 248.
- 300. New England Bird Life | being a | Manual | of | New England Ornithology | revised and edited from the Manuscript of | Winfrid A. Stearns | Member of the Nuttall Ornithological Club etc. | By | Dr. Elliott Coues U. S. A. | Member of the Academy etc. | Part I. Oscines | Boston | Lee and Shepard Publishers | New York Charles T. Dillingham | 1881 1 vol. 8vo. Title and pp. 1–324, figg. 1–56.

304- Brimophilies Kprysogeices per éco elove, \$65 - Magaziofenia o prégiés qual, o Kiby a hinol of onol. 432 - Beof blocus = Kéru-Dispar, & 20065 - Barte 296 - Sporgetila = Topos - sical \$2260 - Ilon

## VALUABLE BOOKS

# NATURAL HISTORY, GEOLOGY, &c.,

ATREDUCED PRICES.

FOR SALE BY

### ESTES & LAURIAT,

301-305 Washington Street, opposite Old South, Boston.

AUDUBON'S BIRDS OF AMERICA. By JOHN J. AUDUBON. 350 Figures of Birds, all life-size, in one vol., elephant folio. With descriptive text; 8 vols. bound in 4. Together, 5 vols., all bound uniform in heavy half Russia binding. New York, 1857. Reduced from \$250.00 to \$165.00. This grand work is now out of print and never can be produced again.

This magnificent work is in one elephant folio volume, measuring three feet five inches long by two feet three inches wide; containing upwards of three hundred and fifty beautifully colored figures of birds, portraying their action while in motion, all of which are life-size; together with a large number of illustrations of the plants and trees most frequented by the birds; a great variety of charming landscapes adding much to the beauty of the work. The descriptive text consists of eight volumes bound in four royal octavo volumes, giving the habits of the birds, anecdotes relating to them, by the great contribuleits who much this existence the study of this large and works life. eight volumes bound in four royal octavo volumes, giving the habits of the birds, aneedotes relating to them, by the great ornithologist who made this science the study of his long and useful life, with full descriptive text of all the birds, delineations of American scenery, plants, trees, &c.; also an account of many very exciting narratives of personal adventures, hair-breadth escapes, — being the experience of Mr. Audubon himself while in pursuit of the birds, — and it is, without exception, one of the most fascinating, valuable, entertaining, and remarkable works published. This set is bound in heavy Russia half binding, library style, and offered for sale at the above low price, — a very great bargain and a rare opportunity. If ordered from a distance, will be carefully packed in a case made expressly. The present offers a rare chance to lovers of works of art to procure, at a small price, one of the most costly and valuable works ever issued.

#### WILSON'S AMERICAN ORNITHOLO-

GY; or, Natural History of the Bir's of the United States. New and enlarged edition. 3 vols. 8vo. With portrait of Wilson, and 103 plates, exhibiting nearly 400 figures of birds, accurately engraved and beautifully printed in colors. New edition. London, 1877.

Reduced from \$25.00 to \$12.50.

This is by far the best edition of the American Ornithology, both on account of the beautiful plates, and the interesting notes of the editor; and every ornithologist should, if possible, procure this edition.

FISHES OF THE BRITISH ISLANDS (History of the). By Jonathan Couch, F.L.S. Illustrated with 256 carefully colored plates. 4 vols. Royal 8vo. Cloth. London, 1877.

Reduced from \$42.00 to \$23.00.

The author, who is well known as one of the first practical authorities on British fishes, has for fifty years been observing, noting, and drawing with his own pencil, the various fish which live in British waters, —a vast labor, in which he has been assisted by scientific friends living in various portions of the United Kingdom. The drawings are beautifully colored to life, and some of the portraits (especially of the dog-fish) are really marvellous, rendering the recognition of a fish a work of the greatest case. recognition of a fish a work of the greatest ease.

SMALLER BRITISH BIRDS. With descriptions of their Nests, Eggs, Habits, &c., &c. By H. G. and H. B. Adams. Illustrated with 30 beautiful full page colored plates. Containing hundreds of figures of birds and their eggs, well colored to nature. Small 4to, gilt edges. London, 1874. Reduced from \$12.00 to \$5.00.

The present work is an effort to produce a book which shall give a concise yet sufficiently full description of the smaller British birds; not a scientific work, but one essentially popular in its character, rendered attractive by lifelike portraits of the birds, drawn and colored as closely to nature as the eye and the hand of the artist can make them. BRITISH GRASSES (Natural History of). By E. J. Lowe, Esq. Illustrated with 74 finely colored plates. 8vo. Cloth. 1874. Reduced from \$10.00 to \$6.00.

This is a work not only valuable to the botanical student for its pictorial accuracy, but of use also to the landed proprietor and the farmer, pointing out to them those grasses which are useful and lucrative in husbandry, and teaching them the varied soils and positions upon which they thrive, and explaining their qualities and the several uses to which they are applied in many branches of manufacture and in-

RITISH AND EXOTIC FERNS (Natural History of). By E. J. Lowe, Esq. Illustrated with 479 finely colored plates. 8 vols. BRITISH Royal 8vo. Cloth. London, 1872.

Reduced from \$60.00 to \$32.00.

A book which should contain ample means of studying and identifying the Exotic species, accessible to persons of moderate means, has hitherto been a desideratum. This want the present work promises most hopefully to fill. It is admirably "got up;" the plates are carefully and prettily executed; there is a neat illustrative woodcut at the head of cuted; there is a neat illustrative woodcut at the head of each description, and the letterpress is full and practical, without being deficient in scientific accuracy. It is really the cheapest work for its excellence we have ever seen, and should be "in the hands of every gardener and every private person who cultivates these charming objects."

NEW AND RARE FERNS (Natural History of). Containing Species and Varieties not included in other works. By E. J. Lowe. Illustrated with 72 colored plates and numerous woodcuts. 8vo. Cloth. London, 1871.

Reduced from \$10.00 to \$6.00.

So many new ones have been introduced, that it has been deemed necessary to publish a separate volume. This work will be found to contain colored plates or woodcut illustrations of one hundred and fifty-one new species.

BRITISH MOSSES. Their Homes, Aspects, Structure, and Uses. Containing a Colored Figure of each species, etched from Nature. By F. E. Tripp. Illustrated with beautifully colored . plates. 2 vols. Royal 8vo. Cloth. London, 1874. Reduced from \$25.00 to \$13.00.

It is a book to read, to ponder, to mark, learn, and inwardly digest.... Let those who want to know the "moral" of mosses inquire within the covers of the volume. He will there find that these humble plants have their uses, their virtues, and their mission.

NATURAL HISTORY (Museum of): being a Popular Account of the Structure, Habits, and Classification of the various departments of the Animal Kingdom. By Sir John Richardson and Animal Amgdon. By Sh John Michaelson and others. With a History of the American Fauna, by Joseph B. Holder, M.D., Fellow of the New York Academy of Sciences. Illustrated with hundreds of steel engravings, also many finely colored with the Cloth ored plates, and numerous woodcuts. 4to. Cloth, gilt, extra. New York, 1880.

Reduced from \$24.00 to \$10.50.

Not only written in a free, familiar, teaching style, but exquisitely illustrated and beautifully got up. Gives a large amount of information on the subject of animal life, and we commend these handsome volumes very heartily to all interested in natural history.

STRUTT'S SYLVA BRITANNICA AND SCOTICA; or, portraits of Forest Trees distinguished for their antiquity, magni-tude, or beauty. Drawn from Nature, and etched by Jacob George Strutt. Imperial folio. Com-prising 50 very large and highly finished etchings. Half bound morocco, extra, gilt edges. London, Reduced from \$45.00 to \$18.00. 1826.

Some of the etchings resemble the paintings by Waterloo very strikingly; but the whole are various, beautiful, and interesting.

BUTTERFLIES AND MOTHS (History of). Accurately delineating every known Species, with the English as well as the Scientific Names, accompanied by full Descriptions, Date of Appearance, Lists of the Localities they haunt, their Food in the Caterpillar State, and other Features of their Habits and Modes of Existence, &c. By Rev. F. O. Morris, B.A. The plates contain nearly 2,000 exquisitely colored specimens. In 4 vols. Royal 8vo. Cloth. London. Pachysed from \$60 Otto \$32.00 Reduced from \$60.00 to \$32.00.

Speaking of entomology, we should place Mr. Morris's "History of British Moths" at the head. It gives a colored figure of every known British moth, together with dates appearance, localities, description, and food of caterpillar. It forms a handsome work for a library, and will, we should be a local description of the property of th hope, lead many to commence the fascinating study of entomology.

OUR NATIVE FERNS; or, a History of the British Species and their Varieties. By E. J. Lowe. With 79 colored and 909 wood engravings. 2 vols. 8vo. Cloth. London, 1874. Reduced from \$20.00 to \$12.00.

The importance and value of this work may be inferred from the fact that it contains descriptions of 1,294 varieties of British ferns, with 79 colored plates of species and varieties, and 909 wood engravings, containing much interesting information. The localities are described, each synonyme given, and a description of the proper method of cultivation. There are 184 varieties figured.

BRITISH SEA-WEEDS. Drawn from Professor Harvey's "Phycologia Britannica." With Descriptions, and Amateur's Synopsis, Rules for Laying on Sea-weeds, an Order for Arranging them in the Herbarium, and an Appendix of New Species. By Mrs. Alfred Gatty. Illus-trated with 80 exquisitely colored plates, contain-ing 381 figures. 2 vols. 8vo. Cloth. London, 1874. Reduced from \$24.00 to \$13.00.

In her present work she has endeavored, and we think most successfuly, to translate the terms and phrases of science into the language of amateurs. Mrs. Gatty's familiarity with the plants themselves has enabled her to do this office without failing into the errors to which a mere compiler in separating from the beaten track would be liable.

HISTORY OF BRITISH BUTTER-FLIES. By the Rev. F. O. Morris, B.A. Illustrated with 72 beautifully colored plates. Royal 8vo. Cloth. London, 1876.

Reduced from \$10.00 to \$6.00.

With colored illustrations of all the species, and separate figures of the male and female, where there is any obvious difference between them, and also of the under side, together with the caterpillar and chrysalis, and a full description of each, with copious accounts of their several habits, localities, and times of appearance, together with details as to their preservation, &c., with new and valuable information, — the result of the author's experience for many years.

ALPINE PLANTS. Descriptions and 200 accurately colored figures (drawn and engraved expressly for this work) of some of the most striking and beautiful of the Alpine Flowers. Edited by David Wooster. 2 vols. Royal 8vo. Cloth. London. Reduced from \$20.00 to \$13.00.

The manner in which "Alpine Plants" is produced is creditable alike to author and artist. The literary portion is not the mere dry botanical description often found in such works, but a popular description of the plant, - instructions as to its culture and treatment, — with any interesting information in connection with it that can be obtained. . . . We heartily commend this work to all lovers of flowers.

OWEN (Richard). Comparative Anatomy and Physiology of Vertebrates. Vol. I. Fishes and Reptiles. Vol. II. Birds and Mammals. Vol. III. Mammals. 3 vols. 8vo. Cloth. Illus-III. Mammals. 3 vols. 8vo. Cloth. Illustrated with an immense number of beautiful woodcuts. London, Longmans, 1866-68. Reduced from \$24.00 to \$10.50.

This work has long been the highest authority on this subject, and has received the pruise of such men as Tyndall, Huxley, and others. Humboldt speaks of Owen as the greatest anatomist of his age, and he is generally called the Cuvier of England and the "Newton of natural history."

AMERICAN NATURALIST (The). popular illustrated Magazine of Natural History, Edited by A. S. Packard, Jr., E. S. Morse, A. Hyatt, and F. W. Putnam. Numerous illustrations, many full-paged. 10 vols. (Complete from commencement to 1878.) 8vo. Cloth. Salem, 1868-77. Reduced from \$50.00 to \$25.00. The Naturalist contains departments of Geography and

Travel, Microscopy, and Proceedings of Scientific Societies. A digest of the contents of foreign scientific journals and transactions is also given, together with the latest home and foreign scientific news.

GEOLOGY OF NEW HAMPSHIRE. A report comprising the results of Exploration or-dered by the Legislature, by C. H. Hitchcock and J. H. Huntington. Illustrated with nearly 250 illustrations, maps, diagrams (many of which are full-page). 3 vols. Royal 8vo. 1,200 pages. Half morocco. With large and valuable Atlas completing the work, forming the fourth volume. Concord, 1874. Reduced from \$40.00 to \$25.00.

This great work, which is without a doubt the most valuable report ever published, contains many articles of interest to the general reader, as well as the geologist, as many of the articles were prepared by gentlemen eminent in their specialties, such as the Natural History and Botany, and a History of the Explorations among the White Mountains, &c.


ATURALIST'S LIBRARY, JAR-DINE'S. By Sir William Jardine. 42 vols. Foolscap. 8vo. 1,200 colored plates. With numerous Portraits and Memoirs of Eminent NATURALIST'S LIBRARY, Naturalists. Extra cloth, top edges, gilt. (Sold only in sets.) London, 1865.

Reduced from \$84.00 to \$36.00. Contents: British Birds; Sun Birds; Humming Birds; Contents: British Brids; Sun Brids; Humming Brids; Game Birds; Pigeons; Parrots; Flycatchers; Peacocks; Lions; Tigers; British Quadrupeds; Dogs, 2 vols; Rumi-nating Animals, vol. 1 (Deer, Antelopes, &c.); Ruminating Animals, vol. 2 (Goats, Sheep); Seals; Whales, &c.; Mon-keys; British Butterflies; British Moths, &c.; Foreign Butterflies; Foreign Moths; Beetles; Bees; Introduction, and Foreign Fishes; British Fishes, 2 vols.; Perch, &c;

and Foreign Fishes; British Fishes, 2 vois.; Perch, &c; Fishes of Guiana, 2 vols.

Sir William Jardine's coadjutors in this admirable series were Swainson, Selby, Macgillivray, Waterhouse, Duncan, Hamilton, Smith, and others.

This book is, perhaps, the most interesting, the most beautiful, and the cheapest series ever offered to the public.











3 9088 00001 5677